

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 August 2002 (22.08.2002)

PCT

(10) International Publication Number
WO 02/064550 A1

(51) International Patent Classification⁷: C07C 307/04,
311/08, A61K 31/18, A61P 3/10

(21) International Application Number: PCT/US02/04501

(22) International Filing Date: 12 February 2002 (12.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/783,636 14 February 2001 (14.02.2001) US
Not furnished 8 February 2002 (08.02.2002) US

(71) Applicant: ABBOTT LABORATORIES [US/US];
Dept. 377/AP6D-2, 100 Abbott Park Road, Abbott Park,
IL 60064-6050 (US).

(72) Inventors: LINK, James, T.; Apt. 1405, 1630 Chicago
Avenue, Chicago, IL 60201 (US). SORESENSEN, Bryan,
K.; 2620 N. Lewis Avenue, Waukegan, IL 60087 (US).

PATEL, Jyoti, R.; 1408 Newgate Court, Libertyville,
IL 60048 (US). ARENSEN, David, L.; 431 E. Austin
Avenue, Libertyville, IL 60048 (US). LI, Gaoquan; 793
Sharon Avenue, Park City, IL 60085 (US).

(74) Agents: COLLINS, Daniel, W. et al.; Dept. 377/AP6D-2,
100 Abbott Park Road, Abbott Park, IL 60064-6050 (US).

(81) Designated States (national): CA, JP, MX.


(84) Designated States (regional): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: GLUCOCORTICOID RECEPTOR MODULATORS



(57) Abstract: Compounds of formula (I) (I), or pharmaceutically acceptable salts thereof are novel glucocorticoid receptor modulators and are useful for treating type II diabetes in a mammal.



WO 02/064550 A1

GLUCOCORTICOID RECEPTOR MODULATORS

5

FIELD OF THE INVENTION

The present invention relates to non-steroidal compounds which are selective modulators of steroid receptors, in particular, the glucocorticoid receptor. In particular, the present invention is directed to compounds and methods for using such compounds to treat mammals requiring glucocorticoid receptor modulation.

10

BACKGROUND OF THE INVENTION

15

20

25

Glucocorticoids are lipid soluble hormones synthesized in the adrenal cortex (Neville and O'Hare, *The Adrenal Gland*, James, Ed. New York, Raven, 1-65, (1979)). These molecules readily pass through cell membranes and enter the cytoplasm of target tissues, where they bind to glucocorticoid receptors sequestered in the cytoplasm by complexation with heat shock proteins. Upon binding of the hormone to its receptor, the receptor undergoes a conformational change which results in dissociation of heat shock proteins, and translocation of the ligand bound glucocorticoid receptor into the nucleus where it can either initiate or repress specific gene transcription. Transcriptional activation occurs when the ligand bound receptor complex homodimerizes, and the homodimeric receptor ligand complex binds to chromosomal deoxyribonucleic acid (DNA) at sequence specific sites in the promoter region of regulated genes (Beato, *Cell*, 56, 335-344 (1989); and Yamamoto, *Annu. Rev. Genet.*, 19, 209-215 (1989)). Excesses or deficiencies of these ligands can have profound physiological consequences. As an example, glucocorticoid excess results in Cushing's Syndrome, while glucocorticoid insufficiency results in Addison's Disease.

30

Biologically relevant glucocorticoid receptor agonists include cortisol and corticosterone. Many synthetic glucocorticoid receptor agonists exist including dexamethasone, prednisone and prednisilone. By definition, glucocorticoid receptor antagonists bind to the receptor and prevent glucocorticoid receptor agonists from binding and eliciting GR mediated events, including transcription. RU486 is an example of a non-selective glucocorticoid receptor antagonist.

35

Type II diabetes (also referred to as non insulin-dependent Diabetes Mellitus) is a debilitating disease characterized by an abnormal elevation of blood glucose levels driven by three factors: increased hepatic glucose production, inadequate clearance of glucose via insulin mediated pathways, and decreased uptake of circulating glucose by tissues (DeFronzo, *Diabetes Review* 5(3), 177-269, (1997)). Administration of agents that decrease hepatic glucose production are a fundamental approach to controlling blood glucose (De

Feo et al., *Am. J. Physiol.* 257, E35-E42 (1989); Rooney, et al., *J. Clin. Endocrinol. Metab.* 77, 1180-1183 (1994); and Dinneen et al., *J. Clin. Invest.*, 92, 2283-2290 (1993)).

Glucocorticoids have been shown to have major influences on glucose production.

Glucocorticoid excess aggravates established diabetes while glucocorticoid deficiency
5 reduces blood glucose and improves glucose control in diabetic mice (Boyle, *Diabetes Review*, 1(3), 301-308, (1993); Naeser, *Diabetologia*, 9, 376-379 (1973); and Solomon et al., *Horm. Metab. Res.*, 9, 152-156 (1977)). The underlying mechanism responsible for these effects is the glucocorticoid-induced upregulation of key hepatic enzymes required for gluconeogenesis (Exton et al., *Recent Prog. Horm. Res.*, 26, 411-457 (1970); and Kraus-
10 Friedmann, *Physiol. Rev.*, 64, 170-259 (1984)).

Among the genes that glucocorticoids up-regulate are several genes that play key roles in gluconeogenesis and glycogenolysis, particularly phosphoenolpyruvyl carboxykinase (PEPCK) and glucose-6-phosphatase (Hanson and Patel, *Adv. Enzymol.*, Meister, Ed. New York, John Wiley and Sons, Inc., 203-281 (1994); and Argaud et al.,
15 *Diabetes* 45, 1563-1571 (1996)). Mifepristone (RU486), a potent GR antagonist reduces messenger ribonucleic acid (mRNA), levels of PEPCK and glucose-6-phosphate in the liver, and causes a 50% reduction of plasma glucose levels in obese diabetic db/db transgenic mice (Friedman et al., *J. Biol. Chem.* 272(50), 31475-31481 (1997)). While steroid-based GR antagonists have been useful in demonstrating efficacy for in vivo glucose lowering
20 effects, the utility of such agents is limited due to side effects resulting from potent cross-reactivity with other steroid receptors, in particular progesterone receptor (PR) and mineralocorticoid receptor (MR). US 5,929,058 discloses a method for treating type II diabetes by administering a combination of steroidal-agents that exhibit mineralcorticoid receptor agonist activity and glucocorticoid receptor antagonist activity. Pharmaceutical
25 agents that function as glucocorticoid receptor (GR) antagonists represent a novel approach to controlling type II diabetes.

Other examples of proteins that interact with the glucocorticoid receptor are the transcription factors AP-1 and NF κ -B. Such interactions result in inhibition of AP-1- and NF κ -B- mediated transcription and are believed to be responsible for some of the anti-
30 inflammatory activity of endogenously administered glucocorticoids. In addition, glucocorticoids may also exert physiologic effects independent of nuclear transcription.

Glucocorticoid selective non-steroidal agents that antagonize functional activity mediated by the glucocorticoid receptor may be useful for treating mammals suffering from type II diabetes and for treating symptoms of type II diabetes including hyperglycemia,
35 inadequate glucose clearance, obesity, hyperinsulinemia, hypertriglyceridemia and high circulating glucocorticoid levels. In addition to diabetes, glucocorticoid receptor modulators are useful to treat diseases such as obesity, Syndrome X, Cushing's Syndrome, Addison's disease, inflammatory diseases such as asthma, rhinitis and arthritis, allergy,


autoimmune disease, immunodeficiency, anorexia, cachexia, bone loss or bone frailty, and wound healing.

US Patent No. 5,929,058 discloses a method for treating type II diabetes by administering a combination of steroidal-agents that exhibit mineralcorticoid receptor agonist activity and glucocorticoid receptor antagonist activity.

Although there are glucocorticoid receptor therapies in the art, there is a continuing need for and a continuing search in this field of art for selective glucocorticoid receptor therapies. Thus, the identification of non-steroidal compounds which have specificity for one or more steroid receptors, but which have reduced or no cross-reactivity for other steroid or intracellular receptors, is of significant value in this field.

SUMMARY OF THE INVENTION

In its principle embodiment, the present invention discloses compounds of formula (I) that are useful as glucocorticoid receptor modulators:



or a pharmaceutically acceptable salt or prodrug thereof, wherein

L_A is a covalent bond or $C(X_{A1})(X_{A2})$;

L_D is a covalent bond or $C(X_{D1})(X_{D2})$;

X_{A1} is selected from the group consisting of hydrogen, alkoxy, alkoxycarbonyl, alkoxyalkyl, alkyl, alkylcarbonyl, cyano, halogen, hydroxy, hydroxyalkyl, nitro and $(NR_{10}R_{11})$ carbonyl;

X_{A2} is selected from the group consisting of hydrogen and alkyl;

or X_{A1} and X_{A2} together are oxo;

X_{D1} is selected from the group consisting of hydrogen, alkoxy, alkoxycarbonyl, alkoxyalkyl, alkyl, alkylcarbonyl, cyano, halogen, hydroxy, hydroxyalkyl, nitro and $(NR_{10}R_{11})$ carbonyl;

X_{D2} is selected from the group consisting of hydrogen and alkyl;

or X_{D1} and X_{D2} together are oxo;

L_B is absent or selected from the group consisting of a covalent bond, alkenylene, alkylene, alkynylene, $-(CH_2)_mC(O)(CH_2)_n-$, $-(CH_2)_mC(O)(CH_2)_nCH=CH-$, $-(CH_2)_mC(O)(CH_2)_nC\equiv C-$, $-(CH_2)_mCH(OH)(CH_2)_n-$, $-(CH_2)_mCH(OH)(CH_2)_nCH_2CH=CH-$, $-(CH_2)_mCH(OH)(CH_2)_nCH_2C\equiv C-$, $-(CH_2)_mO(CH_2)_n-$, $-(CH_2)_mO(CH_2)_nCH_2CH=CH-$,
 5 $-(CH_2)_mO(CH_2)_nCH_2C\equiv C-$, $-(CH_2)_mS(CH_2)_n-$, $-(CH_2)_mS(O)(CH_2)_n-$, $-(CH_2)_mS(O)_2(CH_2)_n-$, $-(CH_2)_mN(R_7)(CH_2)_n-$, $-(CH_2)_mN(R_7)C(O)(CH_2)_n-$, $-(CH_2)_mC(O)N(R_7)(CH_2)_n-$, $-O(CH_2)_mC(O)N(R_7)(CH_2)_n-$, $-(CH_2)_mC(O)O(CH_2)_n-$, $-O(CH_2)_mC(O)(CH_2)_n-$, $-(CH_2)_mN(R_7)S(O)_2(CH_2)_n-$, $-(CH_2)_mS(O)_2N(R_7)(CH_2)_n-$, $-(CH_2)_mS(O)_2O(CH_2)_n-$ and

10 $-(CH_2)_mOS(O)_2(CH_2)_n-$ wherein each group is inserted as drawn with the left end attached to A and the right end attached to B;

m is an integer of 0-6;

n is an integer of 0-6;

L_C is absent or selected from the group consisting of a covalent bond, alkenylene, alkylene, alkynylene, $-(CH_2)_pC(O)(CH_2)_q-$, $-(CH_2)_pC(O)(CH_2)_qCH=CH-$, $-(CH_2)_pC(O)(CH_2)_qC\equiv C-$, $-(CH_2)_pCH(OH)(CH_2)_q-$, $-(CH_2)_pCH(OH)(CH_2)_qCH_2CH=CH-$, $-(CH_2)_pCH(OH)(CH_2)_qCH_2C\equiv C-$, $-(CH_2)_pO(CH_2)_q-$, $-(CH_2)_pO(CH_2)_qCH_2CH=CH-$, $-(CH_2)_pO(CH_2)_qCH_2C\equiv C-$, $-(CH_2)_pS(CH_2)_q-$, $-(CH_2)_pS(O)(CH_2)_q-$, $-(CH_2)_pS(O)_2(CH_2)_q-$, $-(CH_2)_pN(R_7)(CH_2)_q-$, $-(CH_2)_pN(R_7)C(O)(CH_2)_q-$, $-(CH_2)_pC(O)N(R_7)(CH_2)_q-$,
 15 $-O(CH_2)_pC(O)N(R_7)(CH_2)_q-$, $-(CH_2)_pC(O)O(CH_2)_q-$, $-O(CH_2)_pC(O)(CH_2)_q-$, $-(CH_2)_pN(R_7)S(O)_2(CH_2)_q-$, $-(CH_2)_pS(O)_2N(R_7)(CH_2)_q-$, $-(CH_2)_pS(O)_2O(CH_2)_q-$ and $-(CH_2)_pOS(O)_2(CH_2)_q-$ wherein each group is inserted as drawn with the left end attached to B and the right end attached to C;

p is an integer of 0-6;

25 q is an integer of 0-6;

L_E is absent or selected from the group consisting of a covalent bond, alkenylene, alkylene, alkynylene, $-(CH_2)_rC(O)(CH_2)_s-$, $-(CH_2)_rC(O)(CH_2)_sCH=CH-$, $-(CH_2)_rC(O)(CH_2)_sC\equiv C-$, $-(CH_2)_rCH(OH)(CH_2)_s-$, $-(CH_2)_rCH(OH)(CH_2)_sCH_2CH=CH-$, $-(CH_2)_rCH(OH)(CH_2)_sCH_2C\equiv C-$, $-(CH_2)_rO(CH_2)_s-$, $-(CH_2)_rO(CH_2)_sCH_2CH=CH-$,
 30 $-(CH_2)_rO(CH_2)_sCH_2C\equiv C-$, $-(CH_2)_rS(CH_2)_s-$, $-(CH_2)_rS(O)(CH_2)_s-$, $-(CH_2)_rS(O)_2(CH_2)_s-$, $-(CH_2)_rN(R_7)(CH_2)_s-$, $-(CH_2)_rN(R_7)C(O)(CH_2)_s-$, $-(CH_2)_rC(O)N(R_7)(CH_2)_s-$, $-O(CH_2)_rC(O)N(R_7)(CH_2)_s-$, $-(CH_2)_rC(O)O(CH_2)_s-$, $-O(CH_2)_rC(O)(CH_2)_s-$, $-(CH_2)_rN(R_7)S(O)_2(CH_2)_s-$, $-(CH_2)_rS(O)_2N(R_7)(CH_2)_s-$, $-(CH_2)_rS(O)_2O(CH_2)_s-$ and
 35 $-(CH_2)_rOS(O)_2(CH_2)_s-$ wherein each group is inserted as drawn with the left end attached to D and the right end attached to E;

r is an integer of 0-6;

s is an integer of 0-6;

- L_F is absent or selected from the group consisting of a covalent bond, alkenylene, alkylene, alkynylene, $-(CH_2)_uC(O)(CH_2)_v-$, $-(CH_2)_uC(O)(CH_2)_vCH=CH-$, $-(CH_2)_uC(O)(CH_2)_vC\equiv C-$, $-(CH_2)_uCH(OH)(CH_2)_v-$, $-(CH_2)_uCH(OH)(CH_2)_vCH_2CH=CH-$, $-(CH_2)_uCH(OH)(CH_2)_vCH_2C\equiv C-$, $-(CH_2)_uO(CH_2)_v-$, $-(CH_2)_uO(CH_2)_vCH_2CH=CH-$, $-(CH_2)_uO(CH_2)_vCH_2C\equiv C-$, $-(CH_2)_uS(CH_2)_v-$, $-(CH_2)_uS(O)(CH_2)_v-$, $-(CH_2)_uS(O)_2(CH_2)_v-$, $-(CH_2)_uN(R_7)(CH_2)_v-$, $-(CH_2)_uN(R_7)C(O)(CH_2)_v-$, $-(CH_2)_uC(O)N(R_7)(CH_2)_v-$, $-O(CH_2)_uC(O)N(R_7)(CH_2)_v-$, $-(CH_2)_uC(O)O(CH_2)_v-$, $-O(CH_2)_uC(O)(CH_2)_v-$, $-(CH_2)_uN(R_7)S(O)_2(CH_2)_v-$, $-(CH_2)_uS(O)_2N(R_7)(CH_2)_v-$, $-(CH_2)_uS(O)_2O(CH_2)_v-$ and $-(CH_2)_uOS(O)_2(CH_2)_v-$ wherein each group is inserted as drawn with the left end attached to E and the right end attached to F;

u is an integer of 0-6;

v is an integer of 0-6;

- A and D are each independently selected from the group consisting of aryl, cycloalkyl and heterocycle;

B, C, E and F are each independently absent or each independently selected from the group consisting of aryl, cycloalkyl and heterocycle;

- R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} and R_{F5} are each independently absent or each independently selected from the group consisting of hydrogen, alkenyl, alkenylthio, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxyacarbonyl, alkoxyacarbonylalkoxy, alkoxyacarbonylalkyl, alkoxyacarbonylalkylthio, alkyl, alkylcarbonyl, alkylcarbonylalkoxy, alkylcarbonylalkyl, alkylcarbonylalkylthio, alkylcarbonyl(NR₁₁)sulfonylalkyl, alkylcarbonyloxy, alkylcarbonylthio, alkylsulfinyl, alkylsulfinylalkyl, alkylsulfonyl, alkylsulfonylalkyl, alkylsulfonyl(NR₁₀)carboxylalkyl, alkylsulfonyl(NR₁₀)carboxylalkoxy, alkylsulfonyl(NR₁₀)alkyl(NR₁₁)-, alkylthio, alkylthioalkyl, alkylthioalkoxy, alkynyl, alkynyloxy, alkynylthio, carboxy, carboxyalkoxy, carboxylalkoxyalkoxy, carboxyalkyl, cyano, cyanoalkoxy, cyanoalkyl, cyanoalkylthio, cycloalkylalkoxy, cycloalkyloxy, ethylenedioxy, formyl, formylalkoxy, formylalkyl, haloalkenyl, haloalkenyloxy, haloalkoxy, haloalkyl, haloalkynyl, haloalkynyloxy, halogen, hydroxy, hydroxyalkoxy, hydroxyalkyl, mercapto, mercaptoalkoxy, mercaptoalkyl, methylenedioxy, nitro, -NR₈R₉, (NR₈R₉)alkoxy, (NR₈R₉)alkyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkoxy, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl, (NR₁₀R₁₁)sulfonylalkyl, -NR₁₀S(O)₂R₁₂, -NR₁₀S(O)₂NR₁₃R₁₄, and -S(O)₂OH;

R_1 , R_2 , and R_3 are each independently selected from the group consisting of hydrogen, alkoxyacarbonyl, alkoxy, alkoxyalkyl, alkyl, alkylcarbonyl, carboxy, halogen, hydroxyalkyl, -NR₁₀R₁₁ and (NR₁₀R₁₁)alkyl;

R_4 is selected from the group consisting of hydrogen, alkenyl, alkoxy, alkoxyalkenyl, alkoxyalkoxy, alkoxyalkyl, alkoxyalkynyl, alkoxycarbonyl, alkoxycarbonylalkoxy, alkoxycarbonylalkenyl, alkoxycarbonylalkyl, alkoxycarbonylalkynyl, alkyl, alkylcarbonyl, alkylcarbonylalkenyl, alkylcarbonylalkoxy, alkylcarbonylalkyl, alkylcarbonylalkynyl, alkynyl, carboxy, carboxyalkenyl, carboxyalkyl, carboxyalkynyl, haloalkoxy, haloalkyl, haloalkenyl, haloalkynyl, halogen, hydroxyalkyl, - $NR_{10}R_{11}$, $(NR_{10}R_{11})$ alkyl, $(NR_{10}R_{11})$ carbonyl, $(NR_{10}R_{11})$ carbonylalkyl, $(NR_{10}R_{11})$ carbonylalkenyl and $(NR_{10}R_{11})$ carbonylalkynyl;

R_5 is selected from the group consisting of hydrogen and alkyl; and

R_6 is selected from the group consisting of hydrogen, alkoxycarbonyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylsulfonyl, arylalkoxycarbonyl, arylalkylcarbonyl, arylalkylsulfonyl, arylcarbonyl, arylsulfonyl, cycloalkylcarbonyl, cycloalkylalkylcarbonyl, cycloalkylsulfonyl, cycloalkylalkylsulfonyl, heterocyclecarbonyl, heterocyclealkylcarbonyl, heterocyclesulfonyl, heterocyclealkylsulfonyl, $(NR_{13}R_{14})$ carbonyl and $(NR_{13}R_{14})$ sulfonyl;

R_8 and R_9 are each independently selected from the group consisting of hydrogen, alkenyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkylcarbonyl, alkylsulfonyl, alkynyl, carboxyalkylcarbonyl; cyanoalkyl, formyl, hydroxy, hydroxyalkyl, - $NR_{10}R_{11}$, $(NR_{10}R_{11})$ carbonyl and carboxyalkyl wherein the alkyl portion of carboxyalkyl is optionally substituted with one or two substituents selected from the group consisting of alkylthio, aryl, heterocycle, hydroxy, carboxy, - $NR_{10}R_{11}$ and $(NR_{10}R_{11})$ carboxy;

R_{10} and R_{11} are each independently selected from the group consisting of hydrogen and alkyl;

R_{12} is selected from the group consisting of alkoxy, alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl; and

R_{13} and R_{14} are each independently selected from the group consisting of hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl.

In another embodiment, compounds of the present invention have formula (I) wherein L_A is a covalent bond; L_D is a covalent bond; A is aryl; D is aryl; and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} , R_{F5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , L_E , L_F , B, C, E, and F are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (I) wherein L_A is a covalent bond; L_D is a covalent bond; A is heterocycle; D is aryl; and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} , R_{F5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , L_E , L_F , B, C, E, and F are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (I) wherein L_A is a covalent bond; L_D is a covalent bond; A is heterocycle; D is heterocycle;

In another embodiment, compounds of the present invention have formula (I)

5 wherein L_A is $C(X_{A1})(X_{A2})$; L_D is a covalent bond; A is aryl; D is aryl; and X_{A1} , X_{A2} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} , R_{F5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , L_E , L_F , B, C, E, and F are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (I) wherein L_A is $C(X_{A1})(X_{A2})$; L_D is a covalent bond; A is heterocycle; D is heterocycle; and X_{A1} , X_{A2} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} , R_{F5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , L_E , L_F , B, C, E, and F are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (I) wherein L_A is $C(X_{A1})(X_{A2})$; L_D is $C(X_{D1})(X_{D2})$; A is heterocycle; D is heterocycle; and X_{A1} , X_{A2} , X_{D1} , X_{D2} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} , R_{F5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , L_E , L_F , B, C, E, and F are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (I)
 35 wherein L_A is $C(X_{A1})(X_{A2})$; L_D is $C(X_{D1})(X_{D2})$; A is aryl; D is aryl; and X_{A1} , X_{A2} , X_{D1} , X_{D2} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{E1} , R_{E2} , R_{E3} , R_{E4} , R_{E5} , R_{F1} , R_{F2} , R_{F3} , R_{F4} , R_{F5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , L_E , L_F , B, C, E, and F are as defined in formula (I).

DETAILED DESCRIPTION OF THE INVENTION

Definition of Terms

As used throughout this specification and the appended claims, the following terms
5 have the following meanings.

The term, "modulator," as used herein, refers to a chemical entity, whether a synthesized chemical entity or a natural endogenous chemical entity, that interacts with a steroid receptor, wherein said interaction elicits a response by the receptor or blocks the response of the receptor when interacted with by a natural endogenous chemical entity, i.e.,
10 cortisol or corticosterone.

The term "alkenyl," as used herein, refers to a straight or branched chain hydrocarbon containing from 2 to 10 carbons and containing at least one carbon-carbon double bond formed by the removal of two hydrogens. Representative examples of alkenyl include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl.
15

The term "alkenyloxy," as used herein, refers to an alkenyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkenyloxy include, but are not limited to, allyloxy, 2-butenyloxy, 3-butenyloxy and 3-pentenlyloxy.
20

The term "alkenylthio," as used herein, refers to an alkenyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom. Representative examples of alkenylthio include, but are not limited, allylsulfanyl, 2-butenylsulfanyl, 3-butenylsulfanyl and 3-pentenylsulfanyl.

The term "alkenylene," denotes a divalent group derived from a straight or branched chain hydrocarbon of from 2 to 10 carbon atoms containing at least one double bond. Representative examples of alkenylene include, but are not limited to, -CH=CH-, -CH=CH₂CH₂-, -CH₂-CH=CH-, -CH₂CH₂CH=CH- and -CH=C(CH₃)CH₂-.
25

The term "alkoxy," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy, tert-butoxy, pentyloxy, and hexyloxy.
30

The term "alkoxyalkenyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined herein. Representative examples of alkoxyalkenyl include, but are not limited to, 3-methoxy-1-propenyl and 4-methoxy-1-butenyl.
35

The term "alkoxyalkoxy," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through another alkoxy group, as defined

herein. Representative examples of alkoxyalkoxy include, but are not limited to, tert-butoxymethoxy, 2-ethoxyethoxy, 2-methoxyethoxy, and methoxymethoxy.

5 The term "alkoxyalkoxyalkoxy," as used herein, refers to an alkoxyalkoxy group, as defined herein, appended to the parent molecular moiety through another alkoxy group, as defined herein. Representative examples of alkoxyalkoxyalkoxy include, but are not limited to, (2-methoxyethoxy)methoxy and (2-methoxyethoxy)ethoxy.

10 The term "alkoxyalkoxyalkyl," as used herein, refers to an alkoxyalkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkoxyalkoxyalkyl include, but are not limited to, tert-butoxymethoxymethyl, ethoxymethoxymethyl, (2-methoxyethoxy)methyl, and 2-(2-methoxyethoxy)ethyl.

The term "alkoxyalkyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkoxyalkyl include, but are not limited to, tert-butoxymethyl, 2-ethoxyethyl, 2-methoxyethyl, and methoxymethyl.

15 The term "alkoxyalkynyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkynyl group, as defined herein. Representative examples of alkoxyalkynyl include, but are not limited to, 3-ethoxy-1-propynyl and 4-methoxy-1-butyne.

20 The term "alkoxycarbonyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkoxycarbonyl include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, and tert-butoxycarbonyl.

25 The term "alkoxycarbonylalkenyl," as used herein, refers to an alkoxycarbonyl group, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined herein. Representative examples of alkoxycarbonylalkenyl include, but are not limited to, 3-methoxy-3-oxo-1-propenyl and 3-ethoxy-3-oxo-1-propenyl.

30 The term "alkoxycarbonylalkoxy," as used herein, refers to an alkoxycarbonyl group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of alkoxycarbonylalkoxy include, but are not limited to, 3-methoxy-3-oxopropoxy, 3-ethoxy-3-oxopropoxy, 2-methoxy-2-oxoethoxy and 4-methoxy-4-oxobutoxy.

35 The term "alkoxycarbonylalkyl," as used herein, refers to an alkoxycarbonyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkoxycarbonylalkyl include, but are not limited to, 3-methoxycarbonylpropyl, 4-ethoxycarbonylbutyl, and 2-tert-butoxycarbonylethyl.

The term "alkoxycarbonylalkynyl," as used herein, refers to an alkoxycarbonyl group, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined herein. Representative examples of alkoxycarbonylalkynyl include, but are not limited to, 3-methoxy-3-oxo-1-propynyl and 3-ethoxy-3-oxo-1-propynyl.

5 The term "alkoxysulfonyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of alkoxysulfonyl include, but are not limited to, methoxysulfonyl, ethoxysulfonyl and propoxysulfonyl.

10 The term "alkyl," as used herein, refers to a straight or branched chain hydrocarbon containing from 1 to 10 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, n-heptyl, n-octyl, n-nonyl, and n-decyl.

15 The term "alkylcarbonyl," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkylcarbonyl include, but are not limited to, acetyl, 1-oxopropyl, 2,2-dimethyl-1-oxopropyl, 1-oxobutyl, and 1-oxopentyl.

20 The term "alkylcarbonylalkenyl," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined herein. Representative examples of alkylcarbonylalkenyl include, but are not limited to, 3-oxo-1-butenyl and 3-oxo-1-pentenyl.

25 The term "alkylcarbonylalkoxy," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of alkylcarbonylalkoxy include, but are not limited to, 3-oxopentyloxy, 3-oxobutoxy and 2-oxopropoxy.

30 The term "alkylcarbonylalkyl," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkylcarbonylalkyl include, but are not limited to, 2-oxopropyl, 3,3-dimethyl-2-oxopropyl, 3-oxobutyl, and 3-oxopentyl.

35 The term "alkylcarbonylalkynyl," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an alkynyl group, as defined herein. Representative examples of alkylcarbonylalkynyl include, but are not limited to, 3-oxo-1-pentynyl and 3-oxo-1-pentenyl.

 The term "alkylcarbonylalkylthio," as used herein, refers to an alkylcarbonylalkyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom, as defined herein. Representative examples of alkylcarbonylalkylthio include, but are not limited to, (2-oxopropyl)sulfanyl, (3-oxobutyl)sulfanyl and (3-oxopentyl)sulfanyl.

The term "alkylcarbonyloxy," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkylcarbonyloxy include, but are not limited to, acetyloxy, ethylcarbonyloxy, and tert-butylcarbonyloxy.

5 The term "alkylcarbonylthio," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom. Representative examples of alkylcarbonylthio include, but are not limited to, acetylsulfanyl, propionylsulfanyl and (2,2-dimethylpropanoyl)sulfanyl.

10 The term "alkylene," denotes a divalent group derived from a straight or branched chain hydrocarbon of from 1 to 10 carbon atoms. Representative examples of alkylene include, but are not limited to, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2-$.

The term "alkylsulfinyl," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a sulfinyl group, as defined herein. Representative examples of alkylsulfinyl include, but are not limited to, methylsulfinyl and ethylsulfinyl.

15 The term "alkylsulfinylalkyl," as used herein, refers to an alkylsulfinyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkylsulfinylalkyl include, but are not limited to, methylsulfinylmethyl and ethylsulfinylmethyl.

20 The term "alkylsulfonyl," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of alkylsulfonyl include, but are not limited to, methylsulfonyl and ethylsulfonyl.

25 The term "alkylsulfonylalkyl," as used herein, refers to an alkylsulfonyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of alkylsulfonylalkyl include, but are not limited to, methylsulfonylmethyl and ethylsulfonylmethyl.

30 The term "alkylthio," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom. Representative examples of alkylthio include, but are not limited, methylsulfanyl, ethylsulfanyl, tert-butylsulfanyl, and hexylsulfanyl.

35 The term "alkylthioalkoxy," as used herein, refers to an alkylthio group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of alkylthioalkoxy include, but are not limited, 2-methylsulfanylethoxy and 2-ethylsulfanylethoxy.

The term "alkylthioalkyl," as used herein, refers to an alkylthio group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

Representative examples of alkylthioalkyl include, but are not limited, methylsulfanylmethyl and 2-(ethylsulfanyl)ethyl.

The term "alkynyl," as used herein, refers to a straight or branched chain hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond. Representative examples of alkynyl include, but are not limited, to acetylenyl, 1-propynyl, 2-propynyl, 3-butynyl, 2-pentynyl, and 1-butynyl.

The term "alkynyloxy," as used herein, refers to an alkynyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkynyloxy include, but are not limited to, 3-pentynyloxy, 3-butynyloxy and 2-propynyloxy.

The term "alkynylthio," as used herein, refers to an alkynyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom. Representative examples of alkynylthio include, but are not limited, 2-propynylsulfanyl, 3-butynylsulfanyl and 3-pentynylsulfanyl.

The term "alkynylene," denotes a divalent group derived from a straight or branched chain hydrocarbon of from 2 to 10 carbon atoms containing at least one triple bond. Representative examples of alkynylene include, but are not limited to, $-C\equiv C-$, $-CH_2C\equiv C-$, $-CH(CH_3)CH_2C\equiv C-$, $-C\equiv CCH_2-$, and $-C\equiv CCH(CH_3)CH_2-$.

The term "aryl," as used herein, refers to a monocyclic-ring system or a bicyclic-ring system wherein one or more of the fused rings are aromatic. Representative examples of aryl include, but are not limited to, anthracenyl, azulenyl, fluorenyl, indanyl, indenyl, naphthyl, phenyl, and tetrahydronaphthyl.

The aryl groups of this invention can be substituted with 1, 2, 3, 4 or 5 substituents independently selected from alkenyl, alkenylthio, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkoxy, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkoxy, alkylcarbonylalkyl, alkylcarbonylalkylthio, alkylcarbonyloxy, alkylcarbonylthio, alkylsulfinyl, alkylsulfinylalkyl, alkyl sulfonyl, alkylsulfonylalkyl, alkylthio, alkylthioalkyl, alkylthioalkoxy, alkynyl, alkynyloxy, alkynylthio, carboxy, carboxyalkoxy, carboxyalkyl, cyano, cyanoalkoxy, cyanoalkyl, cyanoalkylthio, ethylenedioxy, formyl, formylalkoxy, formylalkyl, haloalkenyl, haloalkenyloxy, haloalkoxy, haloalkyl, haloalkynyl, haloalkynyloxy, halogen, hydroxy, hydroxyalkoxy, hydroxyalkyl, mercapto, mercaptoalkoxy, mercaptoalkyl, methylenedioxy, nitro, $-NR_8R_9$, $(NR_8R_9)alkoxy$, $(NR_8R_9)alkyl$, $(NR_8R_9)carbonyl$, $(NR_8R_9)carbonylalkoxy$, $(NR_8R_9)carbonylalkyl$, $(NR_{10}R_{11})sulfonyl$, $(NR_{10}R_{11})sulfonylalkyl$, $-NR_{10}S(O)_2R_{12}$, $-NR_{10}S(O)_2NR_{13}R_{14}$, and $-S(O)_2OH$.

The term "arylalkoxy," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein.

Representative examples of arylalkoxy include, but are not limited to, 2-phenylethoxy, 3-naphth-2-ylpropoxy, and 5-phenylpentyloxy.

5 The term "arylalkoxycarbonyl," as used herein, refers to an arylalkoxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylalkoxycarbonyl include, but are not limited to, benzyloxycarbonyl and naphth-2-ylmethoxycarbonyl.

10 The term "arylalkyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, and 2-naphth-2-ylethyl.

The term "arylalkylcarbonyl," as used herein, refers to an arylalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylalkylcarbonyl include, but are not limited to, phenylacetyl and 3-phenylpropanoyl.

15 The term "arylalkylsulfonyl," as used herein, refers to an arylalkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of arylalkylsulfonyl include, but are not limited to, benzylsulfonyl and 2-phenylethylsulfonyl.

20 The term "arylcarbonyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylcarbonyl include, but are not limited to, benzoyl, 4-cyanobenzoyl, and naphthoyl.

25 The term "aryloxy," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through an oxy moiety, as defined herein. Representative examples of aryloxy include, but are not limited to, phenoxy, naphthyloxy, 3-bromophenoxy, 4-chlorophenoxy, 4-methylphenoxy, and 3,5-dimethoxyphenoxy.

30 The term "arylsulfonyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of arylsulfonyl include, but are not limited to, phenylsulfonyl, 4-bromophenylsulfonyl and naphthylsulfonyl.

The term "carbonyl," as used herein, refers to a -C(O)- group.

The term "carboxy," as used herein, refers to a -CO₂H group.

35 The term "carboxyalkenyl," as used herein, refers to a carboxy group, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined herein. Representative examples of carboxyalkenyl include, but are not limited to, carboxymethoxy, carboxyethenyl and 3-carboxy-1-propenyl.

The term "carboxyalkoxy," as used herein, refers to a carboxy group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein.

Representative examples of carboxyalkoxy include, but are not limited to, carboxymethoxy, 2-carboxyethoxy and 3-carboxypropoxy.

The term "carboxyalkyl," as used herein, refers to a carboxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

5 Representative examples of carboxyalkyl include, but are not limited to, carboxymethyl, 2-carboxyethyl, and 3-carboxypropyl.

The term "carboxyalkynyl," as used herein, refers to a carboxy group, as defined herein, appended to the parent molecular moiety through an alkynyl group, as defined herein. Representative examples of carboxyalkynyl include, but are not limited to,

10 carboxymethoxy, carboxyethynyl and 3-carboxy-1-propynyl.

The term "cyano," as used herein, refers to a -CN group.

The term "cyanoalkoxy," as used herein, refers to a cyano group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein.

15 Representative examples of cyanoalkoxy include, but are not limited to, cyanomethoxy, 2-cyanoethoxy and 3-cyanopropoxy.

The term "cyanoalkyl," as used herein, refers to a cyano group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of cyanoalkyl include, but are not limited to, cyanomethyl, 2-cyanoethyl, and 3-cyanopropyl.

20 The term "cyanoalkylthio," as used herein, refers to a cyanoalkyl group, as defined herein, appended to the parent molecular moiety through a sulfur atom, as defined herein. Representative examples of cyanoalkylthio include, but are not limited to, cyanomethylsulfanyl, 2-cyanoethylsulfanyl and 3-cyanopropylsulfanyl.

The term "cycloalkyl," as used herein, refers to a saturated cyclic hydrocarbon group containing from 3 to 8 carbons. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

The cycloalkyl groups of this invention can be substituted with 1, 2, 3, 4 or 5 substituents independently selected from alkenyl, alkenylthio, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkoxy, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkoxy, alkylcarbonylalkyl, alkylcarbonylalkylthio, alkylcarbonyloxy, alkylcarbonylthio, alkylsulfinyl, alkylsulfinylalkyl, alkyl sulfonyl, alkylsulfonylalkyl, alkylthio, alkylthioalkyl, alkylthioalkoxy, alkynyl, alkynyloxy, alkynylthio, carboxy, carboxyalkoxy, carboxyalkyl, cyano, cyanoalkoxy, cyanoalkyl, cyanoalkylthio, formyl, formylalkoxy, formylalkyl, haloalkenyl, haloalkenyloxy, haloalkoxy, haloalkyl, haloalkynyl, haloalkynyloxy, halogen, hydroxy, hydroxyalkoxy, hydroxyalkyl, mercapto, mercaptoalkoxy, mercaptoalkyl, nitro, -NR₈R₉, (NR₈R₉)alkoxy, (NR₈R₉)alkyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkoxy, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl,

(NR₁₀R₁₁)sulfonylalkyl, -NR₁₀S(O)₂R₁₂, -NR₁₀S(O)₂NR₁₃R₁₄, and -S(O)₂OH.

The term "cycloalkylalkoxy," as used herein, refers to a cycloalkyl group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of cycloalkylalkoxy include, but are not limited to,
5 cyclopropylmethoxy, 2-cyclobutylethoxy, 2-cyclopropylethoxy, cyclopentylmethoxy, cyclohexylmethoxy, and 4-cycloheptylbutyl.

The term "cycloalkylalkyl," as used herein, refers to a cycloalkyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of cycloalkylalkyl include, but are not limited to,
10 cyclopropylmethyl, 2-cyclobutylethyl, cyclopentylmethyl, cyclohexylmethyl, and 4-cycloheptylbutyl.

The term "cycloalkylalkylcarbonyl," as used herein, refers to cycloalkylalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of cycloalkylalkylcarbonyl include, but are not
15 limited to, cyclopropylcarbonyl, cyclohexylacetyl and 3-cyclohexylpropanoyl.

The term "cycloalkylalkylsulfonyl," as used herein, refers to cycloalkylalkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of cycloalkylalkylsulfonyl include, but are not limited to, cyclohexylmethylsulfonyl and 2-cyclohexylethylsulfonyl.

The term "cycloalkylcarbonyl," as used herein, refers to cycloalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of cycloalkylcarbonyl include, but are not limited to, cyclopropylcarbonyl, 2-cyclobutylcarbonyl, and cyclohexylcarbonyl.

The term "cycloalkyloxy," as used herein, refers to cycloalkyl group, as defined herein, appended to the parent molecular moiety through an oxy group, as defined herein. Representative examples of cycloalkyloxy include, but are not limited to, cyclopropyloxy and cyclobutyloxy.

The term "cycloalkylsulfonyl," as used herein, refers to cycloalkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of cycloalkylsulfonyl include, but are not limited to,
30 cyclohexylsulfonyl and cyclobutylcarbonyl.

The term "ethylenedioxy," as used herein, refers to a -O(CH₂)₂O- group wherein the oxygen atoms of the ethylenedioxy group are attached to two adjacent carbon atoms of the parent molecular moiety forming a six membered ring.

35 The term "formyl," as used herein, refers to a -C(O)H group.

The term "formylalkoxy," as used herein, refers to a formyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom, as defined herein.

Representative examples of formylalkoxy include, but are not limited to, 2-formylethoxy and 3-formylpropoxy.

The term "formylalkyl," as used herein, refers to a formyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

5 Representative examples of formylalkyl include, but are not limited to, formylmethyl and 2-formylethyl.

The term "halo" or "halogen," as used herein, refers to -Cl, -Br, -I or -F.

The term "haloalkenyl," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined
10 herein. Representative examples of haloalkenyl include, but are not limited to, 2-bromoethenyl, 3-bromo-2-propenyl and 1-bromo-2-propenyl.

The term "haloalkenyloxy," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkenyloxy group, as defined herein. Representative examples of haloalkenyloxy include, but are not limited to, 3-bromo-
15 2-propenyloxy and 4-bromo-3-butenyloxy.

The term "haloalkoxy," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of haloalkoxy include, but are not limited to, chloromethoxy, 2-fluoroethoxy, trifluoromethoxy, and pentafluoroethoxy.

20 The term "haloalkyl," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of haloalkyl include, but are not limited to, chloromethyl, 2-fluoroethyl, trifluoromethyl, pentafluoroethyl, and 2-chloro-3-fluoropentyl.

The term "haloalkynyl," as used herein, refers to at least one halogen, as defined
25 herein, appended to the parent molecular moiety through an alkynyl group, as defined herein. Representative examples of haloalkynyl include, but are not limited to, 2-bromoethynyl, 3-bromo-2-propynyl and 1-bromo-2-propynyl.

The term "haloalkynyloxy," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkynyloxy group, as defined
30 herein. Representative examples of haloalkynyloxy include, but are not limited to, (3-bromo-2-propynyl)oxy and (4-bromo-3-butynyl)oxy.

The term "heterocycle" or "heterocyclic," as used herein, refers to a monocyclic or bicyclic ring system. Monocyclic ring systems are exemplified by any 3- or 4-membered ring containing a heteroatom independently selected from oxygen, nitrogen and sulfur; or a
35 5-, 6- or 7-membered ring containing one, two or three heteroatoms wherein the heteroatoms are independently selected from nitrogen, oxygen and sulfur. The 5-membered

ring has from 0-2 double bonds and the 6- and 7-membered rings have from 0-3 double bonds. Representative examples of monocyclic ring systems include, but are not limited to, azetidiny, azepiny, aziridiny, diazepiny, 1,3-dioxolany, dioxany, dithiany, furyl, imidazolyl, imidazoliny, imidazolidiny, isothiazolyl, isothiazoliny, isothiazolidiny, isoxazolyl, isoxazoliny, isoxazolidiny, morpholiny, oxadiazolyl, oxadiazoliny, oxadiazolidiny, oxazolyl, oxazoliny, oxazolidiny, piperaziny, piperidiny, pyranly, pyraziny, pyrazolyl, pyrazoliny, pyrazolidiny, pyridyl, pyrimidiny, pyridaziny, pyrroly, pyrroliny, pyrrolidiny, tetrahydrofurany, tetrahydrothienyl, tetraziny, tetrazolyl, thiadiazolyl, thiadiazoliny, thiadiazolidiny, thiazolyl, thiazoliny, thiazolidiny, thienyl, thiomorpholiny, 1,1-dioxidothiomorpholiny (thiomorpholine sulfone), thiopyranly, triaziny, triazolyl, and trithianly. Bicyclic ring systems are exemplified by any of the above monocyclic ring systems fused to an aryl group as defined herein, a cycloalkyl group as defined herein, or another heterocyclic monocyclic ring system. Representative examples of bicyclic ring systems include but are not limited to, for example, benzimidazolyl, benzothiazolyl, benzothienyl, benzoxazolyl, benzofurany, benzopyranly, benzothiopyranly, benzodioxiny, 1,3-benzodioxolyl, cinnoliny, indazolyl, indolyl, indoliny, indoliziny, naphthyridiny, isobenzofurany, isobenzothienyl, isoindolyl, isoindoliny, isoquinoliny, phthalaziny, pyranopyridyl, quinoliny, quinoliziny, quinoxaliny, quinazoliny, tetrahydroisoquinoliny, tetrahydroquinoliny, and thiopyranopyridyl.

The heterocycles of this invention can be substituted with 1, 2 or 3 substituents independently selected from alkenyl, alkenylthio, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxyalkoxyalkoxy, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkoxy, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkoxy, alkylcarbonylalkyl, alkylcarbonylalkylthio, alkylcarbonyloxy, alkylcarbonylthio, alkylsulfinyl, alkylsulfinylalkyl, alkyl sulfonyl, alkylsulfonylalkyl, alkylthio, alkylthioalkyl, alkylthioalkoxy, alkynyl, alkynyloxy, alkynylthio, carboxy, carboxyalkoxy, carboxyalkyl, cyano, cyanoalkoxy, cyanoalkyl, cyanoalkylthio, ethylenedioxy, formyl, formylalkoxy, formylalkyl, haloalkenyl, haloalkenyloxy, haloalkoxy, haloalkyl, haloalkynyl, haloalkynyloxy, halogen, hydroxy, hydroxyalkoxy, hydroxyalkyl, mercapto, mercaptoalkoxy, mercaptoalkyl, methylenedioxy, nitro, $-NR_8R_9$, $(NR_8R_9)alkoxy$, $(NR_8R_9)alkyl$, $(NR_8R_9)carbonyl$, $(NR_8R_9)carbonylalkoxy$, $(NR_8R_9)carbonylalkyl$, $(NR_{10}R_{11})sulfonyl$, $(NR_{10}R_{11})sulfonylalkyl$, $-NR_{10}S(O)_2R_{12}$, $-NR_{10}S(O)_2NR_{13}R_{14}$, and $-S(O)_2OH$.

The term "heterocyclealkyl," as used herein, refers to a heterocycle, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of heterocyclealkyl include, but are not limited to, pyridin-3-ylmethyl and 2-pyrimidin-2-ylpropyl.

The term "heterocyclealkylcarbonyl," as used herein, refers to a heterocyclealkyl, as

defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclealkylcarbonyl include, but are not limited to, 3-(2-pyrimidinyl)propanoyl and 2-pyridinylacetyl.

5 The term "heterocyclealkylsulfonyl," as used herein, refers to a heterocyclealkyl, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocyclealkylsulfonyl include, but are not limited to, 4-morpholinylsulfonyl and (2-pyrimidinylmethyl)sulfonyl.

10 The term "heterocyclecarbonyl," as used herein, refers to a heterocycle, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclecarbonyl include, but are not limited to, 1-piperidinylcarbonyl, 4-morpholinylcarbonyl, pyridin-3-ylcarbonyl and quinolin-3-ylcarbonyl.

15 The term "heterocyclesulfonyl," as used herein, refers to a heterocycle, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocyclesulfonyl include, but are not limited to, 1-piperidinylcarbonyl, 1-piperidinylsulfonyl, 4-morpholinylsulfonyl, pyridin-3-ylsulfonyl and quinolin-3-ylsulfonyl.

The term "hydroxy," as used herein, refers to an -OH group.

20 The term "hydroxyalkoxy," as used herein, refers to a hydroxy group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein, wherein the alkyl portion of the alkoxy group is optionally substituted with one or two hydroxy groups. Representative examples of hydroxyalkoxy include, but are not limited to, 2-hydroxyethoxy, 3-hydroxypropoxy and 2-ethyl-4-hydroxyheptyloxy.

25 The term "hydroxyalkyl," as used herein, refers to a hydroxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of hydroxyalkyl include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, and 2-ethyl-4-hydroxyheptyl.

The term "mercapto," as used herein, refers to a -SH group.

30 The term "mercaptoalkoxy," as used herein, refers to a mercapto group, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of mercaptoalkoxy include, but are not limited to, 2-mercaptoethoxy and 3-mercaptopropoxy.

35 The term "mercaptoalkyl," as used herein, refers to a mercapto group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of mercaptoalkyl include, but are not limited to, 2-mercaptoethyl and 3-mercaptopropyl.

The term "methylenedioxy," as used herein, refers to a -OCH₂O- group wherein the oxygen atoms of the methylenedioxy are attached to the parent molecular moiety through

two adjacent carbon atoms.

The term "nitrogen protecting group" or "N-protecting group," refers to groups intended to protect an amino group against undesirable reactions during synthetic procedures. Commonly used nitrogen protecting groups are disclosed in T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 3rd edition, John Wiley & Sons, New York (1999). Preferred nitrogen protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, phenylsulfonyl, benzyl, t-butoxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).


The term "nitro," as used herein, refers to a -NO₂ group.

The term "oxo," as used herein, refers to a =O moiety.

The term "sulfinyl," as used herein, refers to a -S(O)- group.

The term "sulfonyl," as used herein, refers to a -SO₂- group.

In one embodiment, compounds of the present invention have formula (II)



(II),

or a pharmaceutically acceptable salt or prodrug thereof, wherein X_{A1}, X_{D1}, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}, R_{D5}, R₁, R₂, R₃, R₄, R₅ and R₆ are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (II) wherein R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen; R₄ is alkyl; R₆ is alkylsulfonyl; and R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (II) wherein R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen; R₄ is alkyl; R₆ is (NR₁₂R₁₃)sulfonyl; and R₁₂, R₁₃, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (II) wherein R₂, R₃, R₄, R₅, X_{A1} and X_{D1} are each hydrogen; R₁ is alkoxy; R₆ is alkylsulfonyl; and R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (II) wherein R₁, R₂, R₃, R₅ and X_{D1} are each hydrogen; X_{A1} is cyano; R₄ is alkyl; R₆ is alkylsulfonyl; and R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (II)

wherein R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen; R₄ is selected from alkenyl, alkoxyalkyl, alkoxycarbonylalkenyl and hydroxyalkyl; R₆ is alkylsulfonyl; and R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula (I).

Representative compounds of formula (II) include, but are not limited to:

- 5 N-{3-[bis(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-[3-(dibenzylamino)-2-methylphenyl]methanesulfonamide;
N-[3-(dibenzylamino)-2-methylphenyl]ethanesulfonamide;
N-[3-(dibenzylamino)-2-methylphenyl]-2-propanesulfonamide;
N-{3-[benzyl(4-methoxycarbonylbenzyl)amino]-2-
10 methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-nitrobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
15 N-{3-[benzyl(2-cyanobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(3-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[(1,3-benzodioxol-5-ylmethyl)(benzyl)amino]-2-
20 methylphenyl}methanesulfonamide;
N-{3-[benzyl(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-chlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
25 N-{3-[[4-(allyloxy)benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[[4-(allyloxy)benzyl](2,4-difluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[(2-cyanobenzyl)(2-fluoro-4-methoxybenzyl)amino]-2-
methylphenyl}methanesulfonamide;
30 N-{3-[(2,4-difluorobenzyl)(4-methoxybenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[(2,4-difluorobenzyl)(2-fluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[bis(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
35 N-{3-[(2-cyanobenzyl)(2,4-difluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-methylbenzyl)amino]-2-methylphenyl}methanesulfonamide;

- N-{3-[benzyl(4-methylbenzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-{3-[benzyl(4-chlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-(3-{benzyl[4-(trifluoromethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide;
 5 N-(3-{benzyl[2-fluoro-4-(trifluoromethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide;
 N-{3-[benzyl(2,4-dichlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-{3-[benzyl(4-{[3-bromo-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
 10 N-(3-{benzyl[4-(methoxymethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide;
 N-(3-{benzyl[4-(hydroxymethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide;
 N-(3-{benzyl[4-(2-hydroxyethoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
 15 N-{3-[(2,4-difluorobenzyl)(4-propoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-[3-(dibenzylamino)-2-ethylphenyl]methanesulfonamide;
 N'-[3-(dibenzylamino)-2-methylphenyl]-N,N-dimethylsulfamide;
 20 N-{3-[bis(2-bromobenzyl)amino]-4-methoxyphenyl}methanesulfonamide;
 N-(3-{(2-bromobenzyl)[cyano(phenyl)methyl]amino}-2-methylphenyl)methanesulfonamide;
 N-[3-(dibenzylamino)-2-((E)-3-ethoxy-3-oxo-1-propenyl)phenyl]methanesulfonamide;
 25 N-[3-(dibenzylamino)-2-(hydroxymethyl)phenyl]methanesulfonamide;
 N-[3-(dibenzylamino)-2-vinylphenyl]methanesulfonamide;
 N-[3-(dibenzylamino)-2-(methoxymethyl)phenyl]methanesulfonamide; and
 N-[3-(dibenzylamino)-2-(ethoxymethyl)phenyl]methanesulfonamide.
 The following additional compounds, representative of formula (II), may be
 30 prepared by one skilled in the art using known synthetic methodology or by using synthetic methodology described in the Schemes and Examples contained herein.
 N-{3-[benzyl(4-chloro-2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-{3-[benzyl(2-chloro-4-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
 35 N-{3-[benzyl(2,4-dichlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-{3-[benzyl(4-bromo-2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[(2,4-difluorobenzyl)(4-ethoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-(3-{(2,4-difluorobenzyl)[4-(3-pentynyloxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

5 N-{3-[benzyl(4-isopropoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-(3-{benzyl[4-(methylsulfanyl)benzyl]amino}-2-methylphenyl)methanesulfonamide;

N-[3-((2,4-difluorobenzyl){4-[(3-methyl-2-butenyl)oxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;


10 N-{3-[[4-(3-butenyloxy)-2-fluorobenzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-(3-{(2,4-difluorobenzyl)[4-(2-pentynyloxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

15 N-{3-[(2,4-difluorobenzyl)(4-propoxybenzyl)amino]-2-methylphenyl}methanesulfonamide; and

N-{3-[benzyl(2,4,6-trifluorobenzyl)amino]-2-methylphenyl}methanesulfonamide.

In another embodiment, compounds of the present invention have formula (III)



20

or a pharmaceutically acceptable salt or prodrug thereof wherein X_{A1} , X_{D1} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B and B are as defined in formula (I).

25 In another embodiment, compounds of the present invention have formula (III) wherein B is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_n-$; m and n are each 0; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (III)

wherein B is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_n-$; m is 0; n is 1-6; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

5 In another embodiment, compounds of the present invention have formula (III) wherein B is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; m is 0; n is 1-6; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

10 In another embodiment, compounds of the present invention have formula (III) wherein B is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_nCH_2CH=CH-$; m is 0; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and n, R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

15 In another embodiment, compounds of the present invention have formula (III) wherein B is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; m is 0; n is 1-6; R_1 , R_2 , R_3 , R_5 , X_{A1} , X_{D1} are each hydrogen; R_4 is alkyl; and R_6 is alkylsulfonyl, and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

20 In another embodiment, compounds of the present invention have formula (III) wherein B is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mC(O)(CH_2)_n-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl and m, n, R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

25 In another embodiment, compounds of the present invention have formula (III) wherein B is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mS(CH_2)_n-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and m, n, R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

30 In another embodiment, compounds of the present invention have formula (III) wherein B is aryl wherein said aryl is phenyl; L_B is alkylene; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

35 In another embodiment, compounds of the present invention have formula (III) wherein B is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mC(OH)(CH_2)_n-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and m, n, R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (III) wherein B is cycloalkyl; L_B is $-(CH_2)_mO(CH_2)_n-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and m, n, R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (III)

wherein B is heterocycle; L_B is $-(CH_2)_mC(O)(CH_2)_n-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and $m, n, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

Representative compounds of formula (III) include, but are not limited to:

5 N-{3-[[4-(4-bromophenoxy)benzyl](2,4-difluorobenzyl)amino]-2-

methylphenyl}methanesulfonamide;

N-{3-[(4-(4-(3-ethoxy-3-oxopropyl)phenoxy)benzyl)(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

10 N-{3-[(4-(4-(2-carboxyethyl)phenoxy)benzyl)(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[[4-(benzyloxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[(2,4-difluorobenzyl)(4-{[3-phenyl-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;

15 N-{3-[(2,4-difluorobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

methyl 4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}benzoate;

20 N-{3-[(2,4-difluorobenzyl)(2-fluoro-4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[(2,4-difluorobenzyl)(4-(3-methoxyphenoxy)benzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[(2-fluorobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

25 N-{3-[(4-methoxybenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

3-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenyl)propanoic acid;

30 N-[3-(benzyl{4-[3-(2-methoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)acetic acid;

4-[4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}-2-(2-methoxyethoxy)phenyl]butanoic acid;

35 ethyl 4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenyl)butanoate;

N-[3-(benzyl{4-[3-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-

- methylphenyl]methanesulfonamide;
 methyl 4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy} benzoate;
 ethyl N-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 5 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy} benzoyl)-beta-alaninate;
 methyl 3-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy} benzoate;
 N-{3-[benzyl(4-{3-[2-(2-methoxyethoxy)ethoxy]phenoxy}benzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 10 N-[3-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl]acetamide;
 N-[3-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl]methanesulfonami
 de;
 15 N-{3-[benzyl(4-{3-[3-(dimethylamino)propoxy]phenoxy}benzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoic acid;
 3-(4-{4-[(2-bromobenzyl){2-methyl-3-
 20 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid;
 (5-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-bromophenoxy)acetic acid;
 4-{[2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]amino}-4-
 25 oxobutanoic acid;
 5-{[2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]amino}-5-
 oxopentanoic acid;
 N-[2-(3-{4-[(benzyl{2-methyl-3-
 30 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]acetamide;
 N-[2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]methanesulfonamid
 e;
 methyl 2-(3-{4-[(benzyl{2-methyl-3-
 35 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethylcarbamate;
 (3-{4-[(4-chloro-2-fluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
 3-(4-{4-[(2-fluorobenzyl){2-methyl-3-

- [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)propanoic acid;
 3-(4-{4-[[[(4-fluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)propanoic acid;
 3-(4-{4-[[[(4-chloro-2-fluorobenzyl){2-methyl-3-
 5 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)propanoic acid;
 (3-{4-[[[(2,4-difluorobenzyl){2-ethyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetic acid;
 (3-{4-[[[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-5-methylphenoxy)acetic acid;
 10 (3-{4-[[[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-5-methoxyphenoxy)acetic acid;
 (2-chloro-5-{4-[[[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetic acid;
 (4-{4-[(benzyl{2-methyl-3-
 15 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetic acid;
 4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} benzoic acid;
 N-[3-(benzyl{4-[4-(methoxymethoxy)phenoxy]benzyl} amino)-2-
 methylphenyl]methanesulfonamide;
 20 N-[3-(benzyl{4-[4-(3-hydroxypropyl)phenoxy]benzyl} amino)-2-
 methylphenyl]methanesulfonamide;
 4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)butanoic acid;
 N-[3-(benzyl{4-[4-(4-hydroxybutyl)phenoxy]benzyl} amino)-2-
 25 methylphenyl]methanesulfonamide;
 N-{3-[(3-bromobenzyl)(4-phenoxybenzyl)amino]-2-
 methylphenyl}methanesulfonamide
 N-{3-[(4-bromobenzyl)(4-phenoxybenzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 30 N-[3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)propanoyl]glycine;
 N-[3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)propanoyl]-beta-alanine;
 4-{[3-(4-{4-[(benzyl{2-methyl-3-
 35 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)propanoyl]amino} butanoic
 acid;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)butanoyl]glycine;

- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-beta-alanine;
4-{4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]amino}butanoic
5 acid;
(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propanoic acid;
10 2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-4-hydroxybutanoic acid;
N-(3-{benzyl[4-(3-hydroxyphenoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
4-(3-{4-[(benzyl{2-methyl-3-
15 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid;
5-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid;
N-[3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]-N-
20 methylglycine;
N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-N-
methylglycine;
4-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
25 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid;
5-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid;
N-[(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]glycine;
30 N-[(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-beta-alanine;
N-[(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-N-methylglycine;
4-{[(3-{4-[(benzyl{2-methyl-3-
35 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]amino}butanoic
acid;
ethyl 4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoate;

- ethyl 5-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)pentanoate;
 N-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]glycine;
 5 N-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-beta-alanine;
 4-{[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]amino} butanoic
 acid;
 10 N-[5-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)pentanoyl]glycine;
 N-[5-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)pentanoyl]-beta-alanine;
 4-{[5-(3-{4-[(benzyl{2-methyl-3-
 15 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)pentanoyl]amino} butanoic
 acid;
 N-[3-(benzyl{4-[3-(2-hydroxyethoxy)phenoxy]benzyl} amino)-2-
 methylphenyl]methanesulfonamide;
 [2-(3-{4-[(benzyl{2-methyl-3-
 20 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)ethoxy]acetic acid;
 2,4-dideoxy-6-O-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)-D-erythro-hexonic acid;
 (3-{4-[(2-bromobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetic acid;
 25 N-{3-[[4-(3-acetylphenoxy)benzyl](2,4-difluorobenzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 N-(3-{(2,4-difluorobenzyl)[4-(3,4-dimethoxyphenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-(3-{(2,4-difluorobenzyl)[4-(pent-3-ynyloxy)benzyl]amino}-2-
 30 methylphenyl)methanesulfonamide;
 N-(3-{(4-chloro-2-fluorobenzyl)[4-(methylthio)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-{3-[(4-chloro-2-fluorobenzyl)(2-fluorobenzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 35 N-(3-{benzyl[(2'-cyano-1,1'-biphenyl-4-yl)methyl]amino}-2-
 methylphenyl)methanesulfonamide;
 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)-N-(5-

- hydroxypentyl)propanamide;
 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(4-
 hydroxybutyl)propanamide;
 5 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(3-
 hydroxypropyl)propanamide;
 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-
 10 hydroxyethyl)propanamide;
 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(6-
 hydroxyhexyl)propanamide;
 N-(3-{benzyl[4-(3-isopropoxyphenoxy)benzyl]amino}-2-
 15 methylphenyl)methanesulfonamide;
 N-[3-(benzyl{4-[3-(cyclobutyl)oxy]phenoxy}benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 N-(3-{benzyl[4-(3-sec-butoxyphenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 20 N-[3-(benzyl{4-[3-(cyclopentyl)oxy]phenoxy}benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 N-[3-(benzyl{4-[3-(1-methylbutoxy)phenoxy]benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 N-[3-(benzyl{4-[3-(2-methoxy-1-methylethoxy)phenoxy]benzyl)amino]-2-
 25 methylphenyl)methanesulfonamide;
 N-[3-(benzyl{4-[3-(cyclohexyl)oxy]phenoxy}benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 N-{3-[benzyl(4-{3-[(3-methylcyclopentyl)oxy]phenoxy}benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 30 N-[3-(benzyl{4-[3-(2-ethoxy-1-methylethoxy)phenoxy]benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 N-{3-[benzyl(4-{3-[(4-methylcyclohexyl)oxy]phenoxy}benzyl)amino]-2-
 methylphenyl)methanesulfonamide;
 N-[3-(benzyl{4-[3-(cycloheptyloxy)phenoxy]benzyl)amino]-2-
 35 methylphenyl)methanesulfonamide;
 N-(3-{benzyl[4-(3-methoxyphenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-(3-{benzyl[4-(3-ethoxyphenoxy)benzyl]amino}-2-

methylphenyl)methanesulfonamide;

N-(3-{benzyl[4-(3-propoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

5 N-[3-(benzyl{4-[3-(cyclopropylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-(3-{benzyl[4-(3-butoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

N-(3-{benzyl[4-(3-isobutoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

10 N-[3-(benzyl{4-[3-(pent-3-ynloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-{3-[benzyl(4-{3-[(2E)-pent-2-enyloxy]phenoxy}benzyl)amino]}-2-methylphenyl)methanesulfonamide;

15 N-{3-[benzyl(4-{3-[(1-methylcyclopropyl)methoxy]phenoxy}benzyl)amino]}-2-methylphenyl)methanesulfonamide;

N-[3-(benzyl{4-[3-(cyclobutylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

20 N-[3-(benzyl{4-[3-(pentyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(2-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

25 N-[3-(benzyl{4-[3-(3-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(2-ethoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-{3-[benzyl(4-{3-[2-(methylthio)ethoxy]phenoxy}benzyl)amino]}-2-methylphenyl)methanesulfonamide;

30 N-[3-(benzyl{4-[3-(cyclopentylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(hexyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

35 N-[3-(benzyl{4-[3-(3,3-dimethylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(2-isopropoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(cyclohexylmethoxy)phenoxy]benzyl}amino)-2-

methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(3-methoxy-3-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

(3-{4-[(2-methylbenzyl){2-methyl-3-
5 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;

(3-{4-[(4-methylbenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;

(3-{4-[(2,4-dichlorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;

10 (3-{4-[(2-chloro-4-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;

(3-{4-[(3,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;

15 N-[3-(benzyl{4-[4-(4-hydrazino-4-oxobutyl)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide;

N~2~[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-asparagine;

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-valine;

20 N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-tyrosine;

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-methionine;

25 N~2~[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-lysine;

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-serine;

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-phenylalanine;

30 N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-tyrosine;

N~2~[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-glutamine;

35 N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-isoleucine;

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-glutamic acid;

N-[4-(4-{4-[(benzyl{2-methyl-3-

- [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)butanoyl]-D-histidine;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)butanoyl]-L-valine;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 5 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenyl)butanoyl]-L-aspartic acid;
 ethyl 4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoate;
 4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoic acid;
 10 5-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoic acid;
 (5-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-2-ethylphenoxy)acetic acid;
 ethyl 4-{[4-(3-{4-[(benzyl{2-methyl-3-
 15 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]amino} butanoat
 e;
 (5-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-2-hexylphenoxy)acetic acid;
 ethyl N-[4-(3-{4-[(benzyl{2-methyl-3-
 20 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-N-
 methylglycinate;
 N-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-N-
 methylglycine;
 25 ethyl (2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetate;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]glycine;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 30 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-beta-alanine;
 4-{[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]amino} butanoic
 acid;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 35 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-N-
 methylglycine;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-glutamic

acid;

- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-valine;
N-[4-(4-{4-[(benzyl{2-methyl-3-
5 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-serine;
N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-isoleucine;
N~2~-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-glutamine;
10 N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]glycine;
N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-beta-alanine;
4-{[5-(4-{4-[(benzyl{2-methyl-3-
15 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]amino} butanoic acid;
N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-N-methylglycine;
20 N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-L-glutamic acid;
N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-L-valine;
25 N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-L-serine;
N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-L-isoleucine;
4-{[4-(4-{4-[(benzyl{2-methyl-3-
30 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetyl]amino} butanoic acid;
N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetyl]-N-methylglycine;
N-[4-(4-{4-[(benzyl{2-methyl-3-
35 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetyl]-L-valine;
N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetyl]-L-serine;
N~2~-[4-(4-{4-[(benzyl{2-methyl-3-

- [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetyl]-L-asparagine;
N-[3-(benzyl{4-[4-(2-hydrazino-2-oxoethoxy)phenoxy]benzyl} amino)-2-methylphenyl]methanesulfonamide;
N-(3-{benzyl[4-(3-{[(2R)-2,3-dihydroxypropyl]oxy}phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
5 N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamic acid;
N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-L-leucine;
10 N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-L-aspartic acid;
N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-L-valine;
15 N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-L-serine;
N~2~[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamine;
20 (5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-fluorophenoxy)acetic acid;
(2-chloro-5-(4-(((2-fluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
(5-(4-(((2-bromobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
25 (5-(4-(((4-bromobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
(2-chloro-5-(4-(((2,4-dichlorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
30 (2-chloro-5-(4-(((4-chloro-2-fluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
(2-chloro-5-(4-(((2-chloro-4-fluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
(5-(4-(((2-bromo-4-chlorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
35 (2-chloro-5-(4-(((3,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
(5-(4-(((4-bromo-2-fluorobenzyl)(2-methyl-3-

- ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
 N-((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)glycine;
 N-((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 5 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)-beta-alanine;
 4-(((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)amino)butanoic
 acid; -
 4-((4-(5-(4-((benzyl(2-methyl-3-
 10 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-
 chlorophenoxy)butanoyl)amino)butanoic acid;
 4-((4-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic
 acid;
 15 (2R)-2-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)propanoic acid;
 N-(3-((4-(4-chloro-3-hydroxyphenoxy)benzyl)(2,4-difluorobenzyl)amino)-2-
 methylphenyl)methanesulfonamide;
 (2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 20 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
 N-(4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-beta-alanine;
 4-((4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic
 25 acid;
 N-(4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-N-
 methylglycine;
 N-(3-((2,4-difluorobenzyl)[4-(2-phenylethoxy)benzyl]amino)-2-
 30 methylphenyl)methanesulfonamide;
 N-(3-((2,4-difluorobenzyl)[4-(3-furylmethoxy)benzyl]amino)-2-
 methylphenyl)methanesulfonamide;
 N-(3-((2,4-difluorobenzyl)[4-(2-furylmethoxy)benzyl]amino)-2-
 methylphenyl)methanesulfonamide;
 35 N-{3-[[4-(1,3-benzodioxol-5-yl)methoxy]benzyl](2,4-difluorobenzyl)amino]-2-
 methylphenyl}methanesulfonamide ;
 N-{3-[[4-[(6-chloro-1,3-benzodioxol-5-yl)methoxy]benzyl](2,4-
 difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]benzoyl}phenoxy)acetic acid;
 N-{3-[(4-benzoylbenzyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide;
 N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)benzoyl]benzyl}amino)-2-
 5 methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[3-(cyclopentylmethoxy)benzoyl]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 N-(3-{benzyl[4-(4-{[(2R)-2,3-dihydroxypropyl]oxy}benzoyl)benzyl]amino}-2-
 methylphenyl)methanesulfonamide; and N-(3-{benzyl[4-(phenylthio)benzyl]amino}-2-
 10 methylphenyl)methanesulfonamide.

The following additional compounds, representative of formula (III), may be prepared by one skilled in the art using known synthetic methodology or by using synthetic methodology described in the Schemes and Examples contained herein.

N-[3-(benzyl{4-[3-(2-methoxyethoxy)phenoxy]benzyl}amino)-2-
 15 methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[4-(methoxycarbonyl)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[4-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 20 N-(3-{(2,4-difluorobenzyl)[4-(3,4-dimethoxyphenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-{3-[benzyl(4-{3-[2-(2-methoxyethoxy)ethoxy]phenoxy}benzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 N-{3-[(4-methoxybenzyl)(4-phenoxybenzyl)amino]-2-
 25 methylphenyl}methanesulfonamide;
 N-(3-{benzyl[4-(4-(3-carboxypropyl)phenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-{3-[(2-fluorobenzyl)(4-phenoxybenzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 30 N-(3-{benzyl[4-(4-(4-ethoxy-4-oxobutyl)phenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-(3-{(2,4-difluorobenzyl)[4-(3-methoxyphenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-{3-[(4-benzoylbenzyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide;
 35 N-[3-{(2,4-difluorobenzyl){4-[4-(methoxycarbonyl)phenoxy]benzyl}amino}-2-
 methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[4-carboxyphenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;

- N-[3-(benzyl{4-[3-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[[4-(3-acetylphenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
- 5 N-[3-(benzyl{4-[4-(methoxymethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-(3-{benzyl[4-(phenylsulfanyl)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- 10 N-[3-(benzyl{4-[4-(4-hydroxybutyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[(2,4-difluorobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
- N-{3-[(2,4-difluorobenzyl)(2-fluoro-4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
- 15 N-[3-{(2,4-difluorobenzyl){4-[3-(methoxycarbonyl)phenoxy]benzyl}amino}-2-methylphenyl]methanesulfonamide;
- ethyl 3-({4-[4-((2,4-difluorobenzyl)-2-methyl-3-[(methylsulfonyl)amino]anilino)methyl}phenoxy]benzoyl}amino)propanoate;
- N-(3-{(2,4-difluorobenzyl)[4-(4-methoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- 20 N-{3-[[4-(4-acetylphenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
- N-(3-{(2,4-difluorobenzyl)[4-(3-ethoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- 25 N-(3-{(2,4-difluorobenzyl)[4-(3,5-dimethylphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- N-(3-{benzyl[4-(4-(2-carboxyethyl)phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- 4-[4-((2,4-difluorobenzyl)-2-methyl-3-[(methylsulfonyl)amino]anilino)methyl}phenoxy]-N-(2-hydroxyethyl)benzamide;
- 30 4-[4-((2,4-difluorobenzyl)-2-methyl-3-[(methylsulfonyl)amino]anilino)methyl}phenoxy]-N-(3-hydroxypropyl)benzamide;
- N-{3-[(2,4-difluorobenzyl)(4-{[3-phenyl-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 35 N-{3-[[4-(2-cyclopropylethoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
- N-(3-{benzyl[4-(3-(2-carboxyethyl)phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

N-[3-((2,4-difluorobenzyl){4-[4-(hydroxymethyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;


N-{3-[{4-[4-(3-aminopropoxy)phenoxy]benzyl}(benzyl)amino]-2-methylphenyl}methanesulfonamide;

5 2-(5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-fluorophenoxy)-N-(methylsulfonyl)acetamide;

N-(3-((2,4-difluorobenzyl)(4-(4-fluoro-3-(2H-tetrazol-5-ylmethoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide

10

In another embodiment, compounds of the present invention have formula (IV)



or a pharmaceutically acceptable salt or prodrug thereof, wherein X_{A1} , X_{D1} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , B and C are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pO(CH_2)_q-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; and R_6 is alkylsulfonyl; and m , n , p , q , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pO(CH_2)_q-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and m , n , p , q , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is cycloalkyl; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pO(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl, and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

5 In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pC(O)O(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

15 In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pC(O)(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen, R_4 is alkyl; R_6 is alkylsulfonyl, and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-O(CH_2)_pC(O)(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl, and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

25 In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pC(O)N(R_7)(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

30 In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-(CH_2)_pC(O)N(R_7)(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl, and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

35 In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is heterocycle; L_B is $-(CH_2)_mO(CH_2)_n-$; L_C is $-O(CH_2)_pC(O)N(R_7)(CH_2)_q-$; $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl, and $m, n, p, q, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{C1}, R_{C2}, R_{C3}, R_{C4}, R_{C5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (IV)

wherein B is aryl wherein said aryl is phenyl; C is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mC(O)(CH_2)_n-$; L_C is $-(CH_2)_pO(CH_2)_q-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl, and m , n , p , q , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

5 In another embodiment, compounds of the present invention have formula (IV) wherein B is aryl wherein said aryl is phenyl; C is heterocycle; L_B is $-(CH_2)_mC(O)(CH_2)_n-$; L_C is $-(CH_2)_pO(CH_2)_q-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl, and m , n , p , q , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

10 The following compounds, representative of formula (IV), may be prepared by one skilled in the art using known synthetic methodology or by using synthetic methodology described in the Schemes and Examples contained herein.

N-[3-(benzyl{4-[3-(2-phenylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

15 N-{3-[[4-[3-(1,3-benzodioxol-5-ylmethoxy)phenoxy]benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide; and

N-[3-(benzyl{4-[3-(benzyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide N-[3-(benzyl{4-[3-(3-morpholin-4-ylpropoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

20 1-[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]piperidine-2-carboxylic acid;

N-{3-[benzyl(4-{3-[(1-methylpyrrolidin-3-yl)methoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;

25 N-{3-[[4-{3-[(1-acetylpyrrolidin-3-yl)methoxy]phenoxy}benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[benzyl(4-{3-[(2-oxotetrahydrofuran-3-yl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;

30 N-[3-(benzyl{4-[3-(tetrahydrofuran-2-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(tetrahydrofuran-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

N-[3-(benzyl{4-[3-(3-furylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

35 N-[3-(benzyl{4-[3-(2-furylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;


N-[3-(benzyl{4-[3-(thien-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

- N-[3-(benzyl{4-[3-(2-thien-3-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(thien-2-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 5 N-[3-(benzyl{4-[3-(2-thien-2-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[benzyl(4-{3-[2-(4-methyl-1,3-thiazol-5-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 10 N-{3-[benzyl(4-{3-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- N-[3-(benzyl{4-[3-(2-morpholin-4-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-(3-((2,4-difluorobenzyl)(4-(3-(((2S,4R)-6-oxo-4-hydroxytetrahydro-2H-pyran-2-yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide;
- 15 N-(3-(benzyl(4-(4-(((2S,4R)-6-oxo-4-hydroxytetrahydro-2H-pyran-2-yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide;
- N-(3-{benzyl[4-(3-(((2S)-5-oxopyrrolidin-2-yl)methoxy)phenoxy]benzyl)amino]-2-methylphenyl)methanesulfonamide;
- N-{3-[benzyl(4-{3-[(3-methyloxetan-3-yl)methoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 20 N-[3-(benzyl{4-[3-(tetrahydro-2H-pyran-4-yloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(tetrahydrofuran-3-yloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 25 N-(3-{benzyl[4-(3-[(2S)-1-methylpyrrolidin-2-yl)methoxy]phenoxy]benzyl)amino]-2-methylphenyl)methanesulfonamide;
- N-[3-(benzyl{4-[3-(pyridin-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[benzyl(4-{3-[(2S)-pyrrolidin-2-ylmethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 30 N-{3-[benzyl(4-{3-[(1-methylpyrrolidin-3-yl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide 2-[(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)methyl]cyclopropanecarboxylic acid;
- 35 4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)cyclohexanecarboxylic acid benzyl 3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoate 1-[4-(4-

- {4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)butanoyl]-L-proline 1-
5-[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)pentanoyl]-L-proline;
- 5 1-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-3-
carboxylic acid;
- 1-[(4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-4-
10 carboxylic acid;
- 1-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-2-
carboxylic acid;
- 1-[(4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]proline;
- 15 1-[(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetyl]piperidine-4-
carboxamide; 4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)-N-[3-(2-oxopyrrolidin-1-
20 yl)propyl]butanamide;
- 3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)-N-(2-morpholin-4-
ylethyl)propanamide;
- 3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)-N-[3-(2-oxopyrrolidin-1-
25 yl)propyl]propanamide;
- 4-{[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)butanoyl]amino}-1-
methyl-1H-pyrrole-2-carboxylic acid;
- 30 4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)-N-(2-oxotetrahydrofuran-
3-yl)butanamide 2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)-N-[3-(2-oxopyrrolidin-
1-yl)propyl]acetamide;
- 35 4-{[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)butanoyl]amino}-1-
methyl-1H-pyrrole-2-carboxylic acid;
- 5-(4-{4-[(benzyl{2-methyl-3-

- [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-oxotetrahydrofuran-3-yl)pentanamide;
 2-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-oxotetrahydrofuran-3-yl)acetamide;
 5 4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-oxotetrahydrofuran-3-yl)butanamide;
 2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-[3-(2-oxopyrrolidin-1-yl)propyl]acetamide;
 10 2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(1,3-thiazol-5-ylmethyl)acetamide;
 15 2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(3-morpholin-4-ylpropyl)acetamide;
 N-[3-(benzyl{4-[3-(2-phenylethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide; and
 20 N-[3-(benzyl{4-[3-(benzyloxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide N-[3-(benzyl{4-[3-(tetrahydrofuran-2-ylmethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[3-(tetrahydrofuran-3-ylmethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide;
 25 N-{3-[benzyl(4-{3-[2-(2-oxopyrrolidin-1-yl)ethoxy]benzoyl}benzyl)amino]-2-methylphenyl}methanesulfonamide; and
 N-{3-[benzyl(4-{3-[2-(4-methyl-1,3-thiazol-5-yl)ethoxy]benzoyl}benzyl)amino]-2-methylphenyl}methanesulfonamide.
 N-{3-[benzyl(4-(4-(3-benzyloxy-3-oxopropyl)phenoxy)benzyl)amino]-2-methylphenyl}methanesulfonamide; and
 30 N-benzyl-4-[4-((2,4-difluorobenzyl)-2-methyl-3-[(methylsulfonyl)amino]anilino)methyl]phenoxy]benzamide.

In another embodiment, compounds of the present invention have formula (V)



(V),

or a pharmaceutically acceptable salt or prodrug thereof, wherein X_{A1} , X_{D1} , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , L_B , L_C , B and C are as defined in formula (I).

In another embodiment, compounds of the present invention have formula (V) wherein B is aryl wherein said aryl is phenyl; E is aryl wherein said aryl is phenyl; L_B is $-(CH_2)_mO(CH_2)_n-$; L_E is $-(CH_2)_pC(O)(CH_2)_q-$; R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen; R_4 is alkyl; R_6 is alkylsulfonyl; and m , n , p , q , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{B5} , R_{C1} , R_{C2} , R_{C3} , R_{C4} , R_{C5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula (I).

The following compounds, representative of formula (V), may be prepared by one skilled in the art using known synthetic methodology or by using synthetic methodology described in the Schemes and Examples contained herein.

(3-{4-(((4-benzoylbenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl)phenoxy}phenoxy)acetic acid; and
(5-{4-(((4-benzoylbenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy}acetic acid.

Another embodiment of the present invention relates to pharmaceutical compositions comprising a therapeutically effective amount of a compound of formula (I-IV) in combination with a pharmaceutically acceptable carrier.

Another embodiment of the present invention relates to a method of selectively modulating the antagonism effects of the glucocorticoid receptor in a mammal comprising administering an effective amount of a compound of formula (I-IV).

Another embodiment of the present invention relates to a method of treating type II diabetes in a mammal comprising administering a therapeutically effective amount of a compound of formula (I-IV).

Another embodiment of the present invention relates to a method of treating type II diabetes in a mammal comprising administering a therapeutically effective amount of a glucocorticoid receptor antagonist.

Another embodiment of the present invention relates to a method of treating symptoms of type II diabetes including, but not limited to, hyperglycemia, hyperinsulinemia, inadequate, glucose clearance, obesity, hypertension and high glucocorticoid levels in a mammal comprising administering a therapeutically effective amount of a compound of formula (I-IV).

Methods for Radioligand Binding Studies with Human Glucocorticoid and Progesterone

Receptor Cytosol

³[H]-dexamethasone (TRK 645) hereafter referred to as ³[H]-dex was purchased from Pharmacia Amersham, Uppsala, Sweden. Dexamethasone hereafter referred to as dex was purchased from SIGMA. The Costar 96-well polypropylene plates (3794 or 3365) were purchased from Life Technologies AB, Täby, Sweden. The GF/B filter (1450-521), filter cassette (1450-104), MeltiLex scintillating wax (1450-441), sample bag (1450-42), Microbeta™ 1450-PLUS and Microsealer 1495-021 were all purchased from Wallac Oy, Turku, Finland. Human glucocorticoid receptors were extracted from Sf9 cells infected with a recombinant baculovirus transfer vector containing the cloned hGR genes. Recombinant baculovirus was generated utilizing the BAC-TO-BAC expression system (Life Technologies) in accordance to instruction from the supplier. The hGR coding sequences were cloned into a baculovirus transfer vector by standard techniques. The recombinant baculoviruses expressing hGR were amplified and used to infect Sf9 cells. Infected cells were harvested 48hrs post infection. The receptors were extracted from the cell pellet with a phosphate buffer (1mM EDTA, 20mM KPO₄ (pH8), 8.6% Glycerol, 12mM MTG, 20mM Na₂MoO₄). The concentration of hGR in the extract was measured as specific ³[H]-dex binding with the G25-assay as described in J. Steroid Biochem. Molec. Biol. 50, No. 5/6, 313-318, 1994 and estimated to approximately 25nM. The extract was aliquoted and stored at -70°C.

The filter binding assay: Dilution series of the test compounds and dex as reference were made from 10mM (1mM dex) stock solutions in DMSO. 10μl of the dilutions was added in duplicates to the wells. The cell extracts were diluted 10 fold in EPMo + MTG buffer (1mM EDTA, HPO₄ 20mM (pH8), 6mM MTG). The diluted extract was added to the wells (110μl). ³[H]-dex were diluted from the stock solution to 10-10.8nM in EPMo + MTG buffer. 110μl of the diluted ³[H]-dex were added to the wells. The final concentration of hGR in the experiment was estimated to 1nM. All preparations were made in ambient temperature (20-25°C) on ice and with +4°C tempered buffers. The plates were incubated over night at +4°C (15-20 hours).

The incubation was stopped by filtration through GF/B filter on the Tomtec Cellharvester. The filtration on the Tomtec Cellharvester was programmed as follows: 1) Preparation before filtration with EP buffer (1mM EDTA 20mM HPO₄ (pH8)) 2x[Wash/Asp 0.6sec., Asp 0.5sec.] ; 2) Prewet of GF/B filter with EP + PEI buffer (EP

buffer, 0.3% Polyethylenimine) [Asp 0.8sec.]; 3) Filtration/harvesting of the 96-well incubation plate 3x[Wash/Asp 0.6sec., Asp 0.5sec.]. The GF/B filter was dried for at least 1 hour at 65°C. A MeltiLex scintillation wax was melted onto the filter with the Microsealer. The filter was placed in a samplebag, which was thereafter trimmed with scissors to fit the filter cassette. The cassette were placed in the Microbeta and measured for 1min /position, returning ccpm (corrected counts per minute).

For compounds able to displace the ^3H -dex from the receptor an IC_{50} -value (the concentration required to inhibit 50% of the binding of ^3H -dex) was determined by a non-linear four parameter logistic model;

$$b = ((b_{\max} - b_{\min}) / (1 + (I / \text{IC}_{50})^S)) + b_{\min} I$$

where b is the amount of bound ligand as measured by tritium counting, I is added concentration of binding inhibitor, IC_{50} is the concentration for inhibitor at half maximal binding and S is a slope factor (Hagglad, J., Carlsson, B., Kivelä, P., Siitari H., (1995) Biotechniques 18, 146-151). For determinations of the concentration of ^3H -dex in the solutions, regular scintillation counting in a Wallac Rackbeta 1214 was performed using the scintillation cocktail Supermix™ (Wallac).

The Microbeta-instrument generates the mean cpm (counts per minute) value / minute and corrects for individual variations between the detectors thus generating corrected cpm values. It was found that the counting efficiency between detectors differed with less than five percent.

A similar protocol was employed to measure affinity of the compounds of the present invention for progesterone receptor (PR).

Compounds of the present invention are active in the GR binding assay described above, and show selectivity for GR over PR, as indicated in Table 1.

Table 1
Glucocorticoid Receptor and Progesterone Receptor Binding

Example Number	GR Binding (% Inhibition at 1.7 μM)	PR Binding (IC_{50} , nM)
1	88.1	600
2	91.3	7,610
3	85.3	ND
4	78.2	ND
5	75.2	ND
6	74.1	ND
7	87.6	ND
8	75.9	ND
9	88.5	4,050

10	90.8	2,310
11	88.6	>10,000
12	86.6	1,000
13	84.0	1,860
14	82.5	2,230
15	77.9	ND
16	75.6	ND
17	88.9	1,070
18	92.6	1,390
19	91.0	ND
20	84.2	430
21	71.5	ND
22	91.0	1,480
23	92.0	920
24	92.1	2,590
25	87.5	1,790
26	94.6	1,320
27	88.9	560
28	91.3	990
29	92.4	>10,000
30	73.8	ND
31	76.9	ND
32	91.7	1,730
33	92.7	>10,000
34	90.0	>10,000
35	92.3	>10,000
36	85.5	1,200
37	72.5	ND
38	79.6	ND
39	84.5	ND
40	87.0	>10,000
41	81.2	ND
42	85.8	27,200
43	91.1	660
44	89.2	ND
45	92.0	1,720

46	81.0	ND
47	76.2	ND
48	ND	ND
49	84.7	ND
50	92.9	190
62	89.1	246
63	96.2	648
64	93.5	1178
65	94.0	633
66	92.3	ND
67	89.0	667
68	90.4	ND
69	86.9	122
70	93.6	251
71	95.2	537
72	92.0	462
73	95.0	ND
74	97.2	ND
75	98.3	2352
76	96.5	149
77	97.1	249
78	97.3	>1000
79	97.2	1103
80	97.1	>1000
81	96.2	622
82	96.9	ND
83	98.1	177
84	97.9	468
85	96.9	351
86	97.5	216
96	88.0	>1000
97	90.1	772
98	87.4	ND
99	92.0	ND
100	92.7	788
101	92.9	1032

102	ND	ND
103	92.4	78.3
104	95.7	583
105	96.2	685
106	96.0	1292
107	95.9	743
108	96.1	362
109	96.6	642
110	96.5	214
111	95.3	773
112	96.9	ND
113	96.4	ND
114	96.7	1373
115	96.4	165
116	95.3	ND
117	98.1	ND
138	91.8	ND
139	92.5	ND
140	88.1	ND
141	85.4	8050
142	90.4	ND
143	89.3	>10000
144	94.7	1300
145	94.0	357
146	28.1	ND
147	93.5	1326
148	95.6	1275
149	94.7	4287
150	94.3	2554
151	94.3	551
152	94.3	896
153	91.7	255
154	92.3	921
155	92.8	499
156	94.9	ND
157	91.8	ND

158	93.1	ND
231	96.0	ND
232	94.9	ND
233	96.8	ND
234	96.1	ND
235	96.7	ND
236	96.1	ND
237	96.7	ND
238	94.2	ND
239	93.4	ND
240	93.3	ND
241	90.6	ND
242	94.1	ND
243	92.0	ND
244	95.3	ND
245	96.3	ND
246	88.6	ND
247	90.9	ND
248	96.7	ND
249	95.4	ND
250	92.9	ND
251	94.2	ND
252	91.9	ND
253	93.2	ND
254	92.8	ND
255	93.6	ND
256	92.6	ND
257	90.6	ND
258	91.6	ND
259	91.2	ND
260	91.3	ND
261	91.5	ND
262	92.2	ND
263	91.7	ND
264	89.5	ND
265	90.0	ND

266	90.7	ND
267	92.2	ND
268	90.8	ND
269	90.7	ND
270	91.1	ND
271	91.0	ND
272	90.2	ND
273	91.2	ND
274	89.9	ND
275	90.5	ND
276	89.2	ND
277	90.6	ND
278	90.1	ND
279	91.7	ND
280	89.5	ND
281	90.0	ND
282	89.9	ND
283	89.7	ND
284	89.6	ND
285	89.7	ND
286	92.6	ND
287	89.2	ND
288	89.8	ND
289	87.8	ND
290	89.1	ND
291	87.7	ND
292	89.4	ND
305	96.6	8.9
306	97.0	ND
307	97.0	ND
308	95.7	ND
309	97.4	ND
310	97.5	ND
311	97.4	ND
312	97.3	ND
313	96.3	ND

314	96.6	ND
315	97.0	ND
316	96.3	ND
317	96.5	ND
318	95.1	ND
319	96.4	ND
320	96.8	ND
321	96.4	ND
322	98.5	360.5
323	98.2	66.6
324	97.4	122.9
325	97.7	122.2
326	97.0	183.3

The data in Table 1 indicates that the compounds of the present invention are selective for binding to glucocorticoid receptors over progesterone receptors and therefore may be useful for the treatment of type II diabetes and related metabolic disorders.

5 Compounds of the present invention may be used for the treatment of diseases associated with an excess or deficiency of glucocorticoids. Such diseases include but are not limited to the following: diabetes, obesity, Syndrome X, Cushing's Syndrome, Addison's disease, inflammatory diseases such as asthma, rhinitis and arthritis, allergy, autoimmune disease, immunodeficiency, anorexia, cachexia, bone loss or bone frailty, and wound
10 healing.

 For the treatment of diabetes or Syndrome X, compounds of the present invention may be used alone, or in combination with any existing anti-diabetic agent. Agents which may be used in combination with the compounds of the present invention include, but are not limited to insulin, an insulin analog such as mecasecmin and the like, an insulin
15 secretagogue such as nateglinide and the like, a biguanide such as metformin and the like, a sulfonylurea such chlorpropamide, glipizide, glyburide, and the like, an insulin sensitizing agent such as troglitazone, pioglitazone, rosiglitazone, and the like, an α -glucosidase inhibitor such as acarbose, voglibose, miglitol and the like, an aldose reductase inhibitor
20 such as zopolrestat and the like, a metiglinide such as repaglinide and the like, or a glycogen phosphorylase inhibitor. Other such anti-diabetic agents are known to one skilled in the art. The ability of the compounds of the present invention to treat diabetes, alone or in combination with another agent, can be demonstrated according to the methods described by Friedman, J.E., Y. Sun, T. Ishizuka, C. J. Farrell, S.E. McCormack, L.M. Herron, P. Hakimi, P. Lechner, and J.S. Yun, in J. Biol. Chem. 272 (50): 31475-31481, 1997; or,

THIS PAGE BLANK (USPTO)

according to the methods described herein.

For the treatment of obesity, compounds of the present invention may be used alone, or in combination with any existing anti-obesity agent. Agents which may be used in combination with the compounds of the present invention include, but are not limited to
5 fatty acid uptake inhibitors such as orlistat and the like, monoamine reuptake inhibitors such as sibutramine and the like, anorectic agents such as dexfenfluramine, bromocryptine, and the like, sympathomimetics such as phentermine, phendimetrazine, mazindol, and the like, or thyromimetic agents. Other such anti-obesity agents are known to one skilled in the art. The ability of the compounds of the present invention to treat obesity, alone or in
10 combination with another agent, can be demonstrated according to the methods described by Walker, H.C., and D.R. Romsos, in *Am. J. Physiol.* 262 (Endocrinol. Metab. 25): E110-E117, 1992; or according to the methods described by Langley, S.C., and D.A. York, in *Am. J. Physiol.* 259 (Regulatory Integrative Comp. Physiol. 28): R539-R544, 1990.

For the treatment of inflammatory diseases such as asthma, rhinitis and arthritis,
15 compounds of the present invention may be used alone, or in combination with any existing anti-inflammatory agent. Agents which may be used in combination with the compounds of the present invention include, but are not limited to glucocorticoid receptor agonists such as prednisolone, cortisone, dexamethasone and the like, or non-steroidal anti-inflammatory agents such as ibuprofen, ketoprofen, diclofenac, and the like. Other such anti-
20 inflammatory agents are known to one skilled in the art. The ability of the compounds of the present invention to treat inflammatory disease, alone or in combination with another agent, can be demonstrated according to the methods described by Taraye, J.P., M. Barbara, M. Aliaga, and J. Tisne-Versailles, in *Arzneim.-Forsch./Drug Res.* 40 (II)Nr. 10: 1125-1131, 1990. The ability of the compounds of the present invention to treat arthritis, alone or
25 in combination with another agent, can be demonstrated according to the methods described by Smith, R.J., and L.M. Sly, in *J. Pharmacol. Exp. Ther.* 277 (3): 1801-1813, 1996. The ability of the compounds of the present invention to treat asthma, alone or in combination


THF for tetrahydrofuran; and Ts for tosylate or $-S(O)_2-(para-CH_3Ph)$.

Preparation of Compounds of The Invention

The compounds and processes of the present invention will be better understood in connection with the following synthetic schemes and methods which illustrate a means by which the compounds of the invention can be prepared.

The compounds of this invention can be prepared by a variety of procedures and synthetic routes. Representative procedures and synthetic routes are shown in, but are not limited to, Schemes 1-20.

Scheme 1




Diaminobenzenes of general formula (7), wherein $R_1, R_2, R_3, R_4, R_6, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula I, may be prepared as described in Scheme 1. Nitroanilines of general formula (1), purchased or prepared using methodology known to those in the art, may be treated with alkylating agents such as benzyl

diisopropylamine to provide benzyl compounds of general formula (3). Benzyl compounds of general formula (3) may be treated with alkylating agents such as benzyl halides, mesylates, tosylates or triflates of general formula (4) and a base such as diisopropylamine to provide dibenzyl compounds of general formula (5). Dibenzyl compounds of general formula (5) may be reduced with a metal like iron or zinc or reduced by hydrogenation using a catalyst containing Pd, Rh or Pt to provide diamino compounds of general formula (6). Diamino compounds of general formula (6) may be alkylated, acylated, or sulfonylated with alkyl halides, acid chlorides or sulfonyl chlorides to provide diaminobenzenes of general formula (7).

Alternatively, nitroanilines of general formula (1) may be treated with 2.0 equivalents (or greater than 2.0 equivalents) of alkylating agents such as benzyl halides, mesylates, tosylates or triflates of general formula (2) or (4) in the presence of a base such as diisopropylamine to provide symmetrical dibenzyl compounds of general formula (5) directly.

Scheme 2




- Diaminobenzenes of general formula (7), wherein R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula I, may be prepared as described in Scheme 2. Nitroanilines of general formula (1) may be treated with benzaldehydes of general formula (9) under reductive amination conditions well known to those in the art, for example in the presence of a hydride reducing agent like sodium borohydride, sodium cyanoborohydride, sodium triacetoxyborohydride or lithium aluminum hydride, to provide benzyl compounds of general formula (3). Benzaldehydes of general formula (9) may be purchased or prepared, for example from benzoic acids or benzyl alcohols, using methodology well known to those in the art. Benzyl compounds of general formula (3) may be treated with benzaldehydes of general formula (10) under reductive amination conditions well known to those in the art to provide dibenzyl compounds of general formula (5). Dibenzyl compounds of general formula (5) may be processed as described in Scheme 1 to provide diaminobenzenes of general formula (7).

Nitroanilines of general formula (1) may also be treated with 2.0 equivalents (or

greater than 2.0 equivalents) of benzaldehydes of general formula (9) or (10) under reductive amination conditions to provide symmetrical dibenzyl compounds of general formula (5) directly.

5


Scheme 3



Diaminobenzenes of general formula (7), wherein R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula I, may be prepared as described in Scheme 3. Nitroanilines of general formula (1) may be treated with benzaldehydes of general formula (9) under reductive amination conditions as described in Scheme 2 to provide benzyl compounds of general formula (3). Benzyl compounds of general formula (3) may then be treated with alkylating agents such as benzyl halides, mesylates, tosylates or triflates of general formula (2) and a base such as diisopropylamine to provide dibenzyl compounds of general formula (5). Dibenzyl compounds of general formula (5) may be processed as described in Scheme 1 to provide diaminobenzenes of general formula (7).

Nitroanilines of general formula (1) may also be monoalkylated first and the alkylation product subjected to reductive amination conditions second to provide dibenzyl compounds of general formula (5).


5



Diaminobenzenes of general formula (7), wherein R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula I, may also be prepared as described in Scheme 4. Nitroanilines of general formula (11) may be alkylated, acylated, or sulfonylated with alkyl halides, acid chlorides or sulfonyl chlorides to provide nitrobenzenes of general formula (12). Nitrobenzenes of general formula (12) may be reduced with a metal like iron or zinc or reduced by hydrogenation using a catalyst containing Pd, Rh or Pt to provide diamino compounds of general formula (13). Diamino compounds of general formula (13) may be processed as described in Schemes 1, 2 or 3 to provide diaminobenzenes of general formula (7).


15

Scheme 5




- Diaminobenzenes of general formula (19), wherein R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{D1} , R_{D2} , R_{D3} , R_{D4} , R_{D5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} and R_{B5} are as defined in Formula (I), may be prepared as described in Scheme 5. Nitroanilines of general formula (14), prepared as described in Schemes 1-3, may be deprotected to provide (15). A preferred protecting group is allyl which can be removed with tetrakis(triphenylphosphine)palladium(0) and phenylsilane. Other potential protecting groups are listed in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, Inc., New York, 1999.
- Phenol (15) may be treated with an aryl group of general formula (16), copper(II) acetate and a base such as triethylamine to provide nitrobenzenes of general formula (17). Alternatively, phenol (15) may be treated with aryl halides (18), palladium salts, ligands, and bases to provide nitrobenzenes of general structure (17). Nitrobenzenes of general formula (17) may be processed as described in Scheme 1 to provide diaminobenzenes of general formula (19).

Scheme 6




Diaminobenzenes of general formula (21), wherein R is selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl and (NR₁₀R₁₁)sulfonylalkyl and R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{D1}, R_{D2}, R_{D3}, R_{D4}, R_{D5}, R₈, R₉, R₁₀ and R₁₁ are as defined in formula (I), may be prepared as described in Scheme 6. Nitroanilines of general formula (15), prepared as described in Schemes 1-5, may be treated with a base such as sodium hydride and an alkylating agent to provide nitrobenzenes of general formula (18). Alternatively, Mitsunobu reaction of phenols (15) with an alcohol ROH, trialkyl phosphine (like triphenylphosphine), and dialkyl azodicarboxylate (like diethyl azodicarboxylate) can be used to prepare nitrobenzenes of general formula (20). Nitrobenzenes of general formula (20) may be processed as described in Scheme 1 to provide diaminobenzenes of general formula (21).



Diaminobenzenes of general formula (23) and of general formula (24), wherein R_1, R_2, R_3, R_4, R_6 are as defined in Formula (I), Ar_1 and Ar_2 are each independently aryl, as defined herein, and Het_1 and Het_2 are each independently heterocycle, as defined herein, may be prepared as described in Scheme 7. Diaminobenzenes of general formula (13), prepared as described in Scheme 4 or purchased commercially, may be treated an aryl or heterocyclic halide or triflate under Buchwald coupling conditions (Wagaw and Buchwald, JOC (1996) 61, 7240-7241; Yang and Buchwald, J. Organometallic Chem., (1999) 576, 125-146; and Harris, Geis and Buchwald, JOC (1999) 64, 6016-6022) to provide secondary amines of general formula (22). Secondary amines of general formula (22) may be resubjected to Buchwald conditions to provide diamines of general formula (23), or may be treated with a benzylic or heterocyclicmethyl halide in the presence of a base such as sodium carbonate or triethylamine to provide diamines of general formula (24).


Scheme 8



- Diaminobenzenes of general formula (28) wherein R is selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR_8R_9) carbonyl, (NR_8R_9) carbonylalkyl, $(NR_{10}R_{11})$ sulfonyl and $(NR_{10}R_{11})$ sulfonylalkyl and $R_1, R_2, R_3, R_4, R_6, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in Formula (I), and diaminobenzenes of general formula (29), wherein $R_1, R_2, R_3, R_4, R_6, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}, R_{D5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}$ and R_{B5} are as defined in formula I, may be prepared as described in Scheme 8.
- Phenols of general formula (25), purchased or prepared using methodology known to those in the art, may be treated with alkylating agents such as benzyl halides, mesylates, tosylates

or triflates and a base such as potassium carbonate to provide ethers of general formula (26). Alternatively, a Mitsunobu reaction can be used to prepare ethers (26) from phenols (25) and an alcohol (ROH). Aldehydes of general formula (26) may be processed as described in Scheme 3 to provide dibenzyl compounds of general formula (27). Dibenzyl compounds of general formula (27) may be processed as described in Scheme 1 to provide compounds of general formula (28). Compounds of general formula (27) may be deprotected as described in Scheme 5 to provide diaminobenzenes of general formula (15). Compounds of general formula (15) may be processed as described in Scheme 5 to provide diaminobenzenes of general formula (29).


Scheme 9



Diaminobenzenes of general formula (37), wherein R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in Formula (I), may be prepared as described in Scheme 9. Phenols of general formula (30), purchased or prepared using methodology known to those in the art, may be treated with electrophilic agents such as 1-

(chloromethyl)-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate) to provide compounds of general formula (31). Compounds of general formula (31) may be may be protected on oxygen to provide dibenzyl compounds of general formula (32). An allyl protecting group is preferred and other potential protecting groups are listed in T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, Inc., New York, 1999. Compounds of general formula (31) may be alkylated directly using an alkylating agent like allyl bromide and a base like potassium carbonate to give phenols 32. Alternatively, one phenol can be reacted with TsCl and the other with an alkylating agent like allyl bromide and a base like potassium carbonate. The Ts group can then be removed by hydrolysis to give compounds of general formula (32). Phenols of general formula (33) may be purchased, prepared as in the preparation of compounds (32), or prepared using methodology known to those in the art. Phenols of general formula (33) can be reacted with aldehydes (34) to provide ethers of general formula (35). Ethers of general formula (35) may be processed as described in Scheme 3 to provide diaminobenzenes of general formula (36). Compounds of general formula (36) may be processed as described in Scheme 1 to provide diaminobenzenes of general formula (37).


Scheme 10



Diaminobenzenes of general formula (41) wherein R is selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl and (NR₁₀R₁₁)sulfonylalkyl and R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula I, and diaminobenzenes of general formula (43), wherein R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula I, L is a linker group, and Y is an alkyl


- group, may be prepared as described in Scheme 10. L is selected from those groups delineated for R_{B1-B4} but has two attachment sites. P is a protecting group and allyl is preferred. Representative protecting groups are described in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, Inc., New York, 1999.
- 5 Nitroaromatics of general formula (38), prepared via methods described in Scheme 9 or using methodology known to those in the art, may be deprotected to provide phenols of general formula (39). A preferred protecting group is allyl which can be removed with tetrakis(triphenylphosphine)palladium(0) and phenylsilane. Phenols of general formula (39)
- 10 may be alkylated as described in Scheme 6 to provide dibenzyl compounds of general formula (40). Dibenzyl compounds of general formula (40) may be reduced and N-substituted as described in Scheme 1 to provide diamino compounds of general formula (41). Alternatively, compounds of general formula (39) may be subjected to aryl substitution reactions as described in Scheme 6 to provide esters of general formula (42). Esters (42) may be processed as described in Scheme 1 followed by hydrolysis of the ester
- 15 to provide diamino acids compounds of general formula (43).

Scheme 11



Diaminobenzenes of general formula (45) and (47), wherein R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{D1}, R_{D2}, R_{D3}, R_{D4}, R_{D5}, R_{B1}, R_{B2}, R_{B3} and R_{B4} are as defined in formula I, may be prepared as described in Scheme 11. Nitrophenols of general formula (39) may be treated with alkylating agents such as benzyl halides, mesylates, tosylates or triflates and a base to provide compounds of general formula (44). Reduction of the nitro group followed by reaction of the resultant amine is accomplished as described in Scheme 1; treatment with an acid like aqueous hydrochloric acid or trifluoroacetic acid of the like to remove the acetonide protecting group followed by hydrolysis of the ester for example using hydroxide anion in an alcoholic solvent provides dibenzyl dihydroxyacids of general formula (45). Alternatively dibenzyl compounds of general formula (44) may be reduced and reacted on nitrogen as described in Scheme 1 to provide diamino compounds of general formula (46). Removal of the acetonide under acidic conditions followed by acid-catalyzed lactonization provides diaminobenzenes hydroxylactones of general formula (47).


Scheme 12



5 Diaminobenzenes of general formula (51) and (52), wherein $R_1, R_2, R_3, R_4, R_6, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula I, and L is a linker group, may be prepared as described in Scheme 12. L is selected from those groups delineated for R_{B1-B4} but has two attachment sites. X is a halogen like chlorine, bromine, iodine. Halides of general formula (48), prepared according to Schemes 5 or 9, or using methodology known to those in the art, may be treated with zinc reagents such as 3-ethoxy-3-oxopropylzinc bromide of general formula (49) and a palladium catalyst

such as tetrakis(triphenylphosphine)palladium (0) to provide compounds of general formula (50). Compounds of general formula (50) that contain an ester may optionally be hydrolyzed to provide dibenzyl diamino acids of general formula (51), or may optionally be reduced with diisobutylaluminum hydride or another reducing reagent to provide dibenzyl diamino alcohols of general formula (52).


Scheme 13



Diaminobenzenes of general formula (55) and (56), wherein R_7 and R_8 are selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR_8R_9) carbonyl, (NR_8R_9) carbonylalkyl, $(NR_{10}R_{11})$ sulfonyl and $(NR_{10}R_{11})$ sulfonylalkyl and $R_1, R_2, R_3, R_4, R_6, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula I, and L is a linker group, may be prepared as described in Scheme 13. L is selected from those groups delineated for R_{B1-B4} but has two attachment sites. Phenols of general formula (53), prepared using the methods described in Scheme 10 or using methodology known to those in the art, may be coupled with phthalimidoalkyl alcohols or phthalimidoalkyl halides as described in Schemes 6 and 1 to provide compounds of general formula (54). Phthalimide compounds of general formula (54) may be treated with hydrazine to provide primary

amines of general formula (55). Amine compounds of general formula (55) may be alkylated, acylated, or sulfonylated with alkyl halides, acid chlorides or sulfonyl chlorides to provide diaminobenzenes of general formula (56).


5



Diaminobenzenes of general formula (57) and (59), wherein R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula I, L is a linker group, and P is an optional protecting group, and (59), wherein R_7 and R_8 are selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR_8R_9) carbonyl, (NR_8R_9) carbonylalkyl, $(NR_{10}R_{11})$ sulfonyl and $(NR_{10}R_{11})$ sulfonylalkyl and R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula I, and L is a linker group, may be prepared as described in Scheme 14. L is selected from those groups delineated for R_{B1-B4} but has two attachment sites. Representative protecting groups are described in T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Synthesis, John

Wiley & Sons, Inc., New York, 1999. Phenols of general formula (53) may be coupled with protected diols or protected hydroxyalkyl halides as described in Schemes 6 and 1 to provide compounds of general formula (57). Compounds of general formula (57) may be deprotected as described in Scheme 5 to the corresponding alcohol and the alcohols converted to the corresponding halide or alkylsulfonate, for example by treatment with triphenylphosphine and carbon tetrabromide or p-toluenesulfonyl chloride or the like, to provide compounds of general formula (58). Compounds of general formula (58) may be reacted with a primary or secondary amine to provide compounds of general formula (59).


Scheme 15



Diaminobenzenes of general formula (63), wherein R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{A5} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula I, may be prepared as described in Scheme 15. Nitroanilines of general formula (1), purchased or prepared using methodology known to those in the art, may be treated with aldehydes (35) and processed as described in Scheme 3 to provide compounds of general formula (60). Compounds of general formula (61) may be deprotected as described in Scheme 5 and the

resultant alcohol alkylated as described in Scheme 6 to provide compounds of general formula (61). Compounds of general formula (61) may be N-alkylated as described in Schemes 1, 2, or 3 to provide compounds of general formula (62). Nitroaromatic compounds of general formula (62) may be reduced and alkylated, acylated, or sulfonylated with alkyl halides, acid chlorides or sulfonyl chlorides as described in Scheme 1.


Hydrolysis of the ester moiety and hydrolyzed to provides diaminobenzenes of general formula (63).



Diaminobenzenes of general formula (66) and (67), wherein R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula I, may be prepared as described in Scheme 16. Dinitrohalides of general formula (64), purchased or prepared using methodology known to those in the art, may be treated with tin reagents such as trimethyl or tri-n-butyl stannanes, a palladium catalyst (palladium salt plus

ligand), copper salt, and LiCl in a Stille reaction to provide nitroaromatic compounds of general formula (65). The dinitrohalides of general formula (65) may be reduced using methodology known to those in the art to provide nitroanilines of general formula (1). Compounds of general formula (1) may be processed as described in Schemes 1, 2, or 3 to provide diamino compounds of general formula (66). Compounds of general formula (1) may also be processed as described in Schemes 1, 2, 3, or 9 to provide diamino compounds of general formula (67).


Scheme 17



Diaminobenzenes of general formula (71, 72, 73, and 74), wherein R is selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl,


alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl and (NR₁₀R₁₁)sulfonylalkyl and R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{B5}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula I, and P is an optional protecting group, may be prepared as described in Scheme 17. Nitroanilines of general formula (1), purchased or prepared using methodology known to those in the art, may be processed as described in Schemes 1, 2, or 3 to provide ester compounds of general formula (68). Ester compounds of general formula (68) may be hydrolyzed and subjected to amide forming coupling conditions in the presence of *N,O*-dimethylhydroxylamine hydrochloride to provide compounds of general formula (69). Alternatively ester compounds of general formula (68) may be treated with the reagent derived from *N,O*-dimethylhydroxylamine hydrochloride and trimethylaluminum to provide compounds of general formula (69). Compounds of general formula (69) may be treated with aryl lithium or aryl Grignard compounds, prepared using methodology known to those in the art, to provide diamino ketones of general formula (71). For the specific case where R_{B5} is an alcohol, diamino compounds of general formula (72) may be protected on the anilino nitrogen (mesylate protecting group is preferred) for example using methanesulfonyl chloride or the like, and deprotected on oxygen as described in Scheme 5, to provide diaminobenzenes of general formula (73). Phenol compounds of general formula (73) may be reacted with alcohols or alkyl halides as described in Scheme 6 to give phenyl ethers; after hydrolysis, or other deprotection, to provide diphenylketones of general structure (74). Cleavage of the preferred mesylate protecting group occurs under basic hydrolytic conditions.

Scheme 18




5 Diaminobenzenes of general formula (77) wherein R is selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl and (NR₁₀R₁₁)sulfonylalkyl-R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2},

R_{D3} , R_{D4} and R_{D5} are as defined in formula I, and L is a linker group, and (79), wherein R_7 and R_8 are selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR_8R_9) carbonyl, (NR_8R_9) carbonylalkyl, $(NR_{10}R_{11})$ sulfonyl and $(NR_{10}R_{11})$ sulfonylalkyl and R_1 , R_2 , R_3 , R_4 , R_6 , R_{A1} , R_{A2} , R_{A3} , R_{A4} , R_{B1} , R_{B2} , R_{B3} , R_{B4} , R_{D1} , R_{D2} , R_{D3} , R_{D4} and R_{D5} are as defined in formula I, and L is a linker group, may be prepared as described in Scheme 18. L is selected from those groups delineated for R_{B1-B4} but has two attachment sites. Compounds of general formula (53) may be reacted with protected diols or protected hydroxyalkyl halides according to Scheme 6 to provide nitroaromatic compounds of general formula (75). Nitroaromatic compounds of general formula (75) may be deprotected to provide alcohol compounds of general formula (76). Alcohol compounds of general formula (76) may be reacted with alcohols or alkyl halides according to Schemes 6 and 1 to provide diamino compounds of general formula (77). Alternatively, alcohol compounds of general formula (76) may be mesylated with mesyl chloride to provide mesylates of general formula (78). Mesylates of general formula (78) may be reacted with primary or secondary amines according to Scheme 14 to provide diamino compounds of general formula (79).



Diaminobenzenes of general formula (82) and (83), wherein R is selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxy sulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, (NR₈R₉)carbonyl, (NR₈R₉)carbonylalkyl, (NR₁₀R₁₁)sulfonyl and (NR₁₀R₁₁)sulfonylalkyl and R₁, R₂, R₃, R₄, R₆, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4} and R_{D5} are as defined in formula I, and P is a protecting group, may be prepared as described in Scheme 19. Phenols of general formula (53) may be treated with N-protected pyrrolidinol mesylates of general formula (80) and a base to provide compounds of general formula (81). Compounds of general formula (81) may be deprotected to provide amine compounds of general formula (82). The pyrrolidine moiety in the amine compounds of general formula (82) may be alkylated, acylated, or sulfonylated with alkyl halides, acid chlorides or sulfonyl chlorides as described in Schemes 1, 2, or 3; reduction of the nitro group and reaction of the resultant amine with alkyl halides, acid chlorides or sulfonyl chlorides as described in Scheme 1 provides compounds of general formula (83).

Scheme 20



- Diaminobenzenes of general formula (85, 87, 88, and 89), wherein R_7 and R_8 are selected from alkenyl, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylthioalkyl, alkynyl, cyanoalkyl, haloalkenyl, haloalkyl, haloalkynyl, $(\text{NR}_8\text{R}_9)\text{carbonyl}$, $(\text{NR}_8\text{R}_9)\text{carbonylalkyl}$, $(\text{NR}_{10}\text{R}_{11})\text{sulfonyl}$ and $(\text{NR}_{10}\text{R}_{11})\text{sulfonylalkyl}$ and $R_1, R_2, R_3, R_4, R_6, R_{A1}, R_{A2}, R_{A3}, R_{A4}, R_{A5}, R_{B1}, R_{B2}, R_{B3}, R_{B4}, R_{D1}, R_{D2}, R_{D3}, R_{D4}$ and R_{D5} are as defined in formula I, and L_1 and L_2 are independently linkers, may be prepared as described in Scheme 20. L_1 and L_2 are selected from those groups delineated for R_{B1-B4} but have two attachment sites. Esters of general formula (84), prepared by methods described in Scheme 9 or by using methodology known to those in the art, may be hydrolyzed to provide acids of general formula (85). Acid compounds of general formula (85) may be coupled with amines of general formula (86) using appropriate coupling reagents, for example an acid anhydride or sulfonyl halide or carbodiimine reagent or the like, to provide compounds of general formula (87). For the specific case where the compounds of formula (86) are amino esters, the resultant ester compounds of general formula (88) may be hydrolyzed to provide acid compounds of general formula (89).

The compounds and processes of the present invention will be better understood by reference to the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

5

Example 1N-{3-[bis(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamideExample 1A

10

N-(2-methyl-3-nitrophenyl)methanesulfonamide

2-Methyl-3-nitroaniline (10.0 g, 65.7 mmol, purchased from Aldrich) in pyridine (70 mL) was treated with methanesulfonyl chloride (5.09 mL, 65.7 mmol) at 0° C. The reaction mixture was allowed warmed to room temperature and stirred for 2 hours. The reaction mixture was concentrated under reduced pressure and the crude products were diluted with diethyl ether (700 mL). The mixture was washed with 1N HCl (250 mL), water (250 mL) and brine (250 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated under reduced pressure to provide the title compound which was used in the next step without further purification.

20

Example 1BN-(3-amino-2-methylphenyl)methanesulfonamide

The crude product from Example 1A and 10% palladium on carbon (1.4 g) were stirred vigorously in ethyl acetate (70 mL) for 24 hours under one atmosphere of hydrogen. The reaction was placed under a nitrogen atmosphere, filtered through a 1 inch pad of 1:1 Celite:silica gel eluting with ethyl acetate and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, ethyl acetate) to provide the title compound as a brown solid (11.06 g, 84%).

30

Example 1CN-{3-[bis(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 1B (1.525 g, 7.625 mmol) and 2-bromobenzaldehyde (1.5 mL, 12.96 mmol) in dichloroethane (26.7 mL) were treated with glacial acetic acid (1.75 mL, 30.5 mmol). The yellow reaction was stirred for 4 hours at room temperature and was treated with sodium triacetoxyborohydride (3.23 g, 15.25 mmol). After stirring overnight at room temperature, the mixture was poured into saturated aqueous NaHCO₃ (200 mL), extracted with diethyl ether (200 mL) and the phases separated. The organic phase was washed with brine (100 mL), dried (Na₂SO₄), filtered and concentrated under

reduced pressure. The residue was purified by flash chromatography (silica gel, 5:2→1:1 hexanes:ethyl acetate) to provide two products. The title compound, a brown solid, was isolated as the minor product (0.29 g, 7% yield). The major product isolated was the monobenzylated analogue, N-{3-[(2-bromobenzyl)amino]-2-

5 methylphenyl}methanesulfonamide. A portion of the minor product was repurified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column.

¹H NMR (300 MHz, CDCl₃) δ 7.50 (dd, J=8.0, 0.7 Hz, 2H), 7.35 (d, J=8.1 Hz, 2H), 7.18 (m, 2H), 7.07 (m, 2H), 6.03 (s, 1H), 4.30 (s, 4H), 2.90 (s, 3H), 2.16 (s, 3H); MS (APCI+) m/z 539 (M+H)⁺.

Example 2

N-[3-(dibenzylamino)-2-methylphenyl]methanesulfonamide

Example 2A

N,N-dibenzyl-N-(2-methyl-3-nitrophenyl)amine

2-Methyl-3-nitroaniline (3.4 g, 22.3 mmol, purchased from Aldrich) and diisopropylethylamine (19.5 mL, 112 mmol) in DMF (34 mL) were treated with benzyl bromide (8.0 mL, 67 mmol) and heated for 18 hours at 90 °C. After cooling to room temperature, the mixture was diluted with diethyl ether (800 mL). The mixture was washed with saturated ammonium chloride (400 mL), water (2x400 mL), brine (400 mL), dried (Na₂SO₄), filtered and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 10:1 hexanes:ethyl acetate) to provide the title compound as a yellow oil (6.38 g, 86%).

Example 2B

N¹,N¹-dibenzyl-2-methyl-1,3-benzenediamine

The product from Example 2A (3.4 g, 16.2 mmol) in acetic acid (20 mL) was treated with zinc powder (2.68 g, 41 mmol, purchased from Aldrich) and stirred for 48 hours. An additional portion of zinc (1.0g, 15.3 mmol) was added and stirring continued for 4 hours. The reaction was filtered through a pad of Celite with CH₂Cl₂ and concentrated. Purification by flash chromatography (silica gel, 4:1 hexanes:ethyl acetate) provided the title compound as a yellow solid (1.98 g, 64%).

Example 2C

N-[3-(dibenzylamino)-2-methylphenyl]methanesulfonamide

The product from Example 2B was processed as described in Example 1A. The residue was purified by flash chromatography (silica gel, 1:1 hexanes:ethyl acetate) to provide the title compound as a brown solid.

¹H NMR (300 MHz, CDCl₃) δ 7.24 (m, 11H), 7.12 (t, J=8.0 Hz, 1H), 6.90 (d, J=7.8 Hz, 1H), 6.18 (s, 1H), 4.05 (s, 4H), 2.93 (s, 3H), 2.34 (s, 3H); MS (APCI+) m/z 381 (M+H)⁺; HRMS (CI) m/z (calcd for C₂₂H₂₄N₂O₅S₁) 380.1558, observed. 380.155.

Example 3

N-[3-(dibenzylamino)-2-methylphenyl]ethanesulfonamide

The product from Example 2B (0.03 g, 0.1 mmol) in pyridine (2 mL) at 0 °C was treated with ethanesulfonyl chloride (0.01g, 0.12 mmol). The reaction mixture was quenched after 1 hour with water (2 mL), extracted with diethyl ether (5 mL), washed with water (2 x 2 mL), rinsed with brine (2 mL), dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column to provide the title compound.

¹H NMR (500 MHz, DMSO-d₆) δ 1.22 (t, 3H, J=7.3 Hz), 2.40 (s, 3H), 3.02 (dd, 2H, J=14.6, 7.5 Hz), 4.05 (s, 4H), 6.94 (m, 2H), 7.01 (m, 1H), 7.20 (m, 2H), 7.27 (m, 7H), 8.92 (s, 1H); MS (APCI+) m/z 395 (M+H)⁺.

Example 4

N-[3-(dibenzylamino)-2-methylphenyl]-2-propanesulfonamide

The product from Example 2B and isopropylsulfonylchloride were processed as described in Example 3 to provide the title compound.

¹H NMR (500 MHz, DMSO-d₆) δ 1.23 (d, 6H, J=6.5 Hz), 2.39 (s, 3H), 3.18 (m, 1H), 4.04 (s, 4H), 6.91 (dd, 1H, J=7.5, 1.6 Hz), 6.97 (m, 2H), 7.19 (m, 2H), 7.26 (m, 8H), 8.86 (s, 1H); MS (APCI+) m/z 409 (M+H)⁺.

Example 5

N'-[3-(dibenzylamino)-2-methylphenyl]-N,N-dimethylsulfamide

The product from Example 2B and N,N-dimethylsulfamoylchloride were processed as described in Example 3 to provide the title compound.

¹H NMR (500MHz, DMSO-d₆) δ 2.25 (s, 3H), 2.33 (m, 6H), 3.88 (s, 4H), 6.72 (dd, 1H, J=7.7, 1.5 Hz), 6.82 (m, 2H), 7.03 (m, 2H), 7.10 (m, 8H), 8.73 (s, 1H); MS (APCI+) m/z 410 (M+H)⁺.

Example 6

N-{3-[benzyl(4-methoxycarbonylbenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 6A

N-benzyl-2-methyl-3-nitroaniline

5 2-Methyl-3-nitroaniline (3.40 g, 22.3 mmol) and benzaldehyde (3.9 mL, 38 mmol) in dichloroethane (78 mL) were treated with glacial acetic acid (5.1 mL, 89 mmol). The yellow reaction mixture was stirred for 4 hours at room temperature, treated with sodium triacetoxyborohydride (9.5 g, 44.6 mmol) and allowed to stir overnight at room temperature. The mixture was poured into saturated aqueous NaHCO₃ (400 mL),
10 extracted with diethyl ether (400 mL) and the phases separated. The organic phase was washed with brine (200 mL), dried (Na₂SO₄), filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 10:1→4:1 hexanes:ethyl acetate) to provide the title compound as a yellow solid (4.72 g, 87%).

15

Example 6B

N-benzyl-N-(4-methoxycarbonylbenzyl)-2-methyl-3-nitroaniline

20 The product from Example 6A (0.92 g, 3.8 mmol) and diisopropylethylamine (1.7 mL, 9.5 mmol) in DMF (9.5 mL) were treated with methyl 4-(bromomethyl)benzoate (0.74 g, 7.6 mmol) and heated for 18 hours at 90 °C. After cooling to room temperature, the mixture was diluted with diethyl ether (200 mL). The mixture was washed with saturated ammonium chloride (200 mL), water (2x200 mL), brine (150 mL), dried (Na₂SO₄), filtered and the filtrate was concentrated under reduced pressure. The residue
25 was purified by flash chromatography (silica gel, 10:1→4:1 hexanes:ethyl acetate) to provide the title compound as a yellow solid (1.26 g, 85%).

Example 6C

N¹-benzyl-N¹-(4-methoxycarbonylbenzyl)-2-methyl-1,3-benzenediamine

30 The product from Example 6B (1.43 g, 3.67 mmol) and NH₄Cl (0.14 g, 2.57 mmol) in ethanol (18.4 mL) and water (6.4 mL) were treated with iron powder (1.43 g, 25.7 mmol) and heated at 80 °C for one hour. The reaction mixture was allowed to cool to room temperature, diluted with CH₂Cl₂ (100 mL), filtered through Celite with ethyl acetate, and the filtrate was concentrated under reduced pressure to provide the title compound which was used without further purification.

35

Example 6D

N-{3-[benzyl(4-methoxycarbonylbenzyl)amino]-2-methylphenyl}methanesulfonamide

The crude product from Example 6C in pyridine (8.8 mL) was treated with methanesulfonyl chloride (0.28 mL, 3.60 mmoles) at 0 °C. The reaction mixture turned from clear and colorless to yellow. After 30 minutes, the mixture was quenched with water (5.0 mL) and was concentrated under reduced pressure. The residue was dissolved in diethyl ether (200 mL), washed with water (2x100 mL), brine (100 mL), dried (Na₂SO₄), filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 4:1→1:1 hexanes:ethyl acetate) to provide the title compound as a yellow solid (1.33 g, 83%). A small portion of the residue (0.1 g) was repurified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column.

¹H NMR (300 MHz, CDCl₃) δ 7.92 (d, J=6.8 Hz, 2H), 7.21 (m, 8H), 7.11 (t, J=8.0 Hz, 1H), 6.89 (d, J=8.0 Hz, 1H), 6.11 (s, 1H), 4.12 (s, 2H), 4.05 (s, 2H), 3.90 (s, 3H), 2.95 (s, 3H), 2.34 (s, 3H); MS (APCI+) m/z 439 (M+H)⁺.

Example 7

N-{3-[benzyl(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 1-bromo-2-(bromomethyl)benzene were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.50 (dd, J=8.0, 1.1 Hz, 1H), 7.34 (dd, J=7.4, 1.7 Hz, 1H), 7.03-7.29 (m, 9H), 7.01 (dd, J=8.0, 1.2 Hz, 1H), 6.04 (s, 1H), 4.20 (s, 2H), 4.13 (s, 2H), 2.90 (s, 3H), 2.23 (s, 3H); MS (ESI+) m/z 461 (M+H)⁺.

Example 8

N-{3-[benzyl(4-nitrobenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 8A

N-[3-(benzylamino)-2-methylphenyl]methanesulfonamide

The product from Example 1B and benzaldehyde were processed as described in Example 1C to provide the title compound.

Example 8B

N-{3-[benzyl(4-nitrobenzyl)amino]-2-methylphenyl}methanesulfonamide

Glacial acetic acid (0.37 mL, 6.4 mmoles) was added to a solution of Example 8A (0.46 g, 1.6 mmoles) and 4-nitrobenzaldehyde (0.48 g, 3.2 mmoles) in dichloroethane (3.2 mL) and CH₃CN (4.0 mL). The reaction was stirred for 4 hours at room temperature and was treated with sodium triacetoxyborohydride (0.68 g, 3.2 mmoles). The reaction mixture

was stirred overnight at room temperature. The mixture was poured into saturated aqueous NaHCO₃ (150 mL) and extracted with diethyl ether (150 mL). The organic phase was washed with brine (150 mL), dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (5:2→1:1 hexanes:ethyl acetate) to provide the title compound as a brown solid (0.034 g, 5% yield).
5 ¹H NMR (300 MHz, CDCl₃) δ 8.12 (d, J=8.9 Hz, 2H), 7.37 (d, J=8.9 Hz, 2H), 7.27 (m, 6H), 7.12 (t, J=7.8 Hz, 1H), 6.87 (d, J=8.1 Hz, 1H), 6.14 (s, 1H), 4.18 (s, 2H), 4.06 (s, 2H), 2.99 (s, 3H), 2.38 (s, 3H); MS (APCI+) m/z 426 (M+H)⁺.

Example 9

N-{3-[benzyl(4-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 8A and 4-fluorobenzaldehyde were processed as described in Example 1C to provide the title compound.

15 ¹H NMR (300 MHz, CDCl₃) δ 7.07-7.32 (m, 9H), 7.00-6.85 (m, 3H), 6.10 (s, 1H), 4.03 (s, 2H), 4.02 (s, 2H), 2.95 (s, 3H), 2.33 (s, 3H); MS (ESI+) m/z 399 (M+H)⁺; Analysis calculated for C₂₂H₂₃FN₂O₂S 0.75 TFA: C, 58.26; H, 4.95; N, 5.79. Found: C, 58.26; H, 5.17; N, 5.63.

Example 10

N-{3-[benzyl(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 8A and 2,4-difluorobenzaldehyde were processed as described in Example 1C to provide the title compound.

25 ¹H NMR (300 MHz, CDCl₃) δ 7.40-7.18 (m, 6H), 7.11 (m, 2H), 6.92 (m, 1H), 6.72 (m, 2H), 6.07 (s, 1H), 4.07 (s, 4H), 2.93 (s, 3H), 2.30 (s, 3H); MS(ESI+) m/z 417 (M+H)⁺.

Example 11

N-{3-[benzyl(2-cyanobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 1-(bromomethyl)-2-cyanobenzene were processed as described in Examples 6B-D to provide the title compound.

30 ¹H NMR (300 MHz, CDCl₃) δ 7.58 (d, J=7.2 Hz, 1H), 7.46 (m, 2H), 7.24 (m, 8H), 7.13 (t, J=8.0 Hz, 1H), 6.97 (dd, J=7.8, 1.0 Hz, 1H), 6.13 (s, 1H), 4.32 (s, 2H), 4.12 (s, 2H), 2.93 (s, 3H), 2.29 (s, 3H); MS (APCI+) m/z 406 (M+H)⁺.

Example 12

N-{3-[benzyl(4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

35 The product from Example 6A and 1-(chloromethyl)-4-methoxybenzene were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.22 (m, 6H), 7.11 (t, J=8.1 Hz, 1H), 7.08 (d, J=8.9 Hz, 2H),

6.92 (dd, J=8.2, 1.2 Hz, 1H), 6.78 (d, J=8.9 Hz, 2H), 6.10 (s, 1H), 4.10 (s, 2H), 4.05 (s, 2H), 3.77 (s, 3H), 2.92 (s, 3H), 2.29 (s, 3H); MS (APCI+) m/z 411 (M+H)⁺.

Example 13

5 N-{3-[benzyl(4-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 1-(bromomethyl)-4-bromobenzene were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.37 (d, J=8.1 Hz, 2H), 7.21 (m, 6H), 7.12 (t, J=8.0 Hz, 1H), 7.05 (d, J=8.1 Hz, 2H), 6.89 (dd, J=8.1, 1.0 Hz, 1H), 6.11 (s, 1H), 4.06 (s, 2H), 4.03 (s, 2H), 10 2.95 (s, 3H), 2.31 (s, 3H); MS (APCI+) m/z 460 (M+H)⁺.

Example 14

N-{3-[benzyl(3-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

15 The product from Example 6A and 1-(chloromethyl)-3-methoxybenzene were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 7H), 7.12 (t, J=8.0 Hz, 1H), 6.92 (dd, J=8.0, 1.1 Hz, 1H), 6.78 (m, 3H), 6.14 (s, 1H), 4.11 (s, 2H), 4.08 (s, 2H), 3.74 (s, 3H), 2.94 (s, 3H), 2.33 (s, 3H); MS (APCI+) m/z 411 (M+H)⁺.

Example 15

20 N-{3-[(1,3-benzodioxol-5-ylmethyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide

Example 15A

N-(1,3-benzodioxol-5-ylmethyl)-N-(2-methyl-3-nitrophenyl)amine

25 2-Methyl-3-nitroaniline and 1,3-benzodioxole-5-carbaldehyde were processed as described in Example 1C to provide the title compound.

Example 15B

N-{3-[(1,3-benzodioxol-5-ylmethyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide

30 The product from Example 15A and benzyl bromide were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.16-7.37 (m, 7H), 7.05 (d, J=8.1 Hz, 1H), 6.70 (s, 1H), 6.67 (s, 2H), 6.10 (s, 1H), 5.91 (s, 2H), 4.35 (s, 2H), 4.29 (s, 2H), 2.86 (s, 3H), 2.17 (s, 3H); MS (ESI+) m/z 425 (M+H)⁺; Analysis calculated for C₂₃H₂₄N₂O₄S 0.90 TFA: C, 56.51; H, 4.76; 35 N, 5.31. Found: C, 56.59; H, 4.68; N, 5.24.

Example 16

N-{3-[benzyl(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)amino]-2-methylphenyl}methanesulfonamide

Example 16A

5 N-(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)-N-(2-methyl-3-nitrophenyl)amine

2-Methyl-3-nitroaniline and 2,3-dihydro-1,4-benzodioxine-6-carbaldehyde were processed as described in Example 1C to provide the title compound.

Example 16B

10 N-{3-[benzyl(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 16A and benzyl bromide were processed as described in Examples 6B-D to provide the title compound.

15 ¹H NMR (300 MHz, CDCl₃) δ 7.30-7.25 (m, 6H), 7.10 (t, J=8.0 Hz, 1H), 6.89 (dd, J=7.8, 1.0 Hz, 1H), 6.75 (m, 2H), 6.66 (dd, J=8.1, 2.0 Hz, 1H), 6.11 (s, 1H), 4.23 (s, 4H), 4.04 (s, 2H), 3.93 (s, 2H), 2.95 (s, 3H), 2.35 (s, 3H); MS (ESI+) m/z 439 (M+H)⁺; Analysis calculated for C₂₄H₂₆N₂O₄S 0.25 CHCl₃: C, 62.19, H, 5.65; N, 5.98. Found: C, 62.18; H, 5.66; N, 5.88.

Example 17

20 N-{3-[benzyl(2-chlorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 1-(bromomethyl)-2-chlorobenzene were processed as described in Examples 6B-D to provide the title compound.

25 ¹H NMR (300 MHz, CDCl₃) δ 7.08-7.37 (m, 11H), 7.01 (dd, J=7.9, 1.2 Hz, 1H), 6.09 (s, 1H), 4.24 (s, 2H), 4.14 (s, 2H), 2.90 (s, 3H), 2.24 (s, 3H); MS (ESI+) m/z 415 (M+H)⁺; Analysis calculated for C₂₂H₂₃ClN₂O₂S 0.25 TFA: C, 60.94; H, 5.28; N, 6.32. Found: C, 60.99; H, 5.53; N, 6.17.

Example 18

30 N-{3-[benzyl(2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 8A and 2-fluorobenzaldehyde were processed as described in Example 1C to provide the title compound.

35 ¹H NMR (300 MHz, CDCl₃) δ 7.22 (m, 8H), 7.10 (t, J=8.0 Hz, 1H), 6.98 (m, 3H), 6.11 (s, 1H), 4.13 (s, 2H), 4.10 (s, 2H), 2.91 (s, 3H), 2.31 (s, 3H); MS (ESI+) m/z 399 (M+H)⁺; Analysis calculated for C₂₂H₂₃FN₂O₂S 0.20 TFA: C, 63.96; H, 5.55; N, 6.65. Found: C, 64.11; H, 5.69; N, 6.61.

Example 19N-{3-[[4-(allyloxy)benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide

5

Example 19AN-[4-(allyloxy)benzyl]-N-(2-methyl-3-nitrophenyl)amine

2-Methyl-3-nitroaniline and 4-allyloxybenzaldehyde were processed as described in Example 1C to provide the title compound.

10

Example 19BN-{3-[[4-(allyloxy)benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 19A and benzyl bromide were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 6H), 7.11 (t, J=8.0 Hz, 1H), 7.08 (d, J=8.5 Hz, 2H), 6.89 (dd, J=8.1, 1.0 Hz, 1H), 6.81 (d, J=8.8, Hz, 2H), 6.09 (s, 1H), 6.04 (m, 1H), 5.40 (ddd, J=17.3, 3.4, 1.7 Hz, 1H), 5.28 (ddd, J=10.5, 2.7, 1.4 Hz, 1H), 4.50 (dt, J=5.4, 1.4 Hz, 2H), 4.04 (s, 2H), 3.49 (s, 2H), 2.94 (s, 3H), 2.32 (s, 3H); MS (APCI+) m/z 437 (M+H)⁺.

15

Example 20N-{3-[[4-(allyloxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

20

Example 20AN-[4-(allyloxy)benzyl]-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 19A and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

25

Example 20BN-{3-[[4-(allyloxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

30

The product from Example 20A was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.20 (d, J=7.1 Hz, 1H), 7.11 (m, 4H), 6.89 (d, J=7.5 Hz, 1H), 6.82 (d, J=8.8 Hz, 2H), 6.71 (m, 2H), 6.15 (s, 1H), 6.02 (s, 1H), 5.40 (ddd, J=17.3, 3.1, 1.3 Hz, 1H), 5.28 (ddd, J=10.5, 2.7, 1.4 Hz), 4.51 (dt, J=5.4, 1.5 Hz), 4.05 (s, 2H), 3.99 (s, 2H), 2.93 (s, 3H), 2.30 (s, 3H); MS (APCI+) m/z 473 (M+H)⁺.

35

Example 21

N-{3-[(2-cyanobenzyl)(2-fluoro-4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

5

Example 21A

N-(2-fluoro-4-methoxybenzyl)-N-(2-methyl-3-nitrophenyl)amine

2-Methyl-3-nitroaniline and 2-fluoro-4-methoxy-benzaldehyde were processed as described in Example 1C to provide the title compound.

10

Example 21B

N-{3-[(2-cyanobenzyl)(2-fluoro-4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 21A and 2-(bromomethyl)benzonitrile were processed as described in Examples 6B-D to provide the title compound.

15 ¹H NMR (300 MHz, CDCl₃) δ 7.58 (d, J=7.8 Hz, 1H), 7.49 (m, 2H), 7.31 (m, 1H), 7.23 (dd, J=8.1, 1.0 Hz, 1H), 7.12 (t, J=8.0 Hz, 1H), 6.96 (m, 2H), 6.54 (m, 2H), 6.09 (s, 1H), 4.33 (s, 2H), 4.08 (s, 2H), 3.76 (s, 3H), 2.92 (s, 3H), 2.27 (s, 3H); MS (APCI+) m/z 454 (M+H)⁺.

Example 22

20 N-{3-[(2,4-difluorobenzyl)(4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 22A

N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

25 2-Methyl-3-nitroaniline and 2,4-difluorobenzaldehyde were processed as described in Example 1C to provide the title compound.

Example 22B

N-{3-[(2,4-difluorobenzyl)(4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

30 The product from Example 22A and 4-methoxybenzyl chloride were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.21 (dd, J=8.0, 1.0 Hz, 1H), 7.12 (m, 4H), 6.91 (d, J=8.1 Hz, 1H), 6.79 (d, J=8.8 Hz, 2H), 6.71 (m, 2H), 6.10 (s, 1H), 4.10 (s, 2H), 4.04 (s, 2H), 3.78 (s, 3H), 2.93 (s, 3H), 2.28 (s, 3H); MS (APCI+) m/z 447 (M+H)⁺.

35

Example 23

N-{3-[(2,4-difluorobenzyl)(2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 22A and 2-fluorobenzyl bromide were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 4H), 7.11 (t, J=8.1 Hz, 1H), 6.94 (m, 3H), 6.72 (m, 2H), 6.13 (s, 1H), 4.15 (s, 2H), 4.13 (s, 2H), 2.91 (s, 3H), 2.25 (s, 3H); MS (APCI+) m/z 435 (M+H)⁺.

Example 24

N-{3-[bis(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 22A and 2,4-difluorobenzyl bromide were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 1H), 7.14 (m, 3H), 6.99 (dd, J=8.1, 1.0 Hz, 1H), 6.72 (m, 4H), 6.14 (s, 1H), 4.09 (s, 4H), 2.93 (s, 3H), 2.25 (s, 3H); MS (APCI+) m/z 453 (M+H)⁺.

Example 25

N-{3-[(2-cyanobenzyl)(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 22A and 2-(bromomethyl)benzonitrile were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (CDCl₃, 300 MHz) δ 7.59 (m, 1H), 7.48 (m, 2H), 7.32 (m, 1H), 7.24 (m, 1H), 7.11 (m, 2H), 6.95 (m, 1H), 6.10 (s, 1H), 4.32 (s, 2H), 4.13 (s, 2H), 2.93 (s, 3H), 2.26 (s, 3H); MS (APCI+) m/z 442 (M+H)⁺.

Example 26

N-{3-[benzyl(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 26A

N-(2-methyl-3-nitrophenyl)-N-(4-phenoxybenzyl)amine

2-Methyl-3-nitroaniline and 4-phenoxybenzaldehyde were processed as described in Example 1C to provide the title compound.

Example 26B

N-{3-[benzyl(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 26A and benzyl bromide were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.33 (m, 2H), 7.21 (m, 6H), 7.14 (m, 4H), 6.97 (m, 3H), 6.88 (d, J=8.5 Hz, 2H), 6.10 (s, 1H), 4.18 (s, 2H), 4.15 (s, 2H), 2.91 (s, 3H), 2.27 (s, 3H); MS (ESI+) m/z 473 (M+H)⁺.

Example 27N-{3-[benzyl(2-methylbenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 1-(bromomethyl)-2-methylbenzene were processed as described in Examples 6B-D to provide the title compound.

5 ¹H NMR (500 MHz, DMSO-d₆) δ 2.16 (s, 3H), 2.28 (s, 3H), 2.88 (s, 3H), 4.06 (d, 4H, J=17.8 Hz), 7.03 (m, 6H), 7.24 (m, 6H), 8.91 (s, 1H); MS (APCI+) m/z 395 (M+H)⁺.

Example 28N-{3-[benzyl(4-methylbenzyl)amino]-2-methylphenyl}methanesulfonamide

10 The product from Example 6A and 1-(bromomethyl)-4-methylbenzene were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (500 MHz, DMSO-d₆) δ 2.24 (s, 3H), 2.39 (s, 3H), 2.91 (s, 3H), 4.01 (d, 4H, J=18.7 Hz), 6.90-7.30 (m, 12H), 8.94 (s, 1H); MS (APCI+) (M+H)⁺ at m/z 395.

Example 29N-{3-[benzyl(4-chlorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 4-chlorobenzyl bromide were processed as described in Examples 6B-D to provide the title compound.

15 ¹H NMR (500 MHz, DMSO-d₆) δ 2.39 (s, 3H), 2.92 (s, 3H), 4.05 (br.s, 4H), 6.98 (m, 3H), 7.27 (m, 9H), 8.95 (s, 1H); MS (APCI+) m/z 415 (M+H)⁺.

Example 30N-(3-{benzyl[4-(trifluoromethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide

25 The product from Example 6A and 4-trifluoromethylbenzyl bromide were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (500 MHz, DMSO-d₆) δ 2.40 (s, 3H), 2.92 (s, 3H), 4.07 (s, 2H), 4.16 (s, 2H), 7.00 (m, 3H), 7.25 (m, 5H), 7.49 (d, 2H, J=8.1 Hz), 7.63 (d, 2H, J=8.1 Hz), 8.95 (s, 1H); MS (APCI+) m/z 448 (M+H)⁺.

Example 31N-(3-{benzyl[2-fluoro-4-(trifluoromethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 6A and 2-fluoro-4-trifluoromethylbenzyl bromide were processed as described in Examples 6B-D to provide the title compound.

35 ¹H NMR (500 MHz, DMSO-d₆) δ 2.34 (s, 3H), 2.88 (s, 3H), 4.11 (s, 2H), 4.21 (s, 2H), 6.98 (m, 2H), 7.05 (t, 1H, J=8.0 Hz), 7.22 (m, 1H), 7.28 (d, 4H, J=4.4 Hz), 7.50 (m, 2H), 7.56 (d, 1H, J=9.4 Hz), 8.96 (s, 1H); MS (APCI+) m/z 467 (M+H)⁺.

Example 32N-{3-[benzyl(2,4-dichlorobenzyl)amino]-2-methylphenyl}methanesulfonamideExample 32A2,4-dichloro-1-(iodomethyl)benzene

2,4-Dichloro-1-(chloromethyl)benzene (0.8 mmoles) in acetone (2 mL) was treated with NaI (0.48 g) and stirred at room temperature overnight. The solvent was removed under reduced pressure and the residue extracted with DMF to provide the title compound.

Example 32BN-{3-[benzyl(2,4-dichlorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and the product from Example 32A were processed as described in Examples 6B-D to provide the title compound.

¹H NMR (500 MHz, DMSO-d₆) δ 2.32 (s, 3H), 2.89 (s, 3H), 4.12 (s, 2H), 4.16 (s, 2H), 6.98 (t, 2H, J=8.0 Hz), 7.05 (t, 1H, J=7.8 Hz), 7.24 (m, 5H), 7.33 (dd, 1H, J=8.2, 2.0 Hz), 7.40 (d, 1H, J=8.1 Hz), 7.52 (d, 1H, J=1.9 Hz), 8.83-9.05 (br.s, 1H); MS (APCI+) m/z 450 (M+H)⁺.

Example 33N-{3-[(2,4-difluorobenzyl)(4-{[3-phenyl-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamideExample 33AN-(2,4-difluorobenzyl)-N-(4-hydroxybenzyl)-2-methyl-3-nitroaniline

The product from Example 20A (1.19 g, 2.81 mmoles) and Pd(PPh₃)₄ (0.16 g, 0.14 mmoles) in CH₂Cl₂ (11.2 mL) was treated with phenylsilane (0.7 mL, 5.63 mmoles) and allowed to stir for 2 hours at room temperature. The residue was purified by flash chromatography (silica gel, 4:1 hexanes:ethyl acetate) to provide the title compound as a yellow oil (1.03 g, 96%).

Example 33BN-(2,4-difluorobenzyl)-2-methyl-3-nitro-N-(4-{[3-phenyl-2-propenyl]oxy}benzyl)aniline

The product from Example 33A (0.1 g, 26 mmoles) in DMF (0.65 mL) was treated with sodium hydride (0.011 g, 0.26 mmoles). After 10 minutes, the reaction mixture was treated with cinnamyl bromide (0.10 g, 0.52 mmoles) and shaken overnight. The mixture was diluted with diethyl ether (50 mL), washed with saturated ammonium chloride (50 mL)

and water (2x50 mL), brine (50 mL), dried (Na₂SO₄), filtered and the filtrate concentrated under reduced pressure to provide the title compound which was used in the next step without further purification.

5 Example 33C

N-{3-[(2,4-difluorobenzyl)(4-{[3-phenyl-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfona

The crude product from Example 33B was processed as described in Examples 6C and 6D to provide the title compound as a brown solid (0.020 g, 14%).

10 ¹H NMR (300 MHz, CDCl₃) δ 7.50 (m, 2H), 7.33 (m, 2H), 7.21 (m, 2H), 7.13 (m, 3H), 6.94 (d, J=7.8 Hz, 1H), 6.85 (d, J=8.8 Hz, 2H), 6.21 (m, 2H), 6.39 (dt, J=15.9, 5.8 Hz, 1H), 6.09 (s, 1H), 4.67 (dd, J=5.9, 1.5 Hz, 2H), 4.15 (s, 2H), 4.08 (s, 2H), 2.91 (s, 3H), 2.26 (s, 3H); MS (APCI+) m/z 549 (M+H)⁺.

15 Example 34

N-{3-[[4-(benzyloxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 34A

20 N-[4-(benzyloxy)benzyl]-N-(2,4-difluorobenzyl)-2-methyl-3-nitroaniline

The product from Example 33A and benzyl bromide were processed as described in Example 33B to provide the title compound.

Example 34B

25 N-{3-[[4-(benzyloxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 34A was processed as described in Examples 6C and 6D to provide the title compound.

30 ¹H NMR (300 MHz, CDCl₃) δ 7.38 (m, 3H), 7.23 (m, 1H), 7.13 (m, 3H), 6.97 (d, J=7.1 Hz, 1H), 6.86 (d, J=8.8 Hz, 2H), 6.73 (m, 2H), 6.08 (s, 1H), 5.02 (s, 2H), 4.22 (s, 2H), 4.14 (s, 2H), 2.89 (s, 3H), 2.21 (s, 3H); MS(ESI+) m/z 523 (M+H)⁺.

Example 35

35 N-{3-[benzyl(4-{[3-bromo-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

Example 35AN-(4-{[3-bromo-2-propenyl]oxy}benzyl)-N-(2,4-difluorobenzyl)-2-methyl-3-nitroaniline

The product from Example 33A and 1,3-dibromo-1-propene were processed as described in Example 33B to provide the title compound.

5

Example 35BN-{3-[benzyl(4-{[3-bromo-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 35A was processed as described in Examples 6C and 6D to provide the title compound.

10

¹H NMR (300 MHz, CDCl₃) δ 7.24 (m, 5H), 7.19 (d, J=7.8 Hz, 2H), 7.08 (d, J=8.8 Hz, 2H), 6.87 (m, 1H), 6.78 (d, J=8.8 Hz, 1H), 6.44 (m, 2H), 6.14 (s, 1H), 4.45 (d, J=4.4 Hz, 2H), 4.03 (s, 2H), 3.98 (s, 2H), 2.95 (s, 3H), 2.34 (s, 3H); MS (APCI+) m/z 515 (M)⁺.

15

Example 36N-{3-[bis(2-bromobenzyl)amino]-4-methoxyphenyl}methanesulfonamideExample 36AN,N-bis(2-bromobenzyl)-N-(2-methoxy-5-nitrophenyl)amine

20

2-Methoxy-5-nitroaniline and 1-bromo-2-(bromomethyl)benzene were processed as described in Example 2A to provide the title compound.

Example 36BN-{3-[bis(2-bromobenzyl)amino]-4-methoxyphenyl}methanesulfonamide

25

The product from Example 36A was processed as described in Examples 6C and 6D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.55-7.45 (m, 4H), 7.22 (td, J=7.5, 1.1 Hz, 2H), 7.07 (td, J=7.7, 1.7 Hz, 2H), 6.90 (dd, J=8.6, 2.5 Hz, 1H), 6.82 (d, J=8.8 Hz, 1H), 6.60 (d, J=2.0 Hz, 1H), 6.03 (bs, 1H), 4.48 (s, 4H), 3.80 (s, 3H), 2.58 (s, 3H); MS (ESI+) m/z 555 (M+H)⁺;

30

Analysis calculated for C₂₂H₂₂Br₂N₂O₃S 0.40 TFA: C, 45.65; H, 3.76; N, 4.67. Found: C, 45.66; H, 3.84; N, 4.67.

Example 37N-(3-{(2-bromobenzyl)[cyano(phenyl)methyl]amino}-2-methylphenyl)methanesulfonamide

35

The major product from Example 1C (0.21 g), potassium cyanide (0.10 g, 1.06 mmoles), and benzaldehyde (0.14 mL, 1.41 mmoles) in methanol (1.86 mL) and acetonitrile

(1.0 mL) were treated with acetic acid (0.86 mL). The reaction was shaken vigorously overnight and was then concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 4:1→1:1 hexanes:ethyl acetate) to provide the title compound as a brown solid (0.22 g, 83%). A portion of the product (0.1 g) was repurified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column. ¹H NMR (300 MHz, CDCl₃) δ 7.66 (dd, J=7.1, 2.0 Hz, 1H), 7.50 (d, J=7.1 Hz, 1H), 7.32 (m, 7H), 7.08 (m, 3H), 5.91 (s, 1H), 5.25 (s, 1H), 4.56 (d, J=13.6 Hz, 1H), 4.21 (d, J=13.6 Hz, 1H), 2.78 (s, 3H), 1.84 (s, 3H); MS (APCI+) m/z 486 (M+H)⁺.

Example 38

N-(3-{benzyl[4-(methoxymethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide

Example 38A

N-benzyl-N-(4-hydroxymethylbenzyl)-2-methyl-3-nitroaniline

The product from Example 6B (0.55 g, 1.4 mmoles) in THF (3.5 mL) was treated with diisobutylaluminum hydride (DIBAL) (2.94 mL, 2.94 mmoles, 1.0M in hexanes) dropwise at -78 °C. After 30 minutes, an additional portion of DIBAL (1.5 mL) was added dropwise. After an additional 30 minutes, the acetone/dry ice bath was removed and the reaction mixture was carefully quenched with saturated NH₄Cl. The mixture was diluted with diethyl ether (200 mL) and saturated sodium potassium tartrate (250 mL) and stirred vigorously for 2.5 hours. The two phases were separated and the organic phase was washed with brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 4:1→1:1 hexanes:ethyl acetate) to provide the title compound as a yellow oil (0.46 g, 91%).

Example 38B

N-benzyl-N-[4-(methoxymethyl)benzyl]-2-methyl-3-nitroaniline

The product from Example 38A (0.1 g, 0.276 mmoles) in DMF (0.7 mL) was treated with sodium hydride (0.012 g, 0.3 mmoles, 60% dispersion). After 10 minutes, the reaction mixture was treated with iodomethane (0.02 mL, 0.36 mmoles) and then stirred overnight at room temperature. The mixture was diluted with diethyl ether (100 mL), washed with saturated ammonium chloride (50 mL) and water (50 mL) twice, washed with brine (50 mL), dried (Na₂SO₄), filtered and concentrated under reduced pressure to provide the title compound which was used in the next step without further purification.

Example 38C

N-(3-{benzyl[4-(methoxymethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide

The crude product from Example 38B was processed as described in Examples 6C and 6D to provide the title compound as a brown oil (0.056 g, 48%).

¹H NMR (300 MHz, CDCl₃) δ 7.20 (m, 10H), 7.10 (t, J=8.0 Hz, 1H), 6.89 (d, J=7.5 Hz, 1H), 6.10 (s, 1H), 4.41 (s, 2H), 4.05 (s, 4H), 3.39 (s, 3H), 2.95 (s, 3H), 2.34 (s, 3H); MS (APCI+) m/z 425 (M+H)⁺.

Example 39

N-(3-{benzyl[4-(hydroxymethyl)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 6D and diisobutylaluminum hydride were processed as described in Example 38A to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 10H), 7.97 (t, J=8.0 Hz, 1H), 6.88 (dd, J=8.5, 1.0 Hz, 1H), 6.12 (s, 1H), 4.67 (d, J=8 Hz, 2H), 4.05 (s, 2H), 4.03 (s, 2H), 2.96 (s, 3H), 2.37 (s, 3H); MS (APCI+) m/z 411 (M+H)⁺.

Example 40

N-[3-(dibenzylamino)-2-((E)-3-ethoxy-3-oxo-1-propenyl)phenyl)methanesulfonamide

Example 40A

N,N-dibenzyl-N-[2-(1,3-dioxolan-2-yl)-3-nitrophenyl]amine

Benzyl bromide (2.14 mL, 17.98 mmoles) was added to a solution of 2-(1,3-dioxolan-2-yl)-3-nitroaniline (prepared according to Wall, M. E.; Wani, M. C.; Nicholas, A. W.; Manikumar, G.; Tele, C.; Moore, L.; Truesdale, A.; Leitner, P.; Besterman, J. M.; J. Med. Chem. 1993, 36, 2689) (1.888 g, 8.99 mmoles) and diisopropylethylamine (3.1 mL, 17.98 mmoles) in DMF (18.0 mL) and the reaction was heated at 85 °C overnight. After cooling to room temperature, the mixture was diluted with diethyl ether (250 mL), washed with saturated ammonium chloride (200 mL), water (2x200 mL), brine (150 mL), dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 4:1 hexanes:ethyl acetate) to provide the title compound (1.51 g, 43%) as a black oil.

Example 40B

N,N-dibenzyl-N-[2-(formyl)-3-nitrophenyl]amine

The product from Example 40A (1.51 g) was processed as described in Example 6C to provide the title compound (0.76 g, 72%).

Example 40C

ethyl (2E)-3-[2-amino-6-(dibenzylamino)phenyl]-2-propenoate

Sodium hydride (0.046 g, 1.14 mmol, 60% dispersion) was added to a solution of triethylphosphonoacetate (0.23 mL, 1.14 mmol) in THF (1.4 mL) at 0 °C. After 15 minutes, the product from Example 40B (0.3 g, 0.95 mmol) in THF (1.4 mL) was added dropwise and the reaction mixture was stirred overnight. The mixture was diluted with diethyl ether (150 mL), washed with water (100 mL), brine (100 mL), dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 4:1 hexanes:ethyl acetate) to provide the title compound (0.29 g, 78%).

Example 40D

N-[3-(dibenzylamino)-2-((E)-3-ethoxy-3-oxo-1-propenyl)phenyl]methanesulfonamide

The product from Example 40C was processed as described in Example 6D to provide the title compound as a brown solid. A portion of the product (0.1 g) was purified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column.

¹H-NMR (300 MHz, CDCl₃) δ 8.06 (d, J=17.0 Hz), 7.22 (m, 12H), 6.85 (dd, J=7.8, 1.0 Hz, 1H), 6.78 (s, 1H), 6.27 (d, J=17.0, 1H), 4.29 (s, 4H), 2.99 (s, 3H); MS (APCI+) m/z 465 (M+H)⁺.

Example 41

N-(3-{benzyl[4-(2-hydroxyethoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

Example 41A

N-(3-{benzyl[4-(2,3-dihydroxypropoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 19B (2.93 g, 6.72 mmol) and 4-methyl-morpholine N-oxide (0.8660 g, 7.4 mmol) in acetone (30 mL) and water (4.3 mL) were treated with osmium tetroxide (1.7 mL, 0.0168 mmol, 2.5 wt % solution in 2-methyl-2-propanol) and allowed to stir at room temperature overnight. The mixture was then treated with 1,4-Diazabicyclo[2.2.2]octane (0.05 g) and 4-methyl-morpholine N-oxide (0.8660 g, 7.4 mmol) and allowed to stir for an additional 24 hours. Saturated NaHSO₃ (250 mL) and ethyl acetate (250 mL) were added and the reaction mixture was stirred vigorously for 2.5 hours. The mixture was partitioned and the organic phase washed with brine, dried (Na₂SO₄), filtered and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, ethyl acetate) to provide the title compound as a yellow oil (3.0 g, 95%).

Example 41B

N-(3-{benzyl[4-(2-hydroxyethoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 41A (0.6 g, 1.27 mmoles) in benzene (11.6 mL) and ethanol (11.6 mL) was treated with lead tetraacetate (0.6215 g, 1.4 mmoles) at 0 °C. The reaction mixture turned from clear and colorless to light yellow and was stirred for 15 minutes. The mixture was diluted with diethyl ether (100 mL), filtered through a 1 inch pad of 1:1 Celite:silica gel and the filtrate was concentrated under reduced pressure. The residue (0.07 g, .16 mmoles) was dissolved in ethanol (1.0 mL) and treated with sodium borohydride (0.006 g, 0.16 mmoles). After 30 minutes, the reaction mixture was filtered through silica gel using 2:1 ethyl acetate:hexanes and concentrated under reduced pressure. The residue was purified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column to provide the title compound as a brown oil (0.055 g).

¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 10H), 6.75 (dd, J=8.8, 1.7 Hz, 2H), 6.04 (s, 1H), 4.66 (dd, J=5.4, 3.7 Hz, 1H), 4.57 (d, J=10.3 Hz, 2H), 4.52 (d, J=10.3 Hz, 2H), 4.21 (m, 1H), 4.02 (m, 1H), 3.95 (m, 1H), 3.72 (s, 1H), 2.76 (s, 3H), 1.98 (s, 3H); MS (APCI+) m/z 441 (M+H)⁺.

Example 42

N-[3-(dibenzylamino)-2-(hydroxymethyl)phenyl]methanesulfonamide

Example 42A

N-[3-(dibenzylamino)-2-formylphenyl]methanesulfonamide

The product from Example 40B was processed as described in Example 6D to provide the title compound.

Example 42B

N-[3-(dibenzylamino)-2-(hydroxymethyl)phenyl]methanesulfonamide

The product from Example 42A (0.017 g, .043 mmoles) in ethanol (2 mL) was treated with sodium borohydride (0.002 g, 0.053 mmoles). After stirring two hours at room temperature, the reaction mixture was quenched with saturated ammonium chloride and extracted with chloroform. The extracts were combined, dried with sodium sulfate, filtered and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 30% ethyl acetate/hexanes) to provide the title compound (0.015 g, 88%).

¹H NMR (300 MHz, CDCl₃) δ 7.63 (s, 1H), 7.12-7.38 (m, 13H), 4.67 (s, 2H), 4.06 (s, 4H), 2.85 (s, 3H); MS (ESI+) m/z 397 (M+H)⁺.

Example 43

N-{3-[(2,4-difluorobenzyl)(4-propoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 43AN-(2,4-difluorobenzyl)-2-methyl-3-nitro-N-(4-propoxybenzyl)aniline

The product from Example 33A and 1-iodopropane were processed as described in Example 33B to provide the title compound.

5

Example 43BN-{3-[(2,4-difluorobenzyl)(4-propoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 43A was processed as described in Examples 6C and 6D to provide the title compound.

10

¹H NMR (300 MHz, CDCl₃) δ 7.30-7.10 (m, 4H), 6.96 (m, 1H), 6.63-6.83 (m, 5H), 6.07 (s, 1H), 3.96-4.32 (m, 4H), 3.87 (t, J=6.6 Hz, 2H), 2.88 (s, 3H), 2.26 (s, 3H), 1.78 (m, 2H), 1.02 (t, J=7.5 Hz, 3H); MS (ESI+) m/z 475 (M+H)⁺.

Example 44

15

N-[3-(dibenzylamino)-2-vinylphenyl]methanesulfonamideExample 44A1,3-dinitro-2-vinylbenzene

1-Chloro-2,6-dinitrobenzene (1.00 g, 4.94 mmol, purchased from Lancaster) tris(dibenzylideneacetone)dipalladium (0.113 g, 0.123 mmol), tri-2-furylphosphine (0.229 g, 0.987 mmol), copper(I) iodide (0.094 g, 0.494 mmol), and lithium chloride (0.628 g, 14.8 mmol) in N,N-dimethylformamide (15 mL) were treated with tributylethylstannane (2.90 mL, 9.87 mmol). The reaction mixture was degassed with nitrogen, stirred overnight at room temperature and then heated at 80 °C for 4 hours. The reaction mixture was then diluted with ethyl acetate and washed with water and brine. The organic phase was dried with sodium sulfate, filtered and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 10% ethyl acetate/hexanes) to provide the title compound (0.669 g, 70%). MS (DCI) m/z 194 (M+H)⁺.

20

25

30

Example 44B3-nitro-2-vinylaniline

The product from Example 44A (0.669 g, 3.45 mmol) in 3:1 ethanol:water (40 mL) was treated with sodium sulfide nonahydrate (1.66 g, 6.90 mmol) and heated at reflux for 30 minutes. The mixture was allowed to cool to room temperature and then concentrated under reduced pressure. The residue was mixed with water and extracted with chloroform (2X). The chloroform extracts were combined, washed with brine, dried with sodium sulfate and the filtrate concentrated under reduced pressure. The residue was

35

purified by flash chromatography (silica gel, 25% ethyl acetate/hexanes) to provide the title compound (0.137 g, 24%).

MS (DCI) m/z 165 (M+H)⁺.

5

Example 44C

N,N-dibenzyl-N-(3-nitro-2-vinylphenyl)amine

The product from Example 44B and benzyl bromide were processed as described in Example 2A to provide the title compound.

10

Example 44D

N-[3-(dibenzylamino)-2-vinylphenyl]methanesulfonamide

The product from Example 44C was processed as described in Examples 6C and 6D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.10-7.32 (m, 13H), 6.96 (dd, J=18.6, 11.6 Hz, 1H), 6.75 (dd, J=8.2, 1.0 Hz, 1H), 5.77 (dd, J=11.5, 1.4 Hz, 1H), 5.51 (dd, J=18.5, 1.2 Hz, 1H), 2.95 (s, 4H), 1.53 (s, 3H); MS (ESI+) m/z 393 (M+H)⁺.

15

Example 45

N-[3-(dibenzylamino)-2-ethylphenyl]methanesulfonamide

20

The product from Example 44D (0.040g, 0.102 mmoles) and p-toluenesulfonylhydrazide (0.150 g, 1.02 mmoles) in ethylene glycol dimethyl ether (5 mL) at reflux were treated dropwise over a four hour period with a solution of sodium acetate trihydrate (0.232 g, 1.70 mmoles) in water. After the addition, the reaction mixture was allowed to cool to room temperature, poured into water and extracted with methylene chloride (3x). The extracts were combined, dried with sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative HPLC (CH₃CN:0.1% TFA in H₂O) on a YMC ODS Guardpak column.

25

¹H NMR (300 MHz, CDCl₃) δ 7.32 (dd, J=8.0, 1.2 Hz, 1H), 7.35-7.15 (m, 11H), 7.09 (dd, J=8.0, 1.2 Hz, 1H), 6.14 (s, 1H), 4.09 (s, 4H), 2.85 (s, 3H), 2.74 (q, J=7.6 Hz, 2H), 1.00 (t, J=7.6 Hz, 3H); MS (ESI+) m/z 395 (M+H)⁺.

30

Example 46

N-[3-(dibenzylamino)-2-(methoxymethyl)phenyl]methanesulfonamide

35

Example 46A

N,N-dibenzyl-2-formyl-3-nitroaniline

The product from Example 40A (1.88 g, 4.83 mmoles) in tetrahydrofuran (30 mL)

was treated with an aqueous solution of 2N sulfuric acid (1.24 mL) and allowed to stir overnight at 65 °C. The mixture was cooled to room temperature, quenched with saturated sodium bicarbonate, and concentrated under reduced pressure. The residue was mixed with diethylether and washed with water, brine, dried with sodium sulfate, filtered, and the filtrate was concentrated under reduced pressure to provide the title compound. MS (ESI+) at m/z 347 (M+H)⁺.

Example 46B

N,N-dibenzyl-2-hydroxymethyl-3-nitroaniline

The product from Example 46A (1.5 g, 4.33 mmoles) in ethanol (10 mL) and tetrahydrofuran (4 mL) was treated with sodium borohydride (0.164 g, 4.33 mmoles). After stirring for 30 minutes at room temperature, the mixture was quenched with saturated ammonium chloride and diluted with diethyl ether. The diethyl ether was washed with saturated ammonium chloride, brine, dried with sodium sulfate, filtered and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 10:1 to 4:1 hexanes:ethyl acetate) to provide the title compound (1.39 g, 92%). MS (ESI+) m/z 348 (M+H)⁺.

Example 46C

N,N-dibenzyl-2-(methoxymethyl)-3-nitroaniline

The product from Example 46B (0.12 g, 0.33 mmoles) in anhydrous N,N-dimethylformamide (1 mL) was treated with 60% sodium hydride (0.016 g, 0.40 mmoles). After stirring for 10 minutes at room temperature, the mixture was treated with iodomethane (0.031 mL, 0.50 mmoles) and allowed to stir overnight at room temperature. The reaction mixture was quenched with saturated ammonium chloride and extracted with ethyl acetate (3X). The ethyl acetate phases were combined, dried with sodium sulfate, filtered and the filtrate concentrated under reduced pressure to provide the title compound. MS (ESI+) m/z 363 (M+H)⁺.

Example 46D

N-[3-(dibenzylamino)-2-(methoxymethyl)phenyl]methanesulfonamide

The product from Example 46C was processed as described in Examples 6C and 6D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.99 (s, 1H), 7.32 (d, J=8.1 Hz, 1H), 7.30-7.10 (m, 11H), 6.96 (d, J=7.7 Hz, 1H), 4.64 (s, 2H), 4.09 (s, 4H), 3.25 (s, 3H), 2.90 (s, 3H); MS (ESI+) m/z 411 (M+H)⁺.

Example 47N-[3-(dibenzylamino)-2-(ethoxymethyl)phenyl]methanesulfonamideExample 47AN,N-dibenzyl-2-(ethoxymethyl)-3-nitroaniline

The product from Example 46B and iodoethane were processed as described in Example 46C to provide the title compound.

Example 47BN-[3-(dibenzylamino)-2-(ethoxymethyl)phenyl]methanesulfonamide

The product from Example 47A was processed as described in Examples 6C and 6D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 8.17 (s, 1H), 7.34 (d, J=7.8 Hz, 1H), 7.10-7.31 (m, 11H), 6.98 (d, J=8.4 Hz, 1H), 4.66 (s, 2H), 4.14 (bs, 4H), 3.39 (q, J=7.0 Hz, 2H), 2.87 (s, 3H), 1.17 (t, J=6.9 Hz, 3H); MS (ESI+) m/z 425 (M+H)⁺.

Example 48N-{3-[[4-(4-bromophenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamideExample 48AN-[4-(4-bromophenoxy)benzyl]-N-(2,4-difluorobenzyl)-2-methyl-3-nitroaniline

The product from Example 33A (1.0 g, 2.6 mmol), 4 Å molecular sieves, Cu(OAc)₂ (0.71 g, 3.91 mmol) and 4-bromophenylboronic acid (1.57 g, 7.81 mmol) in CH₂Cl₂ (20 mL) was treated with triethylamine (1.81 mL, 13.0 mmol) and the resulting solution was stirred vigorously overnight. An additional equivalent of Cu(OAc)₂, 4-bromophenylboronic acid, and triethylamine were added and the reaction mixture was stirred vigorously for 5 hours. The mixture was filtered through a 1 inch pad of Celite using CH₂Cl₂ and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 10:1 hexanes:ethyl acetate) to provide the title compound as a yellow oil (0.87 g, 62%).

Example 48BN-{3-[[4-(4-bromophenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 48A was processed as described in Examples 6C and 6D to provide the title compound.

¹H NMR (300 MHz, CDCl₃) δ 7.42 (m, 2H), 7.20 (m, 6H), 7.05-7.19 (m, 2H), 6.90 (d, J=8.8 Hz, 2H), 6.84 (m, 2H), 6.68-6.76 (m, 2H), 6.11 (s, 1H), 4.07 (s, 2H), 4.05 (s, 2H), 2.96 (s, 3H), 2.32 (s, 3H); MS(ESI+) (M)⁺ at m/z 587.

5

Example 49

N-{3-[(4-(4-(3-ethoxy-3-oxopropyl)phenoxy)benzyl)(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 48B (0.14 g, 0.23 mmol) and Pd(PPh₃)₄ (0.027 g, 0.023 mmol) was treated with 2-ethoxycarbonyl ethyl zinc bromide (0.5M in THF, 1.87 mL, 0.93 mmol, purchased from Aldrich). The resulting solution was degassed and heated at 75 °C overnight. The mixture was filtered through a 1 inch pad of Celite using diethyl ether and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 4:1 → 1:1 hexanes:ethyl acetate) to provide the title compound (0.11 g, 77%).

¹H NMR (300 MHz, CDCl₃) δ 7.06-7.22 (m, 8H), 6.86-6.93 (m, 4H), 6.69-6.77 (m, 2H), 6.09 (s, 1H), 4.12 (q, J=7.1 Hz, 2H), 4.07 (s, 2H), 4.03 (s, 2H), 2.95 (s, 3H), 2.93 (t, J=8.1 Hz, 2H), 2.61 (t, J=8.1 Hz, 2H), 2.31 (s, 3H), 1.24 (t, J=7.1 Hz, 3H); MS (APCI+) (M+H)⁺ at m/z 609.

20

Example 50

N-{3-[(4-(4-(2-carboxyethyl)phenoxy)benzyl)(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 49 was treated with aqueous sodium hydroxide in methanol and allowed to stir overnight. The mixture was concentrated under reduced pressure, diluted with water and extracted with diethyl ether. The aqueous phase was acidified with 2N aqueous hydrochloric acid and extracted with ethyl acetate. The ethyl acetate phase was dried over sodium sulfate, filtered and the filtrate concentrated under reduced pressure to provide the title compound.

¹H NMR (CDCl₃, 300 MHz) δ 7.09-7.23 (m, 8H), 6.87-6.94 (m, 4H), 6.68-6.77 (m, 2H), 6.14 (s, 1H), 4.10 (s, 2H), 4.05 (s, 2H), 2.95 (t, J=7.6 Hz, 2H), 2.94 (s, 3H), 2.68 (t, J=7.6 Hz, 2H), 2.28 (s, 3H); MS (APCI+) (M+H)⁺ at m/z 581.

35

Example 51

N-{3-[(2,4-difluorobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 26A and 2,4-difluorobenzyl bromide were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ

7.33 (m, 2 H), 7.15 (m, 6 H), 6.94 (m, 5 H), 6.72 (m, 2 H), 6.10 (s, 1 H), 4.10 (s, 2 H), 4.06 (s, 2 H), 2.95 (s, 3 H), 2.30 (s, 3 H); MS (APCI) m/z 509 ($M+H^+$).

5

Example 52

methyl 4-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate

10

Example 52A

methyl 4-(4-{[(2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino)methyl]phenoxy}benzoate

15

The product from Example 33A (.1520 g, .396 mmoles), methyl-4-bromobenzoate (.1702 g, .792 mmoles), K_3PO_4 (.1680 g, .792 mmoles), $Pd(OAc)_2$ (.0070 g, .032 mmoles), and 2-(di-*tert*-butylphosphino)biphenyl (.0140 g, .048 mmoles) was treated with toluene (1.16 mL) under a nitrogen atmosphere. The resulting red solution was degassed, stirred for 10 min, and heated to 100 °C overnight. The black solution was diluted with diethyl ether, extracted with sat. NH_4Cl and sat. $NaCl$, dried (Na_2SO_4), and concentrated *under reduced pressure*. The residue was purified by flash chromatography (10:1 hexane:ethyl acetate) on silica gel to provide the title compound (.1886 g, 92%) as a yellow oil.

20

Example 52B

methyl 4-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate

25

The product from Example 52A was processed as described in Examples 6C and D to provide the title compound. 1H NMR (300 MHz, $CDCl_3$) δ 8.01 (m, 2 H), 7.22 (m, 3 H), 7.11 (m, 2 H), 6.95 (m, 5 H), 6.73 (m, 2 H), 6.08 (s, 1 H), 4.11 (s, 2 H), 4.09 (s, 2 H), 3.90 (s, 3 H), 2.96 (s, 3 H), 2.32 (s, 3 H); MS (APCI) m/z 567 ($M+H^+$).

30

N-{3-[(2,4-difluorobenzyl)(2-fluoro-4-phenoxybenzyl)amino]-2-
methylphenyl}methanesulfonamide

Example 53A

4-(allyloxy)-2-fluorobenzonitrile

35

2-Fluoro-4-hydroxybenzonitrile (5.00 g, 36.5 mmoles) and allyl bromide (3.47 mL,

40.1 mmol) in anhydrous DMF (40 mL) were treated with K_2CO_3 (10.0 g, 72.9 mmol) and stirred overnight at 80°C. Reaction mixture cooled and diluted with ethyl acetate, washed with H_2O (2X) and brine, dried (Na_2SO_4), filtered, and concentrated under reduced pressure to provide the title compound with no further purification.

5

Example 53B

4-(allyloxy)-2-fluorobenzaldehyde

The product from Example 53A (6.42 g, 36.5 mmol) in anhydrous THF (50 mL) was treated with diisobutylaluminum hydride (DIBAL) (37.0 mL, 37.0 mmol, 1.0M in hexanes) dropwise at -78 °C. After 30 minutes, an additional portion of DIBAL (37.0 mL) was added dropwise. After an additional 30 minutes, the acetone/dry ice bath was removed, and the reaction mixture was warmed to 0°C. After an additional 30 minutes, the reaction was carefully quenched with saturated NH_4Cl . The mixture was diluted with diethyl ether (200 mL) and saturated sodium potassium tartrate (250 mL) and stirred vigorously for 2.0 hours. The two phases were separated, and the organic phase was washed with brine, dried (Na_2SO_4), filtered, and the filtrate concentrated under reduced pressure to provide the title compound with no further purification. MS (DCI) m/z 181 ($M+H$)⁺.

10

15

Example 53C

N-[4-(allyloxy)-2-fluorobenzyl]-N-(2-methyl-3-nitrophenyl)amine

The product from Example 53B and 2-methyl-3-nitroaniline were processed as described in Example 6A to provide the title compound. MS (ESI-) m/z 315 ($M-H$)⁻.

20

Example 53D

N-[4-(allyloxy)-2-fluorobenzyl]-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 53C and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound. MS (ESI+) m/z 443 ($M+H$)⁺.

25

Example 53E

4-([(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl)-3-fluorophenol

The product from Example 53D was processed as described in Example 33A to provide the title compound. MS (ESI+) m/z 403 ($M+H$)⁺.

30

Example 53F

N-(2,4-difluorobenzyl)-*N*-(2-fluoro-4-phenoxybenzyl)-*N*-(2-methyl-3-nitrophenyl)amine

The product from Example 53E and phenylboronic acid were processed as described in Example 48A to provide the title compound.

5

Example 53G*N*-(3-[(2,4-difluorobenzyl)(2-fluoro-4-phenoxybenzyl)amino]-2-methylphenyl)methanesulfonamide

The product from Example 53F was processed as described in Examples 6C and D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.36 (m, 2 H), 7.16 (m, 5 H), 6.97 (m, 3 H), 6.56-6.80 (m, 4 H), 6.10 (s, 1 H), 4.11 (s, 2 H), 4.08 (s, 2 H), 2.94 (s, 3 H), 2.27 (s, 3 H); MS (APCI) *m/z* 527 (M+H⁺).

10

Example 54

N-(3-[benzyl(4-bromo-2-fluorobenzyl)amino]-2-methylphenyl)methanesulfonamide
The product from Example 6A and 2-fluoro-4-bromo-1-(bromomethyl)benzene were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 8.94 (s, 1 H), 7.43 (dd, 1 H), 7.30 (dd, 1 H), 7.27 (m, 4 H), 7.20 (m, 2 H), 7.04 (t, 1 H), 6.98 (d, 1 H), 6.94 (d, 1 H), 4.08 (d, 4 H), 2.89 (s, 3 H), 2.34 (s, 3 H); MS (APCI+) *m/z* 477 (M+H⁺).

20

Example 55*N*-(3-[benzyl(2-chloro-4-fluorobenzyl)amino]-2-methylphenyl)methanesulfonamide

The product from Example 6A and 2-chloro-4-fluoro-1-(bromomethyl)benzene were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 8.93 (s, 1 H), 7.41 (dd, 1 H), 7.34 (dd, 1 H), 7.26 (m, 4 H), 7.21 (m, 1 H), 7.12 (t, 1 H), 7.05 (t, 1 H), 6.98 (d, 2 H), 4.13 (d, 4 H), 2.89 (s, 3 H), 2.31 (s, 3 H); MS (APCI+) *m/z* 433 (M+H⁺).

25

30

Example 56*N*-(3-[(2,4-difluorobenzyl)[4-(3-methoxyphenoxy)benzyl]amino]-2-methylphenyl)methanesulfonamide

Example 56A

N-(2,4-difluorobenzyl)-*N*-[4-(3-methoxyphenoxy)benzyl]-*N*-(2-methyl-3-nitrophenyl)amine

The product from Example 33A and 3-methoxyphenylboronic acid were processed as described in Example 48A to provide the title compound.

5

Example 56B

N-(3-{(2,4-difluorobenzyl)[4-(3-methoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 56A was processed as described in Examples 6C and D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.00-7.25 (m, 6 H), 6.92 (m, 3 H), 6.74 (m, 2 H), 6.64 (m, 1 H), 6.54 (m, 2 H), 6.13 (s, 1 H), 4.08 (s, 2 H), 4.04 (s, 2 H), 3.77 (s, 3 H), 2.95 (s, 3 H), 2.30 (s, 3 H); MS (APCI) *m/z* 539 (M+H⁺).

10

Example 57

N-(3-[benzyl(4-{[(2*Z*)-3-bromoprop-2-enyl]oxy}benzyl)amino]-2-methylphenyl)methanesulfonamide

15

Example 57A

4-{[(2*Z*)-3-bromoprop-2-enyl]oxy}benzaldehyde

20

4-Hydroxybenzaldehyde and 1,3-dibromo-1-propene were processed as described in Example 33B to provide the title compound.

Example 57B

N-(4-{[(2*Z*)-3-bromoprop-2-enyl]oxy}benzyl)-*N*-(2-methyl-3-nitrophenyl)amine

25

The product from Example 57A and 2-methyl-3-nitroaniline were processed as described in Example 6A to provide the title compound.

Example 57C

N-(3-[benzyl(4-{[(2*Z*)-3-bromoprop-2-enyl]oxy}benzyl)amino]-2-methylphenyl)methanesulfonamide

30

The product from Example 57B and benzyl bromide were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 6 H), 7.15 (m, 3 H), 6.90 (d, 1 H), 6.80 (d, 2 H), 6.30-6.50 (m, 2 H), 6.11 (s, 1 H), 4.68 (d, 2 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 2.93 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) *m/z* 515, 517 (M+H)⁺.

35

Example 58N-{3-[(2-fluorobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

5 The product from Example 26A and 2-fluorobenzyl bromide were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCL₃) δ 7.32 (m, 2 H), 7.20 (m, 5 H), 7.10 (m, 3 H), 6.95 (m, 6 H), 6.12 (s, 1 H), 4.12 (s, 2 H), 4.06 (s, 2 H), 2.93 (s, 3 H), 2.31 (s, 3 H); MS (APCI) m/z 539 (M+H⁺).

10

Example 59N-{3-[(4-methoxybenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

15 The product from Example 26A and 4-methoxybenzyl chloride were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCL₃) δ 7.32 (m, 2 H), 7.15 (m, 7 H), 6.96 (m, 3 H), 6.87 (m, 2 H), 6.77 (m, 2 H), 6.10 (s, 1 H), 4.15 (s, 2 H), 4.12 (s, 2 H), 3.77 (s, 3 H), 2.91 (s, 3 H), 2.25 (s, 3 H); MS (ESI) m/z 503 (M+H⁺).

Example 60

20 6-O-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-2,4-dideoxy-D-erythro-hexonic acid

Example 60A4-(allyloxy)phenol

25 Allyl iodide (8.2 mL, 89 mmoles) was added to a solution of hydroquinone (7.84 g, 71.2 mmoles) and K₂CO₃ (19.7 g, 142.4 mmoles) in acetone (102 mmoles). The resulting mixture was heated to 40 °C overnight and concentrated *under reduced pressure*. The crude products were filtered through a pad of silica with 1:1 hexanes:ethyl acetate. The residue was purified by flash chromatography on a prepacked Biotage column (hexane to 1:1
30 hexane:ethyl acetate) on silica gel to provide the title compound (4.40 g, 41%).

Example 60B4-[4-(allyloxy)phenoxy]benzaldehyde

35 4-Fluorobenzaldehyde (3.8 mL, 35.1 mmoles) was added to a solution of the product from Example 60A (4.39 g, 29.3 mmoles) and K₂CO₃ (8.90 g, 64.4 mmoles) in DMF (29.3 mL). The resulting mixture was heated to 100 °C for 2.5 days and then cooled to room temperature. The crude products were diluted with diethyl ether, extracted with sat. NH₄Cl,

extracted with H₂O (2X), washed with brine, dried (Na₂SO₄), and concentrated *under reduced pressure*. The residue was purified by flash chromatography on a prepacked Biotage column (hexane→9:1 hexane:ethyl acetate) on silica gel to provide the title compound (3.50 g, 81%) as a clear colorless oil.

5

Example 60C

N-{4-[4-(allyloxy)phenoxy]benzyl}-*N*-(2-methyl-3-nitrophenyl)amine

The product from Example 60B was processed as described in Example 6A to provide the title compound.

10

Example 60D

N-{4-[4-(allyloxy)phenoxy]benzyl}-*N*-benzyl-*N*-(2-methyl-3-nitrophenyl)amine

The product from Example 60C and benzyl bromide was processed as described in Example 6B to provide the title compound.

15

Example 60E

4-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenol

The product from Example 60D was processed as described in Example 33A to provide the title compound.

20

Example 60F

tert-butyl 6-*O*-[4-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]-2,4-dideoxy-3,5-*O*-(1-methylethylidene)-*D*-erythro-hexonate

A solution of the product from Example 60E (.2411 g, .55 mmoles), (3*R*, 5*S*)-6-[methylsulfonyl]oxy]-3,5-*O*-isopropylidene-3,5-dihydroxyhexanoic acid *tert*-butyl ester (prepared according to Jendralla, H.; Granzer, B.; Kerekjarto, B. v.; Krause, R.; Schacht, U.; Baader, E.; Bartmann, W.; Beck, G.; Bergmann, A.; Kessler, D.; Wess, G.; Chen, L.-J.; Granata, S.; Herchen, J.; Kleine, H.; Schüssler, H.; Wagner, K. J. Med. Chem. 1991, 34, 2962) (.1852 g, .55 mmoles), K₂CO₃ (.1515 g, 1.1 mmoles), and 18-Crown-6 (.0072 g, .027 mmoles) in DMSO (1.62 mL) under a nitrogen atmosphere was heated to 80 °C for 18 h. The crude products were diluted with diethyl ether, extracted with sat. NH₄Cl, extracted with H₂O (2X), washed with brine, dried (Na₂SO₄), and concentrated *under reduced pressure*. The residue was purified by flash chromatography (4:1 hexane:ethyl acetate→3:1 hexane:ethyl acetate) on silica gel to provide the title compound (.2460 g, 62%) as a yellow oil.

25

30

35

Example 60G

tert-butyl 6-O-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-2,4-dideoxy-3,5-O-(1-
methylethylidene)-D-erythro-hexonate

5 The product from Example 60F was processed as described in Examples 6C and D to provide the title compound.

Example 60H

6-O-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-2,4-dideoxy-D-erythro-
hexonic acid

10 A solution of the product from Example 60G (.3000 g, .41 mmols) in EtOH (2.0 mL) and THF (1.0 mL) was treated with 3N HCl (.15 mL). The resulting mixture was stirred at rt overnight, neutralized with pH 7 buffer and extracted with ethyl acetate (2x). The combined organic layers were rinsed with H₂O and brine, dried (Na₂SO₄), and
15 concentrated *under reduced pressure*. Half of the crude products were dissolved in EtOH (2.3 mL) and treated with 3 M NaOH (.14 mL). After 2.5 h, the reaction was concentrated *under reduced pressure*, diluted with CH₃CN and H₂O, and purified by preparative HPLC (CH₃CN:0.1% TRIFLUOROACETIC ACID in H₂O) on a YMC ODS Guardpak column. The procedure yielded the title compound (.0937g, 72%) as a white solid. ¹H NMR (300
20 MHz, DMSO-d₆) δ 7.22 (m, 11 H), 6.98 (m, 4 H), 6.82 (d, 2 H), 4.04 (s, 2 H), 3.99 (s, 2 H), 3.94 (m, 2 H), 3.82 (m, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.23 (m, 1 H), 2.08 (m, 1 H), 1.89 (s, 3 H), 1.56 (m, 2 H); MS (ESI) m/z 635 (M+H⁺).

Example 61

N-[3-(benzyl{4-[3-(2-methoxyethoxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide

Example 61A

30 3-(allyloxy)phenyl acetate
Resorcinol monoacetate (10.0 g, 65.7 mmols) in anhydrous DMF (100 mL) was treated with K₂CO₃ (18.2g, 131 mmols) and allyl bromide (6.83 mL, 78.9 mmols). The mixture was heated at 80°C overnight, cooled to room temperature, and diluted with ethyl acetate. The mixture was washed with H₂O, brine, dried (Na₂SO₄), filtered, and the filtrate
35 concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 8% to 10% ethyl acetate:hexanes) to provide the title compound as a colorless oil (7.19 g, 57%). MS (DCI) m/z 193 (M+H)⁺.

Example 61B3-(allyloxy)phenol

5 The product from Example 61A (7.19 g, 37.4 mmol) in THF (30 mL) was treated with 4N NaOH (19 mL, 75 mmol) and MeOH (5 mL). After one hour, the reaction mixture was concentrated under reduced pressure. The residue was dissolved in ethyl acetate, washed with aqueous NH₄Cl, brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure to provide the title compound without further purification.

10

Example 61C4-[3-(allyloxy)phenoxy]benzaldehyde

15 The crude product from Example 61B in anhydrous DMF (40 mL) was treated with K₂CO₃ (15.5 g, 112 mmol) and 4-fluorobenzaldehyde (4.82 mL, 44.9 mmol). The reaction mixture was stirred overnight at 100°C. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate. The mixture was washed with H₂O (2X), brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 8% ethyl acetate:hexanes) to provide the title compound (5.75 g, 60%).

20

Example 61DN-{4-[3-(allyloxy)phenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

25 The product from Example 61C and 2-methyl-3-nitroaniline were processed as described in Example 6A to provide the title compound (7.20 g, 82%).

Example 61EN-{4-[3-(allyloxy)phenoxy]benzyl}-N-benzyl-N-(2-methyl-3-nitrophenyl)amine

30 The product from Example 61D and benzyl bromide were processed as described in Example 6B to provide the title compound

Example 61F3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenol

35 The product from Example 61E was processed as described in Example 33A to provide the title compound. MS (ESI-) m/z 439 (M-H)⁺.

Example 61G

N-benzyl-N-{4-[3-(2-methoxyethoxy)phenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

5 The product from Example 61F (0.100 g, 0.227 mmol), polystyrene supported triphenylphosphine (0.151 g, 0.454 mmol, 3 mmol P/g resin), and di-*t*-butylazodicarboxylate (0.079 g, 0.341 mmol) in THF (2 mL) were treated with 2-methoxyethanol (0.022 g, 0.284 mmol). The reaction mixture was shaken overnight at room temperature. The mixture was diluted with ethyl acetate, washed with H₂O, brine,
10 dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure to provide the title compound without further purification.

Example 61H

N-[3-(benzyl{4-[3-(2-methoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

15 The crude product from Example 61G was processed as described in Examples 6C and D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 6 H), 7.15 (m, 4 H), 6.95 (dd, 1 H), 6.89 (d, 2 H), 6.66 (ddd, 1 H), 6.57 (ddd, 1 H), 6.52 (t, 1 H), 6.15 (s, 1 H), 4.14 (s, 2 H), 4.11 (s, 2 H), 4.05 (m, 2 H), 3.75 (m, 2 H), 3.44 (s, 3 H), 2.92 (s, 3 H), 2.29 (s, 3 H); MS (ESI+) *m/z* 547 (M+H)⁺.
20

Example 62

(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

Example 62A

ethyl [3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]acetate

30 The product from Example 61F (1.29 g, 2.93 mmol), triphenylphosphine (1.54 g, 5.86 mmol), and di-*t*-butylazodicarboxylate (1.01 g, 4.40 mmol) in anhydrous THF (6 mL) were treated with ethyl glycolate (0.35 mL, 3.66 mmol). The reaction was stirred overnight at room temperature. The mixture was diluted with ethyl acetate, washed with H₂O, brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure.
35 The residue was purified by chromatography on a pre-packed Biotage silica gel column (hexanes to 10% ethyl acetate:hexanes) to provide the title compound.

Example 62Bethyl (3-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetate

5 The product from Example 62A was processed as described in Examples 6C and D to provide the title compound. MS (ESI+) m/z 575 (M+H)⁺.

Example 62C(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

10 The product from Example 62B was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 7 H), 7.15 (t, 1 H), 7.05 (m, 3 H), 6.86 (d, 2 H), 6.69 (dd, 1 H), 6.65 (m, 2 H), 6.51 (t, 1 H), 4.63 (s, 2 H), 4.09 (s, 2 H), 3.99 (s, 2 H), 2.92 (s, 3 H), 2.20 (s, 3 H); MS (ESI+) m/z 547 (M+H)⁺.

Example 63

20 4-[4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-(2-methoxyethoxy)phenyl]butanoic acid

Example 63A3-(allyloxy)-4-bromophenyl 4-methylbenzenesulfonate

25 *p*-Toluenesulfonylchloride (2.14 g, 11.2 mmol) was added to a solution of 4-bromoresorcinol (2.12 g, 11.2 mmol) and K₂CO₃ (10.0 g, 72.4 mmol) in acetone (150 mL). The reaction was heated to 60 °C overnight, treated with allyl bromide (2.6 mL, 30.3 mmol), and heated to 60 °C for 24 h (according to Bos, M. E.; Wulff, W. D.; Miller, R. A.; Chamberlin, S.; Brandvoid, T. A. *J. Am. Chem. Soc.* **1991**, *113*, 9293). The reaction was cooled to room temperature, quenched with aqueous NH₄Cl, and concentrated under reduced pressure. The crude products were diluted with diethyl ether, extracted with H₂O, washed with brine, dried (Na₂SO₄), and concentrated *under reduced pressure*. The residue was purified by flash chromatography on a prepacked Biotage column (hexane to 4:1 hexane:ethyl acetate) on silica gel to provide the title compound (3.50 g, 81%) as a clear colorless oil.

Example 63B3-(allyloxy)-4-bromophenol

35 Potassium Hydroxide (11.2 g, 200 mmol) was added to a solution of the product

from Example 63A (3.5 g, 9.14 mmoles) in EtOH (180 mL) and H₂O (180 mL). The reaction was heated to 90 °C for 2 hours, cooled to room temperature, and concentrated under reduced pressure. The residue was purified by flash chromatography on a prepacked Biotage column (hexane to 4:1 hexane:ethyl acetate) on silica gel to provide the title compound (1.76 g, 84%).

Example 63C

4-[3-(allyloxy)-4-bromophenoxy]benzaldehyde

4-Fluorobenzaldehyde (1.15 mL, 10.76 mmoles) was added to a solution of the product from Example 63B (1.76 g, 7.69 mmoles) and K₂CO₃ (3.19 g, 23.1 mmoles) in DMF (7.7 mL). The reaction mixture was heated to 100 °C for 14 h and was cooled to room temperature. The crude products were diluted with diethyl ether, extracted with sat. NH₄Cl, extracted with H₂O (2x), rinsed with brine, dried (Na₂SO₄), and concentrated under reduced pressure. The residue was purified by flash chromatography on a prepacked Biotage column (hexane to 4:1 hexane:ethyl acetate) on silica gel to provide the title compound (2.13 g, 83%) as a white solid.

Example 63D

N-{4-[3-(allyloxy)-4-bromophenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

The product from example 63C was processed as described in Example 6A to provide the title compound.

Example 63E

N-{4-[3-(allyloxy)-4-bromophenoxy]benzyl}-N-benzyl-N-(2-methyl-3-nitrophenyl)amine

The product from example 63D and benzylbromide were processed as described in Example 6B to provide the title compound.

Example 63F

5-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)-2-bromophenol

The product from example 63E was processed as described in Example 33A to provide the title compound.

Example 63G

N-benzyl-N-{4-[4-bromo-3-(2-methoxyethoxy)phenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

The product from example 63F was processed as described in Example 62A

substituting 2-methoxyethanol for ethyl glycolate to provide the title compound.

Example 63H

5 *N*-[3-(benzyl{4-[4-bromo-3-(2-methoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 63G was processed as described in Examples 6C and 6D to provide the title compound.

Example 63I

10 4-[4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-(2-methoxyethoxy)phenyl]butanoic acid

The product from Example 63H was processed as described in Examples 49 and 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.08-7.34 (m, 10 H), 7.04 (d, 1 H), 6.97 (d, 1 H), 6.86 (d, 2 H), 6.47 (m, 2 H), 6.26 (s, 1 H), 4.18 (s, 2 H), 4.15 (s, 2 H),
15 4.01 (t, 2 H), 3.74 (t, 2 H), 3.44 (s, 3 H), 2.90 (s, 3 H), 2.66 (t, 2 H), 2.37 (t, 2 H), 2.28 (s, 3 H), 1.93 (m, 2 H); MS (APCI) m/z 633 (M+H⁺).

Example 64

20 ethyl 4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoate

Example 64A

4-(4-bromophenoxy)benzaldehyde

25 4-bromophenol was processed as described in Example 63C to provide the title compound.

Example 64B

N-[4-(4-bromophenoxy)benzyl]-*N*-(2-methyl-3-nitrophenyl)amine

30 The product from Example 64A was processed as described in Example 6A to provide the title compound.

Example 64C

N-benzyl-*N*-[4-(4-bromophenoxy)benzyl]-*N*-(2-methyl-3-nitrophenyl)amine

35 The product from Example 64B and benzyl bromide were processed as described in Example 6B to provide the title compound.

Example 64D

N-(3-{benzyl[4-(4-bromophenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 64C was processed as described in Examples 6C and 6D to provide the title compound.

Example 64E

5 ethyl 4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoate

The product from Example 64C was processed as described in Example 49 with 4-ethoxycarbonylbutylzincbromide to provide the title compound. ¹H NMR (300 MHz, CDCL₃) δ 7.25 (m, 8 H), 7.13 (m, 4 H), 6.90 (m, 4 H), 6.11 (s, 1 H), 4.13 (dd, 2 H), 4.09 (s, 10 2 H), 4.05 (s, 2 H), 2.94 (s, 3 H), 2.63 (t, 2 H), 2.32 (t, 2 H), 2.32 (s, 3 H), 1.86-2.08 (m, 2 H), 1.26 (t, 3 H); MS (ESI) m/z 587 (M+H⁺).

Example 65

15 N-[3-(benzyl{4-[3-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

Example 65A

20 N-[4-(allyloxy)benzyl]-N-benzyl-N-(2-methyl-3-nitrophenyl)amine

The product from Example 19A and benzyl bromide were processed as described in Example 6B to provide the title compound. MS (ESI⁺) m/z 389 (M+H)⁺.

Example 65B

25 4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenol

The product from Example 65A was processed as described in Example 33A to provide the title compound.

Example 65C

30 N-benzyl-N-[4-(3-bromophenoxy)benzyl]-N-(2-methyl-3-nitrophenyl)amine

The product from Example 65B and 3-bromophenylboronic acid were processed as described in Example 48A to provide the title compound. MS (ESI⁺) m/z 504 (M+H)⁺.

Example 65D

35 N-(3-{benzyl[4-(3-bromophenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 65C was processed as described in Example 6C and D to

provide the title compound.

Example 65E

5 ethyl 3-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoate

The product from Example 65D and 3-ethoxy-3-oxopropylzinc bromide (purchased from Aldrich) were processed as described in Example 49 to provide the title compound. MS (ESI+) m/z 573 (M+H)⁺.

10

Example 65F

N-[3-(benzyl{4-[3-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

15 The product from Example 65E was processed as described in Example 38A to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 7 H), 7.15 (m, 3 H), 6.94 (t, 2 H), 6.88 (d, 2 H), 6.82 (d, 1 H), 6.79 (d, 1 H), 6.24 (s, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.66 (t, 2 H), 2.93 (s, 3 H), 2.65 (m, 2 H), 2.31 (s, 3 H), 1.85 (m, 2 H), 1.35 (br.s, 1 H); MS (ESI+) m/z 531 (M+H)⁺.

20

Example 66

methyl 4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate

25

Example 66A

methyl 4-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)benzoate

30 The product from Example 65B (.4618 g, 1.33 mmoles), methyl-4-bromobenzoate (.5707 g, 2.65 mmoles), K₃PO₄ (.5634 g, 2.65 mmoles), Pd(OAc)₂ (.0238 g, .106 mmoles), and 2-(di-*tert*-butylphosphino)biphenyl (.0475 g, .159 mmoles) was treated with toluene (3.9 mL) under a nitrogen atmosphere. The resulting red solution was degassed, stirred for 10 min, and heated to 100 °C overnight. The black solution was diluted with diethyl ether, extracted with sat. NH₄Cl and sat. NaCl, dried (Na₂SO₄), and concentrated under reduced pressure. The residue was purified by flash chromatography (10:1 to 4:1 hexane:ethyl acetate) on silica gel to provide the title compound (.5904 g, 92%) as a yellow oil.

35

Example 66B

methyl 4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate

The product from Example 66B was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.83-8.11 (m, 2 H), 7.03-7.42 (m, 9 H), 6.92 (m, 5 H), 6.15 (s, 1 H), 4.11 (s, 2 H), 4.10 (s, 2 H), 3.89 (s, 3 H), 2.95 (s, 3 H), 2.34 (s, 3 H); MS (ESI) m/z 531 (M+H)⁺.

Example 67

methyl 3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate

Example 67A

methyl 4-(4-{[(2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino]methyl}phenoxy)benzoate

The product from Example 33A and methyl-3-bromobenzoate were processed as described in Example 52A to provide the title compound. MS (ESI+) m/z 519 (M+H)⁺.

Example 67B

methyl 3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate

The product from Example 67A was processed as described in Example 6C and D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.74 (dt, 1 H), 7.45 (m, 1 H), 7.40 (t, 1 H), 7.25 (m, 2 H), 7.15 (m, 4 H), 7.05 (m, 1 H), 6.90 (d, 2 H), 6.75 (m, 2 H), 6.47 (s, 1 H), 4.18 (s, 2 H), 4.12 (s, 2 H), 3.91 (s, 3 H), 2.89 (s, 3 H), 2.21 (s, 3 H); MS (ESI+) m/z 567 (M+H)⁺.

Example 68

ethyl N-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoyl)-beta-alaninate

Example 68A

methyl 4-(4-(((2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino)methyl)phenoxy)benzoate

The product from Example 33A and methyl-4-bromobenzoate were processed as described in Example 52A to provide the title compound. MS (ESI+) m/z 519 (M+H)⁺.

Example 68B

methyl 4-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)benzoate

The product from Example 68A was processed as described in Example 6C and D to provide the title compound. MS (ESI+) m/z 567 (M+H)⁺.

Example 68C

4-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoic acid

The product from Example 68B was processed as described in Example 50 to provide the title compound.

Example 68D

ethyl N-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoyl)-beta-alaninate

The product from Example 68C (0.050 g, 0.096 mmol), 1-hydroxybenzotriazole hydrate (0.015 g, 0.109 mmol), and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.021 g, 0.109 mmol) in DMF (1.5 mL) were treated with β -alanine ethyl ester hydrochloride (0.121 g, 0.136 mmol) and triethylamine (0.025 mL, 0.181 mmol).

The reaction mixture was shaken at room temperature overnight. The mixture was diluted with ethyl acetate, washed with H₂O, saturated aqueous NaHCO₃ (2X), 1N H₃PO₄ (2X), and brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by preparative HPLC (CH₃CN:0.1% trifluoroacetic acid in H₂O) on a YMC ODS Guardpak column to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.73 (d, 2 H), 7.25 (m, 2 H), 7.15 (m, 2 H), 6.95 (m, 6 H), 6.75 (m, 2 H), 6.15 (s, 1 H), 4.18 (q, 2 H), 4.13 (s, 2 H), 4.11 (s, 2 H), 3.73 (dd, 2 H), 2.96 (s, 3 H), 2.65 (t, 2 H), 2.30 (s, 3 H), 1.28 (t, 3 H); MS (ESI+) m/z 652 (M+H)⁺.

Example 69

N-{3-[benzyl(4-{3-[2-(2-methoxyethoxy)ethoxy]phenoxy}benzyl)amino]-2-
methylphenyl}methanesulfonamide

Example 69A

N-benzyl-N-(4-{3-[2-(2-methoxyethoxy)ethoxy]phenoxy}benzyl)-N-(2-methyl-3-nitrophenyl)amine

5 The product from Example 61F and di(ethylene glycol) methyl ether were processed as described in Example 61G to provide the title compound.

Example 69B

10 N-{3-[benzyl(4-{3-[2-(2-methoxyethoxy)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example was processed as described in Examples 6C and D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 6 H), 7.15 (m, 4 H), 6.93 (dd, 1 H), 6.89 (d, 2 H), 6.64 (ddd, 1 H), 6.57 (ddd, 1 H), 6.51 (t, 1 H), 6.19 (s, 1 H), 4.10 (m, 4 H), 4.05 (m, 2 H), 3.85 (m, 2 H), 3.70 (m, 2 H), 3.55 (m, 2 H), 3.37 (s, 3 H), 2.93 (s, 3 H), 2.32 (s, 3 H); MS (ESI+) m/z 591 (M+H)⁺.

Example 70

20 N-[3-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl]acetamide

Example 70A

25 2-{3-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]propyl}-1H-isoindole-1,3(2H)-dione

The product from Example 61F and N-(3-hydroxypropyl)phthalimide were processed as described in Example 61G to provide the title compound.

Example 70B

30 N-{3-[benzyl(4-{3-[3-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)propoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 70A was processed as described in Examples 6C and D to provide the title compound. MS (ESI+) m/z 676 (M+H)⁺.

Example 70C

35 N-{3-[4-[3-(3-aminopropoxy)phenoxy]benzyl](benzyl)amino]-2-

methylphenyl}methanesulfonamide

The product from Example 70B (0.323 g, 0.479 mmoles) in EtOH (5 mL) was treated with hydrazine hydrate (0.15 mL, 4.79 mmoles). The reaction was heated overnight at 60°C. Reaction mixture concentrated under reduced pressure, and residue dissolved in ethyl acetate. The mixture was washed with H₂O (2X), brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 10% to 20% MeOH:CHCl₃) to provide the title compound (0.160 g, 61%). MS (ESI+) m/z 546 (M+H)⁺.

Example 70DN-[3-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl]acetamide

The product from Example 70C (0.028 g, 0.0514 mmoles) in CH₂Cl₂ (1 mL) was treated with acetic anhydride (0.0060 mL, 0.0617 mmoles) and stirred overnight at room temperature. The reaction was concentrated under reduced pressure, and the residue was purified by preparative HPLC (CH₃CN:0.1% trifluoroacetic acid in H₂O) on a YMC ODS Guardpak column. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 10 H), 6.96 (d, 1 H), 6.89 (d, 2 H), 6.60 (m, 2 H), 6.44 (t, 1 H), 6.29 (s, 1 H), 5.85 (br.s, 1 H), 4.09 (s, 2 H), 4.06 (s, 2 H), 3.98 (t, 2 H), 3.44 (dd, 2 H), 2.91 (s, 3 H), 2.31 (s, 3 H), 1.98 (s, 3 H), 1.90-2.10 (m, 2 H); MS (ESI+) m/z 588 (M+H)⁺.

Example 71N-[3-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl]methanesulfonamide

The product from Example 70C was processed as described in Examples 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.25 (m, 8 H), 7.15 (m, 3 H), 6.85 (d, 2 H), 6.65 (m, 2 H), 6.42 (s, 1 H), 6.39 (t, 1 H), 4.65 (br.s, 1 H), 4.48 (s, 2 H), 4.45 (s, 2 H), 4.01 (t, 2 H), 3.35 (m, 2 H), 2.95 (s, 3 H), 2.78 (s, 3 H), 2.09 (s, 3 H), 2.05 (m, 2 H); MS (ESI-) m/z 622 (M-H)⁻.

Example 72N-{3-[benzyl(4-{3-[3-(dimethylamino)propoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

Example 72A

N-benzyl-N-{4-[3-(3-{[*tert*-butyl(dimethyl)silyl]oxy}propoxy)phenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

5 The product from Example 61F and 1-*t*-butyldimethylsilyloxypropan-3-ol were processed as in Example 61G to provide the title compound.

Example 72B

10 N-[3-(benzyl{4-[3-(3-{[*tert*-butyl(dimethyl)silyl]oxy}propoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 72A was processed as in Example 6C and D to provide the title compound. MS (ESI+) *m/z* 661 (M+H)⁺.

Example 72C

15 N-[3-(benzyl{4-[3-(3-hydroxypropoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

20 The product from Example 72B (0.368 g, 0.558 mmoles) in anhydrous THF (2 mL) was treated with 1.0 M tetrabutylammonium fluoride (0.67 mL, 0.669 mmoles) and mixed overnight at room temperature. Reaction diluted with ethyl acetate, washed with saturated NH₄Cl, H₂O, and brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure to provide the title compound without further purification.

Example 72D

25 N-[3-(benzyl{4-[3-(3-bromopropoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

30 The crude product from Example 72C and triphenylphosphine (0.730 g, 2.78 mmoles) in anhydrous DMF (5 mL) at 0°C were treated with *N*-bromosuccinimide (0.396 g, 2.24 mmoles). Reaction mixed at 0°C for 30 minutes. Reaction diluted with ethyl acetate, washed with H₂O (2X), brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 30% ethyl acetate:hexanes) to provide the title compound (0.214 g, 63%). MS (ESI+) *m/z* 609 (M+H)⁺.

35

Example 72E

N-{3-[benzyl(4-{3-[3-(dimethylamino)propoxy]phenoxy}benzyl)amino]-2-

methylphenyl}methanesulfonamide

The product from Example 72D in anhydrous THF (1 mL) was treated with 2.0 M dimethylamine (1.0 mL, 1.97 mmoles) in THF and heated to 70°C overnight. Reaction mixture concentrated under reduced pressure and the residue was purified by preparative HPLC (CH₃CN:0.1% trifluoroacetic acid in H₂O) on a YMC ODS Guardpak column to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 11 H), 6.86 (d, 2 H), 6.72 (s, 1 H), 6.50-6.70 (m, 2 H), 6.32 (t, 1 H), 4.39 (s, 2 H), 4.35 (s, 2 H), 3.98 (t, 2 H), 3.30 (m, 2 H), 2.91 (s, 3 H), 2.90 (s, 3 H), 2.83 (s, 3 H), 2.25 (m, 2 H), 2.13 (s, 3 H); MS (ESI+) m/z 574 (M+H)⁺.

Example 73N-[3-(benzyl{4-[3-(3-morpholin-4-ylpropoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 72D and morpholine were processed as described in Example 72E to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 11 H), 6.86 (d, 2 H), 6.63 (d, 1 H), 6.55 (m, 2 H), 6.33 (t, 1 H), 4.31 (s, 2 H), 4.28 (s, 2 H), 4.00 (m, 6 H), 3.65 (m, 2 H), 3.25 (m, 2 H), 2.95 (m, 2 H), 2.85 (s, 3 H), 2.25 (m, 2 H), 2.17 (s, 3 H); MS (ESI+) m/z 616 (M+H)⁺.

Example 744-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoic acidExample 74Aethyl 4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoate

The product from Example 48B and 4-ethoxy-4-oxobutylzinc bromide (purchased from Aldrich) were processed as described in Example 49 to provide the title compound. MS (ESI+) m/z 623 (M+H)⁺.

Example 74B4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoic acid

The product from Example 74A was processed as described in Example 50 to

provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.40 (m, 8 H), 6.85 (m, 6 H), 6.16 (d, 1 H), 4.26 (s, 2 H), 4.20 (s, 2 H), 2.76 (s, 3 H), 2.65 (m, 2 H), 2.56 (s, 3 H), 2.35 (m, 2 H), 1.95 (m, 2 H); MS (ESI+) m/z 595 (M+H)⁺.

5

Example 75

4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-[3-(2-oxopyrrolidin-1-yl)propyl]butanamide

10 The product from Example 74B and 1-(3-aminopropyl)-2-pyrrolidinone were processed as described in Example 68D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.00-7.30 (m, 8 H), 6.60-6.90 (m, 6 H), 6.25 (s, 1 H), 4.16 (s, 2 H), 4.11 (s, 2 H), 3.30-3.50 (m, 4 H), 3.20 (m, 2 H), 2.84 (s, 3 H), 2.65 (m, 2 H), 2.45 (m, 2 H), 2.24 (s, 3 H), 1.90-2.10 (m, 6 H), 1.65 (m, 2 H); MS (ESI+) m/z 719 (M+H)⁺.

15

Example 76

3-(4-{4-[(2-bromobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

20

Example 76A

3-[4-(4-formylphenoxy)phenyl]propanoic acid

4-Fluorobenzaldehyde and 3-(4-hydroxyphenyl)propionic were processed as described in Example 61C to provide the title compound.

25

Example 76B

3-[4-(4-{[(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoic acid

The product from Example 76A and 2-methyl-3-nitroaniline were processed as described in Example 6A to provide the title compound.

30

Example 76C

methyl 3-[4-(4-{[(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoate

The product from Example 76B (9.78 g, 24.1 mmols) in MeOH (24 mL) was treated with concentrated H₂SO₄ (1 mL) and heated at reflux overnight. Reaction cooled and quenched with saturated NaHCO₃. Reaction mixture diluted with ethyl acetate, washed with H₂O, brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by flash chromatography (silica gel, 20% ethyl acetate:hexanes) to

35

provide the title compound. MS (ESI-) m/z 419 (M-H)⁻.

Example 76D

5 methyl 3-[4-(4-{[(2-bromobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoate

The product from Example 76C and 2-bromobenzyl bromide were processed as described in Example 6B to provide the title compound.

10

Example 76E

methyl 3-(4-(4-(((2-bromobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenyl)propanoate

15 The product from Example 76D was processed as described in Example 6C and D to provide the title compound.

Example 76F

3-(4-{4-[(2-bromobenzyl)(2-methyl-3-[(methylsulfonyl)amino]phenyl)amino)methyl}phenoxy}phenyl)propanoic acid

20 The product from Example 76E was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.49 (dd, 2 H), 7.10-7.30 (m, 6 H), 7.05 (m, 3 H), 6.85 (m, 4 H), 6.18 (s, 1 H), 4.33 (s, 2 H), 4.18 (s, 2 H), 2.95 (m, 2 H), 2.88 (s, 3 H), 2.65 (m, 2 H), 2.22 (s, 3 H);); MS (ESI+) m/z 625 (M+H)⁺.

25

Example 77

(5-{4-[(benzyl(2-methyl-3-[(methylsulfonyl)amino]phenyl)amino)methyl]phenoxy}-2-bromophenoxy)acetic acid

30

Example 77A

[5-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)-2-bromophenoxy]acetic acid

35 The product from Example 63F and ethyl glycolate were processed as described in Example 62A to provide the title compound.

Example 77B

ethyl (5-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-bromophenoxy)acetate

The product from Example 77A was processed as described in Example 6C and D to provide the title compound.

Example 77C

(5-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-bromophenoxy)acetic acid

The product from Example 77D was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.49 (d, 1 H), 7.40 (d, 2 H), 7.25 (m, 7 H), 7.05 (m, 2 H), 6.80 (d, 2 H), 6.68 (dd, 1 H), 6.59 (d, 1 H), 4.64 (s, 2 H), 4.20-4.40 (br.s, 2 H), 4.00-4.20 (br.s, 2 H), 2.86 (s, 3 H), 2.03 (s, 3 H); MS (ESI+) m/z 625 (M+H)⁺.

Example 78

4-{[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]amino}-4-oxobutanoic acid

Example 78A

2-{2-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]ethyl}-1H-isoindole-1,3(2H)-dione

The product from Example 61F and *N*-(2-hydroxyethyl)-phthalimide were processed as described in Example 62A to provide the title compound.

Example 78B

N-{3-[benzyl(4-{3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 78A was processed as described in Example 6C and D to provide the title compound. MS (ESI+) m/z 662 (M+H)⁺.

Example 78C

N-{3-[{4-[3-(2-aminoethoxy)phenoxy]benzyl}(benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 78B was processed as described in Example 70C to provide the title compound. MS (ESI+) m/z 532 (M+H)⁺.

5

Example 78D

benzyl 4-((2-(3-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)ethyl)amino)-4-oxobutanoate

10

The product from Example 78C and benzyl succinic acid were processed as described in Example 68D to provide the title compound.

Example 78E

15

4-{[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]amino}-4-oxobutanoic acid

20

The product from Example 78D was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 10 H), 7.03 (d, 1 H), 6.87 (d, 2 H), 6.75 (m, 2 H), 6.65 (br.s, 1 H), 6.29 (s, 1 H), 6.21 (t, 1 H), 4.10-4.30 (m, 4 H), 3.91 (t, 2 H), 3.65 (m, 2 H), 2.86 (s, 3 H), 2.65 (m, 2 H), 2.55 (m, 2 H), 2.25 (s, 3 H); MS (ESI+) m/z 632 (M+H)⁺.

25

Example 79

5-{[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]amino}-5-oxopentanoic acid

30

Example 79A

benzyl 5-((2-(3-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)ethyl)amino)-5-oxopentanoate

35

The product from Example 78C and benzyl glutaric acid were processed as described in Example 68D to provide the title compound.

Example 79B

5-{[2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]amino}-5-
oxopentanoic acid

The product from Example 79A was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 10 H), 7.04 (d, 1 H), 6.87 (d, 2 H), 6.75 (br.s, 1 H), 6.65 (m, 2 H), 6.37 (s, 1 H), 6.05 (m, 1 H), 4.10-4.30 (m, 4 H), 3.95 (t, 2 H), 3.63 (dd, 2 H), 2.87 (s, 3 H), 2.41 (t, 2 H), 2.30 (m, 2 H), 2.28 (s, 3 H), 1.95 (m, 2 H); MS (ESI+) m/z 646 (M+H)⁺.

10

Example 80

N-[2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]acetamide

The product from Example 78C was processed as described in Example 70D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 11 H), 7.05 (m, 1 H), 6.88 (d, 2 H), 6.55 (m, 2 H), 6.40 (s, 1 H), 5.95 (br.s, 1 H), 4.00-4.30 (m, 4 H), 3.94 (t, 2 H), 3.65 (m, 2 H), 2.88 (s, 3 H), 2.29 (s, 3 H), 2.00 (s, 3 H); MS (ESI+) m/z 574 (M+H)⁺.

15

Example 81

N-[2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]methanesulfonamide
e

20

The product from Example 78C was processed as described in Example 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.40 (m, 10 H), 7.05 (m, 1 H), 6.87 (d, 2 H), 6.60 (m, 2 H), 6.39 (t, 1 H), 6.30 (s, 1 H), 4.85 (br.s, 1 H), 4.10-4.40 (m, 4 H), 4.05 (m, 2 H), 3.52 (dd, 2 H), 3.01 (s, 3 H), 2.86 (s, 3 H), 2.23 (s, 3 H); MS (ESI+) m/z 610 (M+H)⁺.

25

30

Example 82

methyl 2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethylcarbamate

The product from Example 78C (0.050 g, 0.0942 mmoles) in anhydrous THF (1 mL) was treated with methyl chloroformate (0.009 mL, 0.113 mmoles) and mixed overnight at room temperature. Reaction mixture concentrated under reduced pressure, and the residue was purified by preparative HPLC (CH₃CN:0.1% trifluoroacetic acid in H₂O) on a YMC ODS Guardpak column to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-

35

7.30 (m, 10 H), 6.95 (m, 1 H), 6.89 (d, 2 H), 6.55 (m, 2 H), 6.43 (s, 1 H), 6.26 (d, 1 H), 5.15 (br.s, 1 H), 4.00-4.20 (m, 4 H), 3.96 (t, 2 H), 3.68 (s, 3 H), 3.55 (m, 2 H), 2.90 (s, 3 H), 2.30 (s, 3 H); MS (ESI+) m/z 590 (M+H)⁺.

5

Example 83

(3-{4-[(4-chloro-2-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

10

Example 83A

3-(4-{[(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenol

The product from Example 61D was processed as described in Example 33A to provide the title compound.

15

Example 83B

ethyl [3-(4-{[(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]acetate

The product from Example 83A was processed as described in Example 62A to provide the title compound.

20

Example 83C

ethyl [3-(4-{[(4-chloro-2-fluorobenzyl)(2-methyl-3-
nitrophenyl)amino]methyl}phenoxy)phenoxy]acetate

25

The product from Example 83B and 4-chloro-2-fluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

30

Example 83D

ethyl (3-(4-(((4-chloro-2-fluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetate

The product from Example 83C was processed as described in Example 6C and D to provide the title compound.

35

Example 83E

(3-{4-[(4-chloro-2-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83D was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10 (d, 2 H), 7.10-7.30 (m, 5 H), 7.05 (m, 2 H), 6.86 (d, 2 H), 6.60-6.80 (m, 3 H), 6.48 (t, 1 H), 4.62 (s, 2 H), 4.25 (m, 2 H), 4.05 (m, 2 H), 2.89 (s, 3 H), 2.12 (s, 3 H); MS (ESI+) m/z 599 (M+H)⁺.

5

Example 84

3-(4-{4-[(2-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

10

Example 84A

methyl 3-[4-(4-{[(2-fluorobenzyl)(2-methyl-3-
nitrophenyl)amino]methyl}phenoxy)phenyl]propanoate

15

The product from Example 76C and 2-fluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

Example 84B

methyl 3-(4-(4-(((2-fluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenyl)propanoate

20

The product from Example 84A was processed as described in Example 6C and D to provide the title compound.

Example 84C

3-(4-{4-[(2-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

25

The product from Example 84B was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 8 H), 7.00 (m, 3 H), 6.85 (m, 4 H), 6.20 (s, 1 H), 4.22 (s, 2 H), 4.14 (s, 2 H), 2.94 (t, 2 H), 2.90 (s, 3 H), 2.68 (t, 2 H), 2.27 (s, 3 H); MS (ESI+) m/z 563 (M+H)⁺.

30

Example 85

3-(4-{4-[(4-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

35

Example 85A

methyl 3-[4-(4-{[(4-fluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoate

5 The product from Example 76C and 4-fluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

Example 85B

10 methyl 3-(4-(4-(((4-fluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenyl)propanoate

The product from Example 85A was processed as described in Example 6C and D to provide the title compound.

Example 85C

15 3-(4-{4-[[[(4-fluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

The product from Example 85B was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 8 H), 6.80-7.00 (m, 7 H), 6.26 (s, 1 H), 4.10-4.30 (m, 4 H), 2.94 (t, 2 H), 2.90 (s, 3 H), 2.68 (t, 2 H), 2.24 (s, 3 H); MS (ESI+) m/z 563 (M+H)⁺.

Example 86

25 3-(4-{4-[[[(4-chloro-2-fluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

Example 86A

30 methyl 3-[4-(4-{[(4-chloro-2-fluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoate

The product from Example 76C and 4-chloro-2-fluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

Example 86B

35 methyl 3-(4-(4-(((4-chloro-2-fluorobenzyl)(2-methyl-3-

((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenyl)propanoate

The product from Example 86A was processed as described in Example 6C and D to provide the title compound.

Example 86C3-(4-{4-[[((4-chloro-2-fluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

The product from Example 86B was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.30 (m, 7 H), 6.95 (m, 3 H), 6.85 (m, 4 H), 6.22 (s, 1 H), 4.21 (s, 2 H), 4.14 (s, 2 H), 2.95 (t, 2 H), 2.91 (s, 3 H), 2.68 (t, 2 H), 2.27 (s, 3 H); MS (ESI+) m/z 597 (M+H)⁺.

Example 87(3-{4-[[((2,4-difluorobenzyl){2-ethyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acidExample 87AN-{4-[3-(allyloxy)phenoxy]benzyl}-N-(3-nitro-2-vinylphenyl)amine

The product from Example 44B and the product from Example 61C were processed as in Example 6A to provide the title compound.

Example 87BN-{4-[3-(allyloxy)phenoxy]benzyl}-N-(2,4-difluorobenzyl)-N-(3-nitro-2-vinylphenyl)amine

The product from Example 87A and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound. MS (ESI+) m/z 529 (M+H)⁺.

Example 87C3-(4-{[(2,4-difluorobenzyl)(3-nitro-2-vinylphenyl)amino]methyl}phenoxy)phenol

The product from Example 87B was processed as described in Example 33A to provide the title compound.

Example 87D

[3-(4-{[(2,4-difluorobenzyl)(3-nitro-2-vinylphenyl)amino]methyl}phenoxy)phenoxy]acetic acid

The product from Example 87C was processed as described in Example 62A to provide the title compound. MS (ESI+) m/z 575 (M+H)⁺.

5

Example 87E

[3-(4-{[(3-amino-2-ethylphenyl)(2,4-difluorobenzyl)amino]methyl}phenoxy)phenoxy]acetic acid

10 The product from Example 87D (0.045 g, 0.0784 mmoles) in 1:1 ethyl acetate/THF (4 mL) was added to a degassed flask charged with 10% palladium on carbon catalyst (0.050 g). Reaction mixture was degassed again and stirred vigorously at room temperature under an atmosphere of hydrogen for one hour. The reaction mixture was filtered through a pad of celite, and filtrate concentrated under reduced pressure to provide the title compound
15 with no further purification (0.035 g, 82%). MS (ESI+) m/z 547 (M+H)⁺.

Example 87F

ethyl (3-(4-(((2,4-difluorobenzyl)(2-ethyl-3-

20 ((methylsulfonyl)amino)phenyl)amino)methyl}phenoxy)phenoxy)acetate

The product from Example 87E was processed as described in Example 6D to provide the title compound.

Example 87G

(3-{4-[(2,4-difluorobenzyl){2-ethyl-3-

25 [(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)acetic acid

The product from Example 87F was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.20 (m, 2 H), 7.20-7.40 (m, 4 H), 7.10 (d, 2 H), 6.87 (d, 2 H), 6.75 (m, 2 H), 6.65 (m, 2 H), 6.59 (s, 1 H), 4.63 (s, 2 H),
30 4.24 (s, 2 H), 4.13 (s, 2 H), 2.91 (s, 3 H), 2.62 (q, 2 H), 0.91 (t, 3 H); MS (ESI+) m/z 597 (M+H)⁺.

Example 88

(3-{4-[(benzyl{2-methyl-3-

35 [(methylsulfonyl)amino]phenyl}amino)methyl}benzoyl}phenoxy)acetic acid

Example 88A4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]benzoic acid

The product from Example 6D was processed as described in Example 50 to provide the title compound.

5

Example 88B4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]-N-methoxy-N-methylbenzamide

10 The product from Example 88A and *N, O*-dimethylhydroxylamine hydrochloride were processed as described in Example 68D to provide the title compound. MS (ESI+) *m/z* 468 (M+H)⁺.

15

Example 88C1-(allyloxy)-3-bromobenzene

20 3-Bromophenol (10.0 g, 57.8 mmol) in anhydrous DMF (60 mL) was treated with K₂CO₃ (24.0 g, 173 mmol) and allyl iodide (5.81 mL, 63.6 mmol). Reaction stirred overnight at 80°C. Reaction allowed to cool to room temperature and diluted with ethyl acetate. The mixture was washed with H₂O (2X), brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure to provide the title compound with no further purification.

25

Example 88D*N*-{3-[4-[3-(allyloxy)benzoyl]benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide

30 The product from Example 88C (0.983 g, 3.80 mmol) in anhydrous THF (10 mL) at -78°C was treated dropwise with 1.7 M *t*-butyllithium (4.46 mL, 7.59 mmol). Reaction stirred 30 minutes at -78°C and was then treated with the product from Example 88B (1.00 g, 1.90 mmol). Reaction stirred 15 minutes at -78°C and then allowed to warm to 0°C. Reaction quenched with saturated NH₄Cl and warmed to room temperature. The mixture was diluted with ethyl acetate, washed with H₂O, brine, dried (Na₂SO₄), filtered, and the
35 filtrate concentrated under reduced pressure. The residue was purified by chromatography on a pre-packed Biotage silica gel column (30% ethyl acetate:hexanes) to provide the title compound (0.369 g, 36%). MS (ESI-) *m/z* 539 (M-H)⁻.

Example 88E

N-{3-[4-[3-(allyloxy)benzoyl]benzyl](benzyl)amino]-2-methylphenyl}di-methanesulfonamide

5 The product from Example 88D was processed as described in Example 6D to provide the title compound.

Example 88F

10 N-(3-{benzyl[4-(3-hydroxybenzoyl)benzyl]amino}-2-methylphenyl)di-methanesulfonamide

The product from Example 88E was processed as described in Example 33A to provide the title compound.

Example 88G

15 ethyl (3-(4-((benzyl(3-(bis(methylsulfonyl)amino)-2-methylphenyl)amino)methyl)benzoyl)phenoxy)acetate

The product from Example 88F and ethyl glycolate were processed as described in Example 62A to provide the title compound. MS (ESI+) m/z 665 (M+H)⁺.

Example 88H

20 (3-{4-[(benzyl{2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]benzoyl}phenoxy)acetic acid

25 The product from Example 88G was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.65 (d, 2 H), 7.45 (m, 2 H), 7.10-7.30 (m, 11 H), 7.04 (d, 1 H), 6.50 (s, 1 H), 4.69 (s, 2 H), 4.21 (s, 2 H), 4.18 (s, 2 H), 2.93 (s, 3 H), 2.01 (s, 3 H); MS (ESI+) m/z 559 (M+H)⁺.

Example 89

30 (3-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-5-methylphenoxy)acetic acid

Example 89A

35 3-(allyloxy)-5-methylphenol

Orcinol and allyl iodide were processed as described in Example 88C to provide the

title compound. MS (DCI) m/z 165 (M+H)⁺.

Example 89B

5

4-[3-(allyloxy)-5-methylphenoxy]benzaldehyde

The product from Example 89A and 4-fluorobenzaldehyde were processed as in Example 61C to provide the title compound. MS (DCI) m/z 269 (M+H)⁺.

10

Example 89C

N-{4-[3-(allyloxy)-5-methylphenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

The product from Example 89B and 2-methyl-3-nitroaniline were processed as in Example 6A to provide the title compound. MS (ESI-) m/z 403 (M-H)⁻.

15

Example 89D

N-{4-[3-(allyloxy)-5-methylphenoxy]benzyl}-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 89C and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

20

Example 89E

3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)-5-methylphenol

25

The product from Example 89D was processed as described in Example 33A to provide the title compound.

30

Example 89F

ethyl [3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)-5-methylphenoxy]acetate

The product from Example 89E and ethyl glycolate were processed as described in Example 62A to provide the title compound.

35

Example 89G

ethyl (3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-5-methylphenoxy)acetate

The product from Example 89F was processed as described in Example 6C and D to provide the title compound.

5

Example 89H

(3-{4-[(2,4-difluorobenzyl){2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-5-methylphenoxy)acetic acid

The product from Example 89G was processed as described in Example 50 to provide the
10 title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.15 (m, 3 H), 7.10 (d, 2 H), 7.01 (dd, 1 H),
6.86 (d, 2 H), 6.75 (m, 2 H), 6.61 (s, 1 H), 6.48 (d, 2 H), 6.28 (t, 1 H), 4.59 (s, 2 H), 4.14 (s,
2 H), 4.06 (s, 2 H), 2.91 (s, 3 H), 2.28 (s, 3 H), 2.18 (s, 3 H); MS (ESI+) m/z 597 (M+H)⁺.

15

Example 90

(3-{4-[(2,4-difluorobenzyl){2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-5-methoxyphenoxy)acetic acid

Example 90A

3-(allyloxy)-5-methoxyphenol

20 5-Methoxyresorcinol and allyl iodide were processed as described in Example 88C
to provide the title compound. MS (DCI) m/z 181 (M+H)⁺.

25

Example 90B

4-[3-(allyloxy)-5-methoxyphenoxy]benzaldehyde

The product from Example 90A and 4-fluorobenzaldehyde were processed as in
Example 61C to provide the title compound. MS (DCI) m/z 285 (M+H)⁺.

30

Example 90C

N-{4-[3-(allyloxy)-5-methoxyphenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

The product from Example 90B and 2-methyl-3-nitroaniline were processed as in
Example 6A to provide the title compound. MS (ESI-) m/z 419 (M-H)⁻.

35

Example 90D

N-{4-[3-(allyloxy)-5-methoxyphenoxy]benzyl}-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 90C and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound. MS (APCI+) m/z 547 (M+H)⁺.

5

Example 90E

3-(4-[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)-5-methoxyphenol

10 The product from Example 90D was processed as described in Example 33A to provide the title compound.

Example 90F

15 ethyl [3-(4-[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)-5-methoxyphenoxy]acetate

The product from Example 90E and ethyl glycolate were processed as described in Example 62A to provide the title compound. MS (ESI+) m/z 593 (M+H)⁺.

20

Example 90G

ethyl (3-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-5-methoxyphenoxy)acetate

25 The product from Example 90F was processed as described in Example 6C and D to provide the title compound.

Example 90H

(3-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino]methyl}phenoxy)-5-methoxyphenoxy)acetic acid

30 The product from Example 90G was processed as described in Example 50 to provide the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.25 (m, 2 H), 7.16 (t, 1 H), 7.10 (d, 2 H), 7.05 (d, 1 H), 6.87 (d, 2 H), 6.75 (m, 2 H), 6.64 (s, 1 H), 6.20 (m, 2 H), 6.03 (t, 1 H), 4.58 (s, 2 H), 4.21 (s, 2 H), 4.12 (s, 2 H), 3.74 (s, 3 H), 2.89 (s, 3 H), 2.15 (s, 3 H); MS (ESI+) m/z 613 (M+H)⁺.

35

Example 91

(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

5

Example 91A

3-(allyloxy)-4-chlorophenyl 4-methylbenzenesulfonate

4-Chlororesorcinol was processed as described in Example 63A.

10

Example 91B

3-(allyloxy)-4-chlorophenol

The product from Example 91A was processed as described in Example 63B.

15

Example 91C

4-[3-(allyloxy)-4-chlorophenoxy]benzaldehyde

The product from Example 91B and 4-fluorobenzaldehyde were processed as in Example 61C to provide the title compound. MS (DCI) m/z 289 (M+H)⁺.

20

Example 91D

N-{4-[3-(allyloxy)-4-chlorophenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

The product from Example 91C and 2-methyl-3-nitroaniline were processed as in Example 6A to provide the title compound. MS (ESI-) m/z 423 (M-H)⁻.

25

Example 91E

N-{4-[3-(allyloxy)-4-chlorophenoxy]benzyl}-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

30 The product from Example 91D and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound. MS (ESI+) m/z 573 (M+H)⁺.

35

Example 91F

2-chloro-5-(4-{[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenol

The product from Example 91E was processed as described in Example 33A to provide the title compound.

Example 91G

ethyl [2-chloro-5-(4-[[2,4-difluorobenzyl](2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]acetate

The product from Example 91F and ethyl glycolate were processed as described in Example 62A to provide the title compound. MS (ESI+) m/z 597 (M+H)⁺.

Example 91H

ethyl (2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl}phenoxy)phenoxy)acetate

The product from Example 91G was processed as described in Example 6C and D to provide the title compound.

Example 91I

(2-chloro-5-{4-[[2,4-difluorobenzyl]{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)acetic acid

The product from Example 91H was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.32 (d, 2 H), 7.25 (m, 3 H), 7.20 (m, 2 H), 6.96 (d, 2 H), 6.81 (d, 2 H), 6.77 (s, 1 H), 6.75 (m, 1 H), 6.62 (d, 1 H), 4.65 (s, 2 H), 4.17 (s, 2 H), 3.98 (s, 2 H), 2.91 (s, 3 H), 1.97 (s, 3 H); MS (ESI+) m/z 617 (M+H)⁺.

Example 92

1-[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)ethyl]piperidine-2-carboxylic acid

Example 92A

N-benzyl-N-(2-methyl-3-nitrophenyl)-N-(4-{3-[2-(tetrahydro-2H-pyran-2-yloxy)ethoxy]phenoxy}benzyl)amine

The product from Example 61F and 2-(tetrahydro-2H-pyran-2-yloxy)ethanol (purchased from Fluka) were processed as described in Example 62A to provide the title compound.

Example 92B

2-[3-(4-[(benzyl(2-methyl-3-nitrophenyl)amino)methyl]phenoxy)phenoxy]ethanol

5 The product from Example 92A (0.480 g, 0.845 mmol) in EtOH (2.0 mL) was treated with pyridinium *p*-toluenesulfonate (0.0106 g, 0.0422 mmol) and heated at 55°C for four hours. The reaction was diluted with ethyl acetate, washed with H₂O, brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. The residue was purified by chromatography on a pre-packed Biotage silica gel column (hexanes to 50% ethyl acetate:hexanes) to provide the title compound.

Example 92C

2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl methanesulfonate

15 The product from Example 92B was processed as described in Example 6C and D to provide the title compound. MS (ESI+) *m/z* 611 (M+H)⁺.

Example 92D

ethyl 1-[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]piperidine-2-carboxylate

20 The product from Example 92C (0.057 g, 0.0934 mmol) in anhydrous DMF (1.0 mL) was treated with ethyl pipercolinate (0.072 mL, 0.467 mmol) and heated at 80°C overnight. The reaction was diluted with ethyl acetate, washed with H₂O (2X), brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure.

Example 92E

1-[2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]piperidine-2-carboxylic acid

30 The product from Example 92D was processed as described in Example 50 to provide the title compound. ¹H NMR (500 MHz, DMSO) δ 8.97 (s, 1 H), 7.25 (m, 7 H), 7.20 (m, 1 H), 7.05 (m, 1 H), 6.98 (d, 2 H), 6.93 (d, 2 H), 6.75 (dd, 1 H), 6.62 (t, 1 H), 6.56 (dd, 1 H), 4.35 (m, 2 H), 4.07 (s, 2 H), 4.04 (s, 2 H), 3.65 (m, 2 H), 3.55 (m, 2 H), 3.25 (m, 1 H), 2.92 (s, 3 H), 2.40 (s, 3 H), 2.15 (m, 1 H), 1.50-1.80 (m, 5 H); MS (ESI+) *m/z* 644

(M+H)⁺.

Example 93

N-[3-(benzyl(4-{3-[(1-methylpyrrolidin-3-yl)methoxy]phenoxy}benzyl)amino)-2-methylphenyl]methanesulfonamide

Example 93A

tert-butyl 3-([(methylsulfonyl)oxy]methyl}pyrrolidine-1-carboxylate

N-t-butylcarbonyl-3-hydroxymethylpyrrolidine (0.220 g, 1.09 mmoles) and triethylamine (0.168 mL, 1.20 mmoles) in anhydrous CH₂Cl₂ at -10°C were treated dropwise with methanesulfonyl chloride (0.093 mL, 1.20 mmoles). Reaction stirred one hour at 0°C and overnight at room temperature. Reaction diluted with Na₂CO₃ and extracted with CH₂Cl₂ (2X). Combined extracts were washed with H₂O, brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure to provide the title compound with no further purification. MS (DCI) m/z 280 (M+H)⁺.

Example 93B

tert-butyl 3-{[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]methyl}pyrrolidine-1-carboxylate

The product from Example 93A and Example 61F were processed as described in Example 60F. MS (ESI+) m/z 624 (M+H)⁺.

Example 93C

tert-butyl 3-[(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)methyl]pyrrolidine-1-carboxylate

The product from Example 93B was processed as described in Example 6C and D to provide the title compound.

Example 93D

N-[3-(benzyl{4-[3-(pyrrolidin-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 93C (0.108 g, 0.161 mmoles) was treated with 4 N HCl (2.0 mL/ 8.00 mmoles) in dioxane and stirred at room temperature for one hour. Reaction quenched with saturated NaHCO₃ and extracted with CH₂Cl₂. Extracts dried (Na₂SO₄),

filtered, and the filtrate concentrated under reduced pressure to provide the title compound with no further purification.

Example 93E

5 N-{3-[benzyl(4-{3-[(1-methylpyrrolidin-3-yl)methoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 93D (0.030 g, 0.0525 mmoles) in 1:1 CH₃CN/MeOH (2 mL) was treated with 37% aqueous formaldehyde (0.020 mL, 0.263 mmoles) and sodium cyanoborohydride (0.008 g, 0.131 mmoles). Reaction mixed overnight at room temperature.

10 Reaction quenched with two drops acetic acid and concentrated under reduced pressure. The residue was purified by preparative HPLC (CH₃CN:0.1% TRIFLUOROACETIC ACID in H₂O) on a YMC ODS Guardpak column to provide the title compound. ¹H NMR (500 MHz, DMSO) δ 8.96 (s, 1 H), 7.25 (m, 7 H), 7.20 (m, 1 H), 7.05 (m, 1 H), 6.97 (d, 2 H), 6.92 (d, 2 H), 6.70 (m, 1 H), 6.56 (dd, 1 H), 6.53 (dd, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.95
15 (m, 1 H), 3.50-3.80 (m, 1 H), 3.30-3.50 (m, 1 H), 3.00-3.20 (m, 2 H), 2.92 (s, 3 H), 2.80 (m, 5 H), 2.40 (s, 3 H), 2.00-2.30 (m, 1 H), 1.80 (m, 1 H); MS (ESI+) m/z 586 (M+H)⁺.

Example 94

20 N-{3-[(4-{3-[(1-acetylpyrrolidin-3-yl)methoxy]phenoxy}benzyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 93D was processed as described in Example 70D to provide the title compound. ¹H NMR (400 MHz, CDCl₃) δ (rotamer) 7.11 (d, 1 H), 7.10-7.30 (m, 8 H), 6.95 (m, 1 H), 6.90 (d, 2 H), 6.65 (m, 1 H), 6.60 (m, 2 H), 6.52 (6.45) (t, 1
25 H), 6.35 (6.22) (s, 1 H), 4.08 (s, 2 H), 4.05 (s, 2 H), 3.80-4.00 (m, 2 H), 3.50-3.80 (m, 2 H), 3.30-3.50 (m, 2 H), 2.94 (2.92) (s, 3 H), 2.70 (m, 1 H), 2.33 (2.32) (s, 3 H), 2.15 (m, 1 H), 2.08 (s, 3 H), 1.80 (m, 1 H); MS (ESI+) m/z 614 (M+H)⁺.

Example 95

30 (4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

Example 95A

35 ethyl [4-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]acetate

The product from Example 60E and ethyl glycolate were processed as described in

Example 62A to provide the title compound. MS (ESI+) m/z 527 (M+H)⁺.

Example 95B

5

ethyl 4-(4-((benzyl(2-methyl-3-

((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetate

The product from Example 95A was processed as described in Example 6C and D to provide the title compound. MS (ESI+) m/z 575 (M+H)⁺.

10

Example 95C

(4-{4-[(benzyl{2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl)phenoxy}phenoxy)acetic acid

The product from Example 95B was processed as described in Example 50 to provide the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.25 (m, 6 H), 7.15 (m, 3 H),
15 6.95 (m, 5 H), 6.83 (d, 2 H), 6.29 (s, 1 H), 4.66 (s, 2 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 2.94 (s,
3 H), 2.33 (s, 3 H); MS (ESI+) m/z 547 (M+H)⁺.

Example 96

20

4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoic
acid

The product from Example 52 was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.91-8.22 (m, 2 H), 7.06-7.37 (m, 9 H),
6.94 (m, 5 H), 6.21 (s, 1 H), 4.08 (s, 2 H), 4.07 (s, 2 H), 3.85 (br s, 1H), 2.97 (s, 3 H), 2.37
25 (s, 3 H); MS (ESI) m/z 517 (M+H)⁺.

Example 97

30

N-[3-(benzyl{4-[4-(methoxymethoxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide

Example 97A

35

N-benzyl-N-{4-[4-(methoxymethoxy)phenoxy]benzyl}-N-(2-methyl-3-nitrophenyl)amine

The product from Example 65B and methoxymethyl-4-bromophenyl ether was processed as described in Example 66A to provide the title compound.

Example 97B

N-[3-(benzyl{4-[4-(methoxymethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

5 The product from Example 97A was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, CDCL₃) δ 7.07-7.34 (m, 10 H), 6.92 (m, 3 H), 6.77 (ddd, 1 H), 6.66 (t, 1 H), 6.61 (ddd, 1 H), 6.15 (s, 1 H), 5.14 (s, 2 H), 4.06 (s, 2 H), 4.01 (s, 2 H), 3.47 (s, 3 H), 2.94 (s, 3 H), 2.32 (s, 3 H); MS (ESI) m/z 533 (M+H⁺).

10

Example 98

benzyl 3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoate

15

Example 98A

3-[4-(4-{[(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoic acid

A solution of 4-hydroxyphenylpropionic acid (4.02 g, 24.2 mmol), 4-fluorobenzaldehyde (3.63 mL, 33.9 mmol), and K₂CO₃ (6.69 g, 48.4 mmol) in DMF (24.2 mL) was heated to 100 °C for 48 h. The crude products were diluted with H₂O, washed with diethyl ether, acidified with 3N HCl, and extracted with ethyl acetate. The organic layer was washed with H₂O (2x), rinsed with brine, dried (Na₂SO₄), and concentrated *under reduced pressure* to yield crude aldehyde. The title compound was prepared as described in Example 6A using the crude aldehyde.

25

Example 98B

benzyl 3-[4-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenyl]propanoate

The product from example 98A and benzyl bromide were processed as described in Example 6B to provide the title compound.

30

Example 98C

benzyl 3-(4-{4-[(benzyl{2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoate

35

The product from Example 98B was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, CDCL₃) δ 7.02-7.45 (m, 16 H), 6.88 (m, 5 H), 6.11 (s, 1 H), 5.11 (s, 2 H), 4.04 (s, 2 H), 4.01 (s, 2 H), 2.95 (s, 3 H), 2.95 (t, 2 H),

2.67 (t, 2 H), 2.34 (s, 3 H); MS (ESI) m/z 634 (M^+).

Example 99

5 N-[3-(benzyl{4-[4-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

 The product from Example 98C was processed as described in Example 38A to provide the title compound. ^1H NMR (300 MHz, CDCl_3) δ 7.19 (m, 12 H), 6.90 (m, 4 H), 6.09 (s, 1 H), 3.93-4.24 (br.s, 4 H), 3.69 (t, 2 H), 2.95 (s, 3 H), 2.69 (m, 2 H), 2.32 (s, 3 H),
10 1.88 (m, 2 H), 1.25 (m, 1 H); MS (ESI) m/z 531 ($M+\text{H}^+$).

Example 100

15 4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoic acid

 The product from Example 64E was processed as described in Example 50 to provide the title compound. ^1H NMR (300 MHz, CDCl_3) δ 7.08-7.31 (m, 11 H), 6.89 (m, 5 H), 6.19 (s, 1 H), 4.07 (s, 2 H), 4.03 (s, 2 H), 3.48 (dd, 1 H), 2.94 (s, 3 H), 2.66 (t, 2 H), 2.38 (t, 2 H), 2.33 (s, 3 H), 1.96 (m, 2 H); MS (ESI) m/z 559 ($M+\text{H}^+$).

20

Example 101

25 N-[3-(benzyl{4-[4-(4-hydroxybutyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

 The product from Example 64E was processed as described in Example 38A to provide the title compound. ^1H NMR (300 MHz, CDCl_3) δ 7.26 (m, 7 H), 7.11 (m, 5 H), 6.86 (m, 4 H), 6.11 (s, 1 H), 4.25-4.49 (m, 6 H), 3.41-3.80 (m, 3 H), 2.84 (s, 3 H), 2.63 (m, 2 H), 2.14 (s, 3 H), 1.53-1.90 (m, 2 H); MS (ESI) m/z 545 ($M+\text{H}^+$).

30

Example 102

N-{3-[(3-bromobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

 The product from Example 26A and 3-bromobenzyl bromide were processed as described in Examples 6B-D to provide the title compound. ^1H NMR (300 MHz, CDCl_3) δ
35 7.34 (m, 4 H), 7.22 (m, 1 H), 7.12 (m, 5 H), 6.99 (m, 2 H), 6.90 (m, 4 H), 6.12 (s, 1 H), 4.04 (s, 2 H), 4.02 (s, 2 H), 2.96 (s, 3 H), 2.34 (s, 3 H); MS (APCI) m/z 553 ($M+\text{H}^+$).

Example 103N-[3-[(4-bromobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl]methanesulfonamide

The product from Example 26A and 4-bromobenzyl bromide were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 6.80-7.46 (m, 16 H), 6.12 (s, 1 H), 4.01 (s, 4 H), 2.96 (s, 3 H), 2.33 (s, 3 H); MS (ESI) m/z 553 (M+H⁺).

Example 104N-[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]glycineExample 104A3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid

The product from Example 64D was processed as described in Examples 49 and 50 to provide the title compound.

Example 104BN-[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]glycine

A solution of the product from Example 104A (.065 g, .119 mmol), glycine methyl ester hydrochloride (.0300 g, .24 mmol), and diisopropylethylamine (.042 mL, .24 mmol) in DMF (.3 mL) was treated with EDAC (.032 g, .167 mmol) and HOBt (.0225 g, .167 mmol). The resulting mixture was shaken for 12 h. The crude products were diluted with diethyl ether, extracted with sat. NH₄Cl, extracted with H₂O (2x), rinsed with brine, dried (Na₂SO₄), and concentrated *under reduced pressure*. The residue was purified by flash chromatography on silica gel. The product was dissolved in EtOH (.5 mL) and THF (.5 mL) and treated with 2N NaOH (1.0 mL) for 4 h. The resulting mixture was poured into 3N HCl and extracted with ethyl acetate (2x). The combined organics were rinsed with brine, dried (Na₂SO₄), and concentrated *under reduced pressure*. Purification by preparative HPLC (CH₃CN:0.1% trifluoroacetic acid in H₂O) on a YMC ODS Guardpak column yielded the title compound (.0625g, 87%) as a white solid. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.18 (t, 1 H), 7.23 (m, 9 H), 7.00 (m, 3 H), 6.86 (m, 4 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 3.73 (d, 2 H), 2.91 (s, 3 H), 2.79 (t, 2 H), 2.40 (m, 3 H), 2.39 (s, 3 H);

MS (APCI) m/z 602 ($M+H^+$).

Example 105

5 N-[3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]-beta-alanine

The product from Example 104A and β -alanine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ^1H NMR (300 MHz, DMSO- d_6) δ 8.97 (s, 1 H), 7.90 (t, 1 H), 7.14-7.34 (m, 8 H), 7.01 (m, 4 H), 6.87 (d, 4 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 7.71 (br s, 1H), 3.22 (dd, 2 H), 2.91 (s, 3 H), 2.77 (t, 2 H), 2.34 (m, 7 H); MS (APCI) m/z 616 ($M+H^+$).

Example 106

15 4-{[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]amino}butanoic acid

The product from Example 104A and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.80 (m, 1 H), 7.14-7.35 (m, 8 H), 6.92-7.13 (m, 4 H), 6.87 (m, 4 H), 3.96-4.29 (m, 4 H), 2.96-3.16 (m, 3 H), 2.91 (s, 3 H), 2.76 (m, 2 H), 2.09-2.47 (m, 7 H), 1.49-1.69 (m, 2 H); MS (APCI) m/z 630 (M+H⁺).

25 Example 107

N-[4-(4-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]glycine

The product from Example 100 was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.13 (t, 1 H), 7.13-7.34 (m, 8 H), 7.01 (m, 4 H), 6.89 (m, 4 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 3.72 (d, 2 H), 2.91 (s, 3 H), 2.55 (t, 2 H), 2.39 (s, 3 H), 2.14 (t, 2 H), 2.07 (s, 1 H), 1.80 (m, 2 H); MS (ESI) m/z 616 (M+H⁺).

35 Example 108

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-beta-alanine

The product from Example 100 and β -alanine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ^1H NMR (300 MHz, DMSO- d_6) δ 8.96 (s, 1 H), 7.88 (br.s, 1 H), 7.12-7.35 (m, 8 H), 7.02 (m, 4 H), 6.89 (m, 4 H), 4.04 (m, 4 H), 3.22 (m, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.37 (m, 5 H), 2.05 (m, 2 H), 1.76 (m, 2 H); MS (APCI) m/z 630 ($\text{M}+\text{H}^+$).

Example 109

4-{[4-(4-{[4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl]amino)methyl]phenoxy}phenyl)butanoyl]amino}butanoic
acid

The product from Example 100 and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 104B to provide the title compound. ^1H NMR (300 MHz, DMSO- d_6) δ 8.97 (s, 1 H), 7.80 (br.s, 1 H), 7.13-7.33 (m, 8 H), 7.01 (m, 4 H), 6.88 (m, 4 H), 3.94-4.16 (m, 4 H), 3.06 (m, 4 H), 2.91 (s, 3 H), 2.39 (s, 2 H), 2.20 (m, 2 H), 2.07 (m, 2 H), 1.79 (m, 2 H), 1.61 (m, 2 H); MS (APCI) m/z 644 ($\text{M}+\text{H}^+$).

Example 110

(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl]amino)methyl]phenoxy}phenoxy)acetic acid

Example 110A

ethyl [3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino]methyl}phenoxy)phenoxy]acetate

The product from Example 83B and 2,4 difluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

Example 110B

(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl]amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 110A was processed as described in Examples 6C, 6D, and 50 to provide the title compound. ^1H NMR (300 MHz, CDCL $_3$) δ 7.02-7.38 (m, 7 H), 6.59-7.00 (m, 7 H), 6.47 (m, 1 H), 4.62 (s, 3 H), 4.21-4.42 (br.s, 2 H), 4.15 (br.s, 2 H), 2.88 (s, 3 H), 2.68-3.05 (br.s, 1 H), 2.10 (s, 2 H); MS (APCI) m/z 583 ($\text{M}+\text{H}^+$).

Example 111

2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propanoic acid

5

Example 111A

butyl 2-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]propanoate

The product from Example 61F and butyl lactate was processed as described in Example 62A to provide the title compound.

10

Example 111B

2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propanoic acid

The product from Example 111A was processed as described in Examples 6C, 6D, and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.24 (m, 8 H), 6.88-7.08 (m, 5 H), 6.61 (m, 1 H), 6.50 (ddd, 1 H), 6.44 (t, 1 H), 4.79 (q, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.92 (br.s, 1 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 1.46 (d, 3 H); MS (ESI) m/z 561 (M+H⁺).

20

Example 112

N-{3-[benzyl(4-{3-[(2-oxotetrahydrofuran-3-yl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

25

Example 112A

3-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]dihydrofuran-2(3H)-one

The product from Example 61F and α-hydroxy-γ-butyrolactone were processed as described in Example 62A to provide the title compound.

30

Example 112B

N-{3-[benzyl(4-{3-[(2-oxotetrahydrofuran-3-yl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 112A was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.13-7.34 (m, 8 H), 6.89-7.12 (m, 5 H), 6.80 (dd, 1 H), 6.66 (t, 1 H), 6.56 (ddd, 1 H), 5.32 (dd, 1 H), 4.41 (td, 1 H), 4.25 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 2.91 (s, 3 H), 2.73 (m, 1 H),

2.40 (s, 3 H), 2.23 (m, 1 H); MS (ESI) m/z 573 ($M+H^+$).

Example 113

5 2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-4-hydroxybutanoic acid

The product from Example 112B was processed as described in Example 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.24 (m, 6 H), 6.88-7.13 (m, 5 H), 6.81 (m, 1 H), 6.41-6.70 (m, 6 H), 5.32 (m, 1 H), 4.73 (m, 1 H), 4.41 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.54 (m, 1 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 1.82-2.04 (m, 1 H); MS (ESI) m/z 591 (M+H⁺).

Example 114

15 2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-[3-(2-oxopyrrolidin-
1-yl)propyl]acetamide

The product from Example 62C and 1-(3-aminopropyl)-2-pyrrolidinone was processed as described in Example 68D to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.07 (m, 1 H), 7.23 (m, 8 H), 6.85-7.06 (m, 4 H), 6.45-6.79 (m, 4 H), 4.43 (s, 2 H), 4.08 (s, 2 H), 4.06 (s, 2 H), 2.99-3.39 (m, 6 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.17 (m, 2 H), 1.87 (m, 2 H), 1.56 (m, 2 H); MS (APCI) m/z 671 (M+H⁺).

25 Example 115
N-(3-{benzyl[4-(3-hydroxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

Example 115A

30 ethyl 3-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]propanoate

The product from Example 61F and ethyl 3-bromopropionate was processed as described in Example 33B to provide the title compound.

Example 115B

35 N-(3-{benzyl[4-(3-hydroxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 115A was processed as described in Examples 6C, 6D and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 9.54 (br.s, 1 H),

8.97 (s, 1 H), 6.87-7.33 (m, 13 H), 6.51 (ddd, 1 H), 6.37 (m, 1 H), 6.32 (t, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 2.92 (s, 3 H), 2.40 (s, 3 H); MS (ESI) m/z 489 (M+H⁺).

5

Example 116

4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid

10

Example 116A

ethyl 4-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl]phenoxy}phenoxy]butanoate

A solution of the product from Example 61F (.1000g, .227 mmoles) in DMF (.57 mL) was treated with NaH (.0109 g, .27 mmoles, 60% dispersion). After 15 min, ethyl 4-bromobutyrate (.039 mL, .27 mmoles) was added and the reaction was stirred overnight.

15 The crude products were diluted with diethyl ether, washed with sat. NH₄Cl, extracted with H₂O (2x), washed with brine, dried (Na₂SO₄), and concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel (hexane to 7:1 hexane:ethyl acetate) to provide the title compound as a yellow oil.

20

Example 116B

4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid

The product from Example 116A was processed as described in Examples 6C, 6D, and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.23 (m, 8 H), 6.88-7.08 (m, 5 H), 6.68 (dd, 1 H), 6.49 (m, 2 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.93 (t, 2 H), 3.33-3.61 (br.s, 1 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.35 (t, 2 H), 1.90 (m, 2 H); MS (ESI) m/z 575 (M+H⁺).

30

Example 117

5-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid

35

Example 117A

ethyl 5-[3-(4-{[benzyl(2-methyl-3-nitrophenyl)amino]methyl]phenoxy}phenoxy]pentanoate

The product from Example 61F and ethyl 5-bromovalerate were processed as

described in Example 116A to provide the title compound.

Example 117B

5 5-(3-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid

The product from Example 117A was processed as described in Examples 6C, 6D and 50 to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.06-7.37 (m, 8 H), 6.99 (d, 1 H), 6.89 (d, 2 H), 6.61 (ddd, 1 H), 6.50 (m, 2 H), 6.45 (t, 2 H), 3.99-4.25 (m, 4 H), 3.89 (m, 4 H), 2.89 (s, 3 H), 2.44 (m, 2 H), 2.29 (s, 3 H), 1.67-1.89 (m, 4 H); MS (ESI) m/z 589 (M+H⁺).

Example 118

15 N-[3-(4-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]-N-
methylglycine

The product from Example 104A and sarcosine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.15-7.38 (m, 8 H), 7.02 (m, 4 H), 6.87 (m, 4 H), 4.12 (s, 1 H), 4.06 (s, 2 H), 4.01 (s, 2 H), 3.99 (s, 1 H), 2.99 (s, 2 H), 2.91 (s, 3 H), 2.82 (s, 1 H), 2.76 (m, 2 H), 2.62 (m, 2 H), 2.39 (s, 3 H); MS (ESI) m/z 616 (M+H⁺).

Example 119

25 N-[4-(4-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-N-methylglycine

The product from Example 100 and sarcosine ethyl ester hydrochloride was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.12-7.36 (m, 8 H), 7.02 (m, 4 H), 6.89 (m, 4 H), 4.08 (s, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.98 (s, 1 H), 2.97 (s, 2 H), 2.91 (s, 3 H), 2.81 (s, 1 H), 2.58 (m, 2 H), 2.39 (s, 3 H), 2.33 (m, 2 H), 2.22 (m, 1 H), 1.66-1.88 (m, 2 H); MS (ESI) m/z 630 (M+H⁺).

Example 120

35 4-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid

Example 120A

N-{4-[3-(allyloxy)phenoxy]benzyl}-*N*-(2,4-difluorobenzyl)-*N*-(2-methyl-3-nitrophenyl)amine

5 The product from Example 61D and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound.

Example 120B

10 3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenol

 The product from Example 120A was processed as described in Example 33A to provide the title compound.

Example 120C

15 ethyl 4-[3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino]methyl}phenoxy)phenoxy]butanoate

 The product from Example 120B and ethyl 4-bromobutyrate were processed as described in Example 116A to provide the title compound.

Example 120D

20 4-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino]methyl}phenoxy)phenoxy)butanoic acid

 The product from Example 120C was processed as described in Examples 6C, 6D
25 and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.28 (m, 4 H), 6.86-7.19 (m, 7 H), 6.69 (m, 1 H), 6.49 (m, 2 H), 4.08 (s, 2 H), 4.06 (s, 2 H), 3.94 (t, 2 H), 2.90 (s, 3 H), 2.38 (m, 2 H), 2.33 (s, 3 H), 1.91 (m, 2 H); MS (APCI) m/z 611 (M+H⁺).

Example 121

30 5-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino]methyl}phenoxy)phenoxy)pentanoic acid

Example 121A

35 ethyl 5-[3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-

nitrophenyl)amino)methyl}phenoxy}phenoxy}pentanoate

The product from Example 120B and ethyl 5-bromovalerate were processed as described in Example 116A to provide the title compound.

Example 121B5-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)pentanoic acid

The product from Example 121A was processed as described in Examples 6C, 6D and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.28 (m, 4 H), 6.85-7.20 (m, 7 H), 6.69 (dd, 1 H), 6.47 (m, 2 H), 4.08 (s, 2 H), 4.06 (s, 2 H), 3.92 (t, 2 H), 2.90 (s, 3 H), 2.33 (s, 3 H), 2.26 (t, 2 H), 1.48-1.81 (m, 4 H); MS (ESI) m/z 625 (M+H⁺).

Example 122N-[(3-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)acetyl]glycine

The product from Example 62C was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.37 (t, 1 H), 7.13-7.35 (m, 8 H), 6.88-7.09 (m, 5 H), 6.73 (ddd, 1 H), 6.60 (t, 1 H), 6.54 (ddd, 1 H), 4.50 (s, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.79 (d, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H); MS (APCI) m/z 604 (M+H⁺).

Example 123N-[(3-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)acetyl]-beta-alanine

The product from Example 62C and β-alanine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.10 (t, 1 H), 7.24 (m, 8 H), 6.99 (m, 5 H), 6.71 (m, 1 H), 6.55 (m, 2 H), 4.43 (s, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.31 (dd, 2 H), 2.88 (s, 3 H), 2.40 (t, 4 H), 2.07 (s, 2 H); MS (APCI) m/z 618 (M+H⁺).

Example 124N-[(3-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl}phenoxy}phenoxy)acetyl]-N-methylglycine

The product from Example 62C and sarcosine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.24 (m, 8 H), 6.89-7.09 (m, 5 H), 6.66 (m, 1 H), 6.54 (t, 1 H), 6.47 (m, 1 H), 4.85 (s, 1 H), 4.70 (s, 1 H), 4.14 (s, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.99 (s, 1 H), 3.91 (br.s, 2 H), 3.01 (s, 2 H), 2.91 (s, 2 H), 2.82 (s, 2 H), 2.40 (s, 2 H); MS (APCI) m/z 618 (M+H⁺).

Example 125

4-[(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]amino}butanoic acid

The product from Example 62C and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.10 (t, 1 H), 7.24 (m, 8 H), 6.88-7.10 (m, 5 H), 6.71 (dd, 1 H), 6.55 (m, 2 H), 4.43 (s, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.12 (dd, 2 H), 2.92 (s, 3 H), 2.40 (s, 3 H), 2.18 (t, 2 H), 1.63 (m, 2 H); MS (APCI) m/z 632 (M+H⁺).

Example 126

ethyl 4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoate

The product from Example 116A was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.07-7.31 (m, 10 H), 6.90 (m, 3 H), 6.62 (m, 1 H), 6.54 (m, 1 H), 6.50 (t, 1 H), 6.14 (s, 1 H), 4.13 (q, 2 H), 4.07 (s, 2 H), 4.04 (s, 2 H), 3.96 (t, 2 H), 2.94 (s, 3 H), 2.49 (t, 2 H), 2.34 (s, 3 H), 2.08 (m, 2 H), 1.24 (t, 3 H); MS (APCI) m/z 603 (M+H⁺).

Example 127

ethyl 5-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoate

The product from Example 117A was processed as described in Examples 6C and 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.07-7.31 (m, 10 H), 6.91 (m, 3 H), 6.61 (ddd, 1 H), 6.54 (ddd, 1 H), 6.49 (t, 1 H), 6.14 (s, 1 H), 4.12 (dd, 2 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.92 (m, 2 H), 2.94 (s, 3 H), 2.38 (m, 2 H), 2.33 (s, 3 H), 1.80 (m, 4 H), 1.25 (t, 3 H); MS (APCI) m/z 617 (M+H⁺).

Example 128

N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]glycine

5 The product from Example 116B was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.18 (t, 1 H), 7.14-7.34 (m, 8 H), 6.88-7.09 (m, 5 H), 6.68 (ddd, 1 H), 6.49 (m, 2 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.93 (t, 2 H), 3.82 (br.s, 1 H), 3.72 (d, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.27 (t, 2 H), 1.91 (m, 2 H); MS (APCI) m/z 630 (M-H⁺).

Example 129

N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-beta-alanine

15 The product from Example 116B and β-alanine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.90 (t, 1 H), 7.14-7.36 (m, 8 H), 6.85-7.12 (m, 5 H), 6.67 (ddd, 1 H), 6.48 (m, 2 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.90 (t, 2 H), 3.80 (br.s, 1 H), 3.22 (dd, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.35 (t, 2 H), 2.19 (t, 2 H), 1.76-1.98 (m, 2 H); MS (APCI) m/z 646 (M+H⁺).

Example 130

4-{4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl}amino}butanoic acid

25 The product from Example 116B and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.83 (t, 1 H), 7.13-7.34 (m, 8 H), 6.87-7.09 (m, 5 H), 6.67 (m, 1 H), 6.47 (m, 2 H), 4.27-4.62 (br.s, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.91 (t, 2 H), 3.03 (dd, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.19 (t, 4 H), 1.88 (m, 2 H), 1.59 (m, 2 H); MS (APCI) m/z 660 (M+H⁺).

Example 131

N-[5-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]glycine

The product from Example 117B was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.12 (t, 1 H), 7.23 (m, 8 H), 6.87-7.08 (m, 5 H), 6.68 (m, 1 H), 6.48 (m, 2 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.91 (t, 2 H), 3.90-4.23 (br.s, 1 H), 3.72 (d, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.17 (t, 2 H), 1.52-1.77 (m, 4 H); MS (APCI) m/z 646 (M+H⁺).

Example 132

N-[5-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-beta-alanine

The product from Example 117B and β-alanine ethyl ester hydrochloride were processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 8.97 (s, 1 H), 7.86 (t, 1 H), 7.23 (m, 8 H), 6.88-7.08 (m, 5 H), 6.67 (m, 1 H), 6.48 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.91 (t, 2 H), 3.35 (br.s, 1 H), 3.22 (dd, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.35 (t, 2 H), 2.09 (t, 2 H), 1.50-1.74 (m, 4 H); MS (APCI) m/z 658 (M-H⁻).

Example 133

4-{[5-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]amino}butanoic acid

The product from Example 117B and ethyl 4-aminobutyrate hydrochloride was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 8.97 (s, 1 H), 7.79 (t, 1 H), 7.24 (m, 8 H), 6.87-7.09 (m, 5 H), 6.68 (dd, 1 H), 6.48 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.91 (t, 2 H), 3.27-3.53 (m, 1 H), 3.03 (dd, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.20 (t, 2 H), 2.10 (t, 2 H), 1.49-1.74 (m, 6 H); MS (APCI) m/z 674 (M+H⁺).

Example 134

N-[3-(benzyl{4-[3-(2-hydroxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 92B was processed as described in Example 6C, 6D, and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.24 (m, 8 H), 6.88-7.08 (m, 5 H), 6.70 (m, 1 H), 6.49 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (t, 2 H), 3.67 (t, 2 H), 3.35 (br.s, 1 H), 2.92 (s, 3 H), 2.40 (s, 3 H); MS (APCI) m/z 533 (M+H⁺).

Example 135

5 [2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethoxy]acetic acid

Example 135A

10 ethyl {2-[3-(4-{[benzyl(2-methyl-3-
nitrophenyl)amino]methyl}phenoxy)phenoxy]ethoxy}acetate

The product from Example 92B and ethyl iodoacetate were processed as described in Example 116A to provide the title compound.

Example 135B

15 [2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethoxy]acetic acid

The product from Example 135A was processed as described in Example 6C, 6D, and 50 to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.24 (m, 8 H), 6.88-7.09 (m, 5 H), 6.70 (m, 1 H), 6.50 (m, 2 H), 4.07 (s, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (t, 1 H), 3.78 (m, 2 H), 3.67 (t, 1 H), 3.36 (br.s, 1 H), 2.92 (s, 3 H), 2.40 (s, 3 H); MS (APCI) m/z 591 (M+H⁺).

Example 136

25 2,4-dideoxy-6-O-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-D-erythro-hexonic acid

Example 136A

30 tert-butyl 2,4-dideoxy-6-O-[3-(4-{[(2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino]methyl}phenoxy)phenyl]-3,5-O-(1-methylethylidene)-D-erythro-
hexonate

The product from Example 120B was processed as described in Example 60F to provide the title compound.

35

Example 136B

tert-butyl 2,4-dideoxy-6-O-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-3,5-O-(1-
methylethylidene)-D-erythro-hexonate

5 The product from Example 136A was processed as described in Examples 6C and D to provide the title compound.

Example 136C

2,4-dideoxy-6-O-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-D-erythro-hexonic acid

10 The product from Example 136B was processed as described in Example 60H to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.00-7.21 (m, 7 H), 6.92 (m, 1 H), 6.83 (m, 2 H), 6.71 (m, 2 H), 6.53 (m, 2 H), 6.42 (s, 1 H), 4.30 (br.s, 1 H), 4.16 (br.s, 1 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.62-3.87 (br.s, 4 H), 2.88 (s, 3 H), 2.33-2.58 (br.s, 2 H), 2.25 (s, 3 H), 2.08 (s, 1 H), 1.71 (br.s, 2 H); MS (ESI) m/z 671 (M+H⁺).

15

Example 137

(3-{4-[(2-bromobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

20

Example 137A

ethyl (3-(4-(((2-bromobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetate

25 The product from Example 83B and 2-bromobenzyl bromide were processed as described in Example 6B-D to provide the title compound.

Example 137B

30 The product from Example 137A was treated with 2:1:1 2N aqueous sodium hydroxide:tetrahydrofuran:water overnight. The mixture was acidified with 2N aqueous hydrochloric acid and extracted with ethyl acetate. The ethyl acetate phase was dried over sodium sulfate, filtered and the filtrate concentrated under reduced pressure to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.93 (s, 1 H), 7.55 (d, 1 H), 7.42 (d, 1 H), 7.27 (m, 4 H), 7.15 (t, 1 H), 7.07 (m, 1 H), 7.00 (m, 2 H), 6.92 (d, 2 H), 6.66 (d, 1 H), 6.51 (m, 2 H), 4.64 (s, 2 H), 4.17 (s, 2 H), 4.11 (s, 2 H), 2.89 (s, 3 H), 2.32 (s, 3 H); MS (ESI⁺) m/z 625 (M+H)⁺.

35

Example 138N-{3-[(4-benzoylbenzyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 4-(bromomethyl)benzophenone were processed as described in Examples 6B, 6C, and 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.75 (m, 7H), 7.60 (m, 2H), 7.50 (m, 2H), 7.35 (dd, 1H), 6.25 (m, 2H), 7.10 (t, 2H), 6.9 (d, 2H), 4.15 (s, 2H), 4.10 (s, 2H), 2.97 (s, 3H), 2.39 (s, 3H); MS (ESI+) m/z 485 (M+H)⁺.

Example 139N-(3-{benzyl[4-(phenylthio)benzyl]amino}-2-methylphenyl)methanesulfonamideExample 139AN-benzyl-N-(2-methyl-3-nitrophenyl)-N-[4-(phenylthio)benzyl]amine

4-thiophenoxybenzaldehyde was processed as described in Example 6A to provide the title compound.

Example 139BN-(3-{benzyl[4-(phenylthio)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 139A and benzyl bromide were processed as described in Examples 6B, 6C, and 6D to provide the title compound. ¹H NMR (300 MHz, CDCl₃) δ 7.33 (m, 2H), 7.21 (m, 6H), 7.14 (m, 4H), 6.90 (m, 3H), 6.88 (d, 2H), 6.13 (s, 1H), 4.08 (d, 2H), 4.02 (d, 2H), 2.94 (s, 3H), 2.34 (s, 3H); MS (ESI+) m/z 487 (M+H)⁺.

Example 140N-{3-[[4-(3-acetylphenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 33A and 3-acetylphenylboronic acid were processed as described in Examples 48A, 6C, and 6D to provide the title compound. ¹H NMR (MeOH, 500 MHz) δ 7.72 (m, 1 H), 7.49 (m, 1 H), 7.46 (d, 1 H), 7.28 (m, 2 H), 7.21 (m, 2 H), 7.09 (m, 2 H), 7.02 (m, 1 H), 6.91 (m, 2 H), 6.82 (m, 2 H), 4.15 (s, 2 H), 4.12 (s, 2 H), 2.88 (s, 3 H), 2.55 (s, 3 H), 2.36 (s, 3 H); MS(ESI, +Q1MS) m/e 551.

Example 141

N-(3-((2,4-difluorobenzyl)[4-(3,4-dimethoxyphenoxy)benzyl]amino)-2-methylphenyl)methanesulfonamide

The product from Example 33A and 3,4-dimethoxyphenylboronic acid were processed as described in Examples 48A, 6C, and 6D to provide the title compound. ¹H NMR (MeOH, 500 MHz) δ 7.22 (m, 3 H), 7.08 (m, 2 H), 6.99 (m, 1 H), 6.91 (d, 1 H), 6.83 (m, 4 H), 6.64 (d, 1 H), 6.48 (dd, 1 H), 4.12 (s, 2 H), 4.08 (s, 2 H), 3.81 (s, 3 H), 3.76 (s, 3 H), 2.88 (s, 3 H), 2.35 (s, 3 H); MS(ESI, +Q1MS) m/e 569.

Example 142

N-{3-[(2,4-difluorobenzyl)(4-ethoxybenzyl)amino]-2-methylphenyl}methanesulfonamide

Example 142A

N-(2,4-difluorobenzyl)-N-(4-ethoxybenzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 33A(39 mg, 0.1 mmole) and di-t-butylazo dicarboxylate (35 mg, 0.15 mmole) in tetrahydrofuran (1.5 ml) were added to PPh₃ on resin (70 mg, 0.3 mmole: loading 3 mmole/g) followed by ethanol (5.7 mg, 0.125 mmole) in 50 microlitre tetrahydrofuran. Reaction mixture was shaken at room temperature overnight. Filtered off and solvent removed. Solaris 530 organic synthesis system from PE Biosystems was used to do this reaction. The residue purified by HPLC (CH₃CN:0.1 TFA in H₂O) on a YMC ODS guardpak column.

Example 142B

The product from Example 142A processed as described in Example 6C-D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.22 (d, 1 H), 7.14 (m, 4 H), 6.94 (d, 1 H), 6.78 (d, 2 H), 6.72 (m, 2 H), 6.08 (s, 1 H), 4.14 (s, 2 H), 4.06 (s, 2 H), 3.99 (q, 2 H), 2.91 (s, 3 H), 2.25 (s, 3 H), 1.39 (t, 3 H); MS (ESI+) m/z 461 (M+H)⁺.

Example 143

N-(3-((2,4-difluorobenzyl)[4-(pent-3-ynoxy)benzyl]amino)-2-methylphenyl)methanesulfonamide

The product from Example 33A and 3-pentyn-1-ol were processed as described in Examples 142A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.20 (d, 1 H), 7.11 (m, 4 H), 6.89 (d, 1 H), 6.80 (d, 2 H), 6.72 (m, 2 H), 6.06 (s, 1 H), 4.05 (s, 2 H), 4.01 (m, 4 H), 2.93 (s, 3 H), 2.58 (m, 2 H), 2.29 (s, 3 H), 1.79 (t, 3 H); MS (ESI+)

m/z 499 (M+H)⁺.

Example 144

5 N-(3-((4-chloro-2-fluorobenzyl)[4-(methylthio)benzyl]amino)-2-
 methylphenyl)methanesulfonamide

Example 144A

10 2-Methyl-3-nitroaniline (3.65 g, 24 mmoles) and 2-fluoro-4-chlorobenzaldehyde (6.47 g, 40.8 mmoles) were processed as described in example 6A to provide the product.

Example 144B

15 The product from Example 144A and 4-Methylthio-1-(iodomethyl)benzene prepared as in Example 32A were processed as described in Example 6B-D to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.97 (s, 1 H), 7.33 (dd, 1 H), 7.25 (t, 1 H), 7.18 (m, 5 H), 7.04 (t, 1 H), 6.98 (d, 1 H), 6.92 (d, 1 H), 4.08 (s, 2 H), 4.04 (s, 2 H), 2.90 (s, 3 H), 2.43 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 479 (M+H)⁺.

20

Example 145

N-{3-[(4-chloro-2-fluorobenzyl)(2-fluorobenzyl)amino]-2-
 methylphenyl}methanesulfonamide

25 The product from Example 144A and 2-fluoro-1-(bromomethyl)benzene were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.96 (s, 1 H), 7.33 (dd, 1 H), 7.26 (m, 3 H), 7.19 (dd, 1 H), 7.11 (m, 2 H), 7.05 (t, 1 H), 7.00 (d, 1 H), 6.96 (d, 1 H), 4.13 (d, 4 H), 2.87 (s, 3 H), 2.28 (s, 3 H); MS (ESI+) m/z 451 (M+H)⁺.

30

Example 146

N-(3-{benzyl}[(2'-cyano-1,1'-biphenyl-4-yl)methyl]amino)-2-
 methylphenyl)methanesulfonamide

35 The product from Example 6A and 4'-bromomethyl-2-cyanobiphenyl were processed as described in Examples 6B-D to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 8.97 (s, 1 H), 7.92 (d, 1 H), 7.76 (t, 1 H), 7.60 (d, 1 H), 7.56 (td, 1 H), 7.51 (d, 2 H), 7.45 (d, 2 H), 7.29 (d, 4 H), 7.21 (m, 1 H), 7.04 (m, 2 H), 6.98 (d, 1 H), 4.15 (s, 2 H), 4.10 (s, 2 H), 2.92 (s, 3 H), 2.44 (s, 3 H); MS (ESI-) m/z 480 (M-H).

Example 147

5 3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-morpholin-4-
ylethyl)propanamide

Dicyclohexylcarbodiimide on resin (70 mg) was added to a Robbin's Organic Synthesis Block (Robbins Scientific Corporation) followed by the product from Example 104A (375 microlitre of 0.13 M solution) in 20 % dimethylacetamide in 1,2-dichloroethane. Then HOAt (375 micolitre of 0.23 M solution in 20 % DME in DCE and 4-(2-aminoethyl)morpholine (375 microlitre of 0.2 M solution in 20 % DME/DCE) were added using Gilson Liquid Handler. Reaction block was covered and shaken gently at room temperature overnight. Trisamine resin (57 mg, 4.3 mmole/g loading) was added and shaken two hours. Filtered it off in a Robbin's Collection Block and washed with CH₂Cl₂ (4x0.5 ml). Filtrates concentrated. Residue purified by HPLC to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.95 (s, 1 H), 8.11 (t, 1 H), 7.26 (m, 6 H), 7.20 (m, 3 H), 7.04 (t, 1 H), 6.97 (m, 2 H), 6.89 (m, 4 H), 4.04 (d, 4 H), 3.96 (s, 4 H), 3.63 (s, 4 H), 3.14 (t, 2 H), 3.10 (s, 2 H), 2.92 (s, 3 H), 2.81 (t, 2 H), 2.41 (m, 5 H); MS (ESI+) m/z 657 (M+H)⁺.

20

Example 148

3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-[3-(2-oxopyrrolidin-1-
yl)propyl]propanamide

25 The product from Example 104A and 1-(3-aminopropyl)-2-pyrrolidinone were processed as described in Example 147 to provide the title compound. ¹H NMR (500 MHz, CDCL₃) δ 7.26 (m, 2 H), 7.19 (m, 6 H), 7.11 (m, 3 H), 6.88 (m, 5 H), 6.79 (m, 1 H), 6.16 (s, 1 H), 4.04 (d, 4 H), 3.38 (t, 2 H), 3.22 (t, 2 H), 3.16 (dd, 2 H), 2.95 (m, 5 H), 2.51 (t, 2 H), 2.42 (t, 2 H), 2.33 (s, 3 H), 2.06 (m, 2 H), 1.62 (m, 2 H); MS (ESI+) m/z 669 (M+H)⁺.

30

Example 149

3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(5-
hydroxypentyl)propanamide

35

The product from Example 104A and 5-amino-1-pentanol were processed as described in Example 147 to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ

8.95 (s, 1 H), 7.73 (t, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (t, 1 H), 6.96 (m, 2 H), 6.87 (m, 4 H), 4.03 (d, 4 H), 3.35 (m, 2 H), 3.00 (dd, 2 H), 2.91 (s, 3 H), 2.78 (t, 2 H), 2.39 (s, 3 H), 2.33 (t, 2 H), 1.37 (m, 4 H), 1.23 (m, 2 H); MS (ESI+) m/z 630 (M+H)⁺.

5

Example 150

3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(4-hydroxybutyl)propanamide

10 The product from Example 104A and 4-amino-1-butanol were processed as described in Example 147 to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.95 (s, 1 H), 7.74 (t, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (t, 1 H), 6.96 (m, 2 H), 6.86 (m, 4 H), 4.03 (d, 4 H), 3.35 (m, 2 H), 3.03 (m, 2 H), 2.91 (s, 3 H), 2.78 (t, 2 H), 2.39 (s, 3 H), 2.34 (m, 2 H), 1.37 (m, 4 H); MS (ESI+) m/z 616 (M+H)⁺.

15

Example 151

3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(3-hydroxypropyl)propanamide

20

The product from Example 104A and 3-amino-1-propanol were processed as described in Example 147 to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 8.95 (s, 1 H), 7.75 (t, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (t, 1 H), 6.96 (m, 2 H), 6.87 (m, 4 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 3.35 (m, 2 H), 3.07 (dd, 2 H), 2.91 (s, 3 H), 2.78 (t, 2 H), 2.39 (s, 3 H), 2.34 (t, 2 H), 1.50 (m, 2 H); MS (ESI+) m/z 602 (M+H)⁺.

25

Example 152

3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-hydroxyethyl)propanamide

30

The product from Example 104A and ethanolamine were processed as described in Example 147 to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.95 (s, 1 H), 7.78 (t, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (t, 1 H), 6.97 (m, 2 H), 6.87 (m, 4 H), 4.03 (d, 4 H), 3.35 (m, 2 H), 3.10 (dd, 2 H), 2.91 (s, 3 H), 2.78 (t, 2 H), 2.39 (s, 3 H), 2.36 (m, 2 H); MS (ESI+) m/z 588 (M+H)⁺.

35

Example 153

3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(6-
hydroxyhexyl)propanamide

5 The product from Example 104A and 6-amino-1-hexanol processed as described in Example 147 to provide the title compound. ¹H NMR (500 MHz, DMSO-D) δ 8.95 (s, 1 H), 7.73 (t, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (t, 1 H), 6.96 (m, 2 H), 6.87 (m, 4 H), 4.03 (d, 4 H), 3.35 (m, 2 H), 3.00 (dd, 2 H), 2.91 (s, 3 H), 2.78 (t, 2 H), 2.39 (s, 3 H), 2.33 (t, 10 2 H), 1.37 (m, 4 H), 1.23 (m, 4 H); MS (APCI+) m/z 645 (M+H)⁺.

Example 154

N-(3-{(2,4-difluorobenzyl)[4-(3-furylmethoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide

15 The product from Example 33A and 3-furanmethanol were processed as described in Examples 142A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.49 (m, 1 H), 7.43 (t, 1 H), 7.21 (d, 1 H), 7.12 (m, 4 H), 6.91 (d, 1 H), 6.86 (d, 2 H), 6.73 (m, 2 H), 6.47 (d, 1 H), 6.08 (s, 1 H), 4.91 (s, 2 H), 4.07 (s, 2 H), 4.01 (s, 2 H), 2.94 (s, 3 H), 20 2.29 (s, 3 H); MS (APCI-) m/z 511 (M-H).

Example 155

N-(3-{(2,4-difluorobenzyl)[4-(2-furylmethoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide

25 The product from Example 33A and furfuryl alcohol were processed as described in Examples 142A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.44 (m, 1 H), 7.21 (d, 1 H), 7.13 (m, 4 H), 6.92 (d, 1 H), 6.87 (d, 2 H), 6.72 (m, 2 H), 6.41 (d, 1 H), 6.38 (m, 1 H), 6.05 (s, 1 H), 4.96 (s, 2 H), 3.96-4.18 (m, 4 H), 2.90 (s, 3 H), 2.26 30 (s, 3 H); MS (APCI-) m/z 511 (M-H).

Example 156

N-{3-[[4-(1,3-benzodioxol-5-ylmethoxy)benzyl](2,4-difluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide

35 The product from Example 33A and piperonyl alcohol were processed as described in Examples 142A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃)

δ 7.21 (d, 1 H), 7.13 (m, 4 H), 6.92 (m, 2 H), 6.85 (m, 3 H), 6.80 (d, 1 H), 6.72 (m, 2 H), 6.05 (s, 1 H), 5.96 (s, 2 H), 4.92 (s, 2 H), 4.11 (s, 2 H), 4.03 (s, 2 H), 2.91 (s, 3 H), 2.27 (s, 3 H); MS (APCI-) m/z 565 (M-H).

5

Example 157

N-{3-[{4-[(6-chloro-1,3-benzodioxol-5-yl)methoxy]benzyl}(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 33A and 6-chloropiperonyl alcohol were processed as described in Examples 142A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.22 (d, 1 H), 7.14 (m, 3 H), 6.97 (s, 1 H), 6.94 (d, 1 H), 6.85 (d, 3 H), 6.72 (m, 2 H), 6.06 (s, 1 H), 5.97 (s, 2 H), 5.03 (s, 2 H), 4.11 (s, 2 H), 4.05 (s, 2 H), 2.91 (s, 3 H), 2.26 (s, 3 H); MS (APCI-) m/z 599 (M-H).

15

Example 158

N-(3-{(2,4-difluorobenzyl)[4-(2-phenylethoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 33A and 2-phenylethyl alcohol were processed as described in Examples 142A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.26 (m, 6 H), 7.11 (d, 4 H), 6.90 (d, 1 H), 6.78 (d, 2 H), 6.72 (m, 2 H), 6.04 (s, 1 H), 4.13 (t, 2 H), 4.07 (s, 2 H), 4.01 (s, 2 H), 3.08 (t, 2 H), 2.89 (s, 3 H), 2.27 (s, 3 H); MS (APCI-) m/z 535 (M-H).

25

Example 159

N-(3-{benzyl[4-(3-isopropoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

30

Example 159A

The product from Example 61F (40mg, 0.09 mmole) and di-t-butylazo dicarboxylate (33.6mg, 0.14 mmole) in tetrahydrofuran (1.5 ml) were added to PPh₃ on resin (63 mg, 0.19 mmole: loading 3 mmole/g) followed by isopropanol (6.8 mg, 0.113 mmole) in 50microlitre tetrahydrofuran. Reaction mixture was shaken at room temperature overnight. Filtered off and solvent removed. The residue purified by HPLC.

35

Example-159B

The product from Example 159A processed as described in Example 6C-D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.10-7.32 (m, 10 H), 6.97 (br.s, 1 H), 6.89 (d, 2 H), 6.62 (m, 1 H), 6.51 (m, 2 H), 6.09 (s, 1 H), 4.49 (m, 1 H), 4.00-4.24 (br.s, 4 H), 2.90 (s, 3 H), 2.30 (s, 3 H), 1.32 (s, 3 H), 1.30 (s, 3 H); MS (ESI+) m/z 531 (M+H)⁺.

5

Example 160

N-[3-(benzyl{4-[3-(cyclobutyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

10 The product from Example 61F and cyclobutanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.11-7.36 (m, 10 H), 7.00 (s, 1 H), 6.88 (d, 2 H), 6.53 (m, 2 H), 6.40 (s, 1 H), 6.07 (s, 1 H), 4.59 (m, 1 H), 4.21 (Br.S., 4 H), 2.85 (s, 3 H), 2.40 (m, 2 H), 2.27 (s, 3 H), 2.15 (m, 2 H), 1.86 (m, 1 H), 1.67 (m, 1 H); MS (ESI+) m/z 543 (M+H)⁺.

15

Example 161

N-(3-{benzyl[4-(3-sec-butoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

20 The product from Example 61F and 2-butanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.20 (m, 10 H), 6.97 (s, 1 H), 6.90 (d, 2 H), 6.63 (m, 1 H), 6.51 (m, 2 H), 6.10 (s, 1 H), 4.24 (m, 1 H), 4.14 (s, 4 H), 2.90 (s, 3 H), 2.30 (s, 3 H), 1.73 (m, 1 H), 1.59 (m, 1 H), 1.27 (d, 3 H), 0.95 (t, 3 H); MS (ESI+) m/z 545 (M+H)⁺.

25

Example 162

N-[3-(benzyl{4-[3-(cyclopentyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

30 The product from Example 61F and cyclopentanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 7.12-7.34 (m, 10 H), 7.01 (s, 1 H), 6.88 (s, 2 H), 6.62 (m, 1 H), 6.49 (m, 2 H), 6.07 (s, 1 H), 4.67 (m, 1 H), 4.22 (s, 4 H), 2.86 (s, 3 H), 2.28 (s, 3 H), 1.73-1.93 (m, 6 H); MS (ESI+) m/z 557 (M+H)⁺.

35

Example 163

N-[3-(benzyl{4-[3-(1-methylbutoxy)phenoxy]benzyl}amino)-2-

methylphenyl]methanesulfonamide

The product from Example 61F and 2-pentanol were processed as described in Example 159A to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.12-7.33 (m, 10 H), 6.99 (s, 1 H), 6.89 (d, 2 H), 6.62 (m, 1 H), 6.50 (m, 2 H), 6.08 (s, 1 H), 4.31 (m, 1 H), 4.15 (s, 4 H), 2.88 (s, 3 H), 2.30 (s, 3 H), 1.71 (m, 1 H), 1.40 (m, 3 H), 1.27 (d, 3 H), 0.92 (t, 3 H); MS (ESI+) m/z 559 (M+H)⁺.

Example 164

N-[3-(benzyl{4-[3-(2-methoxy-1-methylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 1-methoxy-2-propanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.18 (m, 9 H), 6.97 (br.s, 1 H), 6.89 (d, 2 H), 6.66 (m, 1 H), 6.54 (m, 2 H), 6.14 (s, 1 H), 4.47 (m, 1 H), 4.10 (s, 4 H), 3.56 (m, 1 H), 3.46 (m, 1 H), 3.39 (s, 3 H), 2.90 (s, 3 H), 2.31 (s, 3 H), 1.28 (d, 3 H); MS (ESI+) m/z 561 (M+H)⁺.

Example 165

N-[3-(benzyl{4-[3-(cyclohexyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and cyclohexanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 6.98 (br.s, 1 H), 6.89 (d, 2 H), 6.65 (m, 1 H), 6.64 (dd, 1 H), 6.50 (m, 2 H), 6.11 (s, 1 H), 4.05-4.30 (m, 5 H), 2.89 (s, 3 H), 2.29 (s, 3 H), 1.97 (m, 2 H), 1.70-1.91 (m, 2 H), 1.55 (m, 3 H), 1.33 (m, 3 H); MS (ESI+) m/z 571 (M+H)⁺.

Example 166

N-[3-[benzyl(4-{3-[(3-methylcyclopentyl)oxy]phenoxy}benzyl)amino]-2-methylphenyl]methanesulfonamide

The product from Example 61F and 3-methylcyclopentanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 6.99 (br.s, 1 H), 6.89 (d, 2 H), 6.59 (m, 1 H), 6.48 (m, 2 H), 6.08 (s, 1 H), 4.68 (m, 1 H), 4.02-4.32 (br.s, 4 H), 2.88 (s, 3 H), 2.25 (m, 4 H), 1.74-2.15 (m, 4 H), 1.38 (m, 1 H), 1.13 (m, 1 H), 1.06 (d, 1 H), 1.00 (d, 2 H); MS (ESI+) m/z 571 (M+H)⁺.

Example 167N-[3-(benzyl{4-[3-(2-ethoxy-1-methylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

5 The product from Example 61F and 1-ethoxy-2-propanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.18 (m, 9 H), 6.97 (br.s, 1 H), 6.89 (d, 2 H), 6.67 (m, 1 H), 6.54 (m, 2 H), 6.13 (s, 1 H), 4.46 (m, 1 H), 3.98-4.24 (br.s, 4 H), 3.60 (m, 1 H), 3.53 (m, 2 H), 3.47 (m, 1 H), 2.90 (s, 3 H), 2.31 (s, 3 H), 1.29 (d, 3 H), 1.18 (t, *J* = 7.0 Hz, 3 H); MS (ESI+) *m/z* 575 (M+H)⁺.

10

Example 168N-{3-[benzyl(4-{3-[(4-methylcyclohexyl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

15 The product from Example 61F and 4-methylcyclohexanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.18 (m, 9 H), 6.99 (br.s, 1 H), 6.88 (d, 2 H), 6.65 (m, 1 H), 6.51 (m, 2 H), 6.09 (s, 1 H), 4.43 (m, 1 H), 4.04-4.32 (br.s, 4 H), 2.88 (s, 3 H), 2.29 (s, 3 H), 1.97 (m, 2 H), 1.65-1.87 (m, 2 H), 1.23-1.59 (m, 4 H), 1.02 (m, 1 H), 0.93 (d, 2 H), 0.90 (d, 1 H); MS (ESI+) *m/z* 585 (M+H)⁺.

20

Example 169N-[3-(benzyl{4-[3-(cycloheptyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

25 The product from Example 61F and cycloheptanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 6.98 (br.s, 1 H), 6.89 (d, 2 H), 6.60 (m, 1 H), 6.49 (m, 2 H), 6.10 (s, 1 H), 4.35 (m, 1 H), 3.99-4.28 (br.s, 4 H), 2.89 (s, 3 H), 2.29 (s, 3 H), 1.99 (m, 2 H), 1.74 (m, 8 H), 1.43 (m, 2 H); MS (ESI+) *m/z* 585 (M+H)⁺.

30

Example 170N-(3-{benzyl[4-(3-methoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

35 The product from Example 61F and methanol were processed as described in

Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.10-7.35 (m, 9 H), 7.01 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (dd, 1 H), 6.54 (dd, 1 H), 6.47 (s, 1 H), 6.09 (s, 1 H), 3.98-4.40 (br.s, 4 H), 3.77 (s, 3 H), 2.84 (s, 3 H), 2.28 (s, 3 H); MS (ESI+) m/z 503 (M+H)⁺.

5

Example 171

N-(3-{benzyl[4-(3-ethoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 61F and ethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.20 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.63 (dd, 1 H), 6.52 (dd, 1 H), 6.49 (t, 1 H), 6.10 (s, 1 H), 4.08-4.35 (br.s, 4 H), 3.99 (dd, 2 H), 2.87 (s, 3 H), 2.27 (s, 3 H), 1.38 (t, 3 H); MS (ESI-) m/z 515 (M-H)⁻.

10

Example 172

N-(3-{benzyl[4-(3-propoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 61F and propanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.20 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (dd, 1 H), 6.52 (m, 2 H), 6.10 (s, 1 H), 4.06-4.32 (br.s, 4 H), 3.88 (t, 2 H), 2.88 (s, 3 H), 2.28 (s, 3 H), 1.70-1.93 (m, 2 H), 1.01 (t, 3 H); MS (ESI+) m/z 531 (M+H)⁺.

15

20

Example 173

N-[3-(benzyl{4-[3-(cyclopropylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and cyclopropylmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 6.98 (br.s, 1 H), 6.89 (d, 2 H), 6.63 (dd, 1 H), 6.52 (m, 2 H), 6.10 (s, 1 H), 4.00-4.32 (br.s, 4 H), 3.76 (d, 2 H), 2.88 (s, 3 H), 2.29 (s, 3 H), 1.25 (m, 1 H), 0.63 (m, 2 H), 0.32 (m, 2 H); MS (ESI-) m/z 541 (M-H)⁻.

25

30

Example 174

N-(3-{benzyl[4-(3-butoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 61F and 1-butanol were processed as described in Example 159A to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.18 (m, 9 H),

35

6.99 (br.s, 1 H), 6.89 (d, 2 H), 6.64 (dd, 1 H), 6.52 (m, 2 H), 6.10 (s, 1 H), 4.06-4.38 (br.s, 4 H), 3.92 (t, 2 H), 2.88 (s, 3 H), 2.28 (s, 3 H), 1.65-1.96 (m, 2 H), 1.46 (m, 2 H), 0.96 (t, 3 H); MS (ESI+) m/z 545 (M+H)⁺.

5

Example 175

N-(3-{benzyl[4-(3-isobutoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 61F and isobutyl alcohol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ
10 7.11-7.34 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (dd, 1 H), 6.50 (m, 2 H), 6.07 (s, 1 H), 3.99-4.39 (br.s, 4 H), 3.67 (d, 2 H), 2.86 (s, 3 H), 2.28 (s, 3 H), 2.07 (m, 1 H), 1.00 (d, 6 H); MS (ESI+) m/z 545 (M+H)⁺.

15

Example 176

N-[3-(benzyl{4-[3-(pent-3-ynloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 3-pentyn-1-ol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ
20 7.08-7.46 (m, 7 H), 6.84 (m, 3 H), 6.65 (m, 2 H), 6.52 (m, 2 H), 6.46 (s, 1 H), 6.04 (s, 1 H), 3.91-4.18 (m, 6 H), 2.76 (s, 3 H), 2.59 (m, 2 H), 2.21 (s, 3 H), 1.78 (t, 2 H); MS (ESI-) m/z 553 (M-H)⁻.

25

Example 177

N-{3-[benzyl(4-{3-[(2E)-pent-2-enyloxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 61F and 2-pentyn-1-ol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ
30 7.20 (m, 9 H), 7.04 (br.s, 1 H), 6.88 (s, 2 H), 6.65 (dd, 1 H), 6.51 (m, 2 H), 6.06 (s, 1 H), 5.63 (m, 2 H), 4.54 (d, 2 H), 3.98-4.42 (br.s, 4 H), 2.84 (s, 3 H), 2.27 (s, 3 H), 2.10 (m, 2 H), 1.00 (m, 3 H); MS (ESI+) m/z 557 (M+H)⁺.

35

Example 178

N-{3-[benzyl(4-{3-[(1-methylcyclopropyl)methoxy]phenoxy}benzyl)amino]-2-

methylphenyl}methanesulfonamide

The product from Example 61F and 2-methylcyclopropanemethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 7.01 (br.s, 1 H), 6.88 (d, 2 H), 6.63 (dd, 1 H), 6.52 (dd, 1 H), 6.49 (t, 1 H), 6.11 (s, 1 H), 4.05-4.42 (br.s, 4 H), 3.79 (m, 1 H), 3.72 (m, 1 H), 2.86 (s, 3 H), 2.26 (s, 3 H), 1.08 (d, 3 H), 0.94 (m, 1 H), 0.73 (m, 1 H), 0.47 (m, 1 H), 0.37 (m, 1 H); MS (ESI-) m/z 555 (M-H)⁻.

Example 179N-[3-(benzyl{4-[3-(cyclobutylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and cyclobutylmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.12-7.33 (m, 9 H), 7.01 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (m, 1 H), 6.51 (m, 2 H), 6.08 (s, 1 H), 3.98-4.50 (br.s, 4 H), 3.88 (d, 2 H), 2.85 (s, 3 H), 2.74 (m, 1 H), 2.27 (s, 3 H), 2.13 (m, 2 H), 1.78-2.02 (m, 4 H); MS (ESI-) m/z 555 (M-H)⁻.

Example 180N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 2-cyclopropylethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.22 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.66 (m, 1 H), 6.51 (m, 2 H), 6.09 (s, 1 H), 4.03-4.39 (br.s, 4 H), 3.99 (t, 2 H), 2.86 (s, 3 H), 2.28 (s, 3 H), 1.65 (dd, 2 H), 0.83 (m, 1 H), 0.47 (m, 2 H), 0.10 (dd, 2 H); MS (ESI+) m/z 557 (M+H)⁺.

Example 181N-[3-(benzyl{4-[3-(pentylloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 1-pentanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.18 (m, 9 H), 6.99 (br.s, 1 H), 6.89 (d, 2 H), 6.64 (m, 1 H), 6.52 (m, 2 H), 6.09 (s, 1 H), 3.99-4.33 (br.s, 4 H), 3.91 (t, 2 H), 2.88 (s, 3 H), 2.30 (s, 3 H), 1.76 (m, 2 H), 1.37 (m, 4 H), 0.92 (t, 3 H); MS (ESI-) m/z 557 (M-H)⁻.

Example 182N-[3-(benzyl{4-[3-(2-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

5 The product from Example 61F and 2-methyl-1-butanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (m, 1 H), 6.51 (m, 2 H), 6.09 (s, 1 H), 3.97-4.40 (br.s, 4 H), 3.78 (m, 1 H), 3.68 (m, 1 H), 2.87 (s, 3 H), 2.29 (s, 3 H), 1.84 (m, 1 H), 1.25 (m, 2 H), 0.99 (d, 3 H), 0.93 (t, 3 H); MS (ESI+) m/z 559 (M+H)⁺.

10

Example 183N-[3-(benzyl{4-[3-(3-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

15 The product from Example 61F and isoamyl alcohol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (m, 1 H), 6.51 (m, 2 H), 6.10 (s, 1 H), 4.00-4.40 (br.s, 4 H), 3.94 (t, 2 H), 2.87 (s, 3 H), 2.28 (s, 3 H), 1.82 (m, 1 H), 1.65 (q, 2 H), 0.94 (d, 6 H); MS (ESI+) m/z 559 (M+H)⁺.

20

Example 184N-[3-(benzyl{4-[3-(2-ethoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

25 The product from Example 61F and 2-ethoxyethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.16 (m, 9 H), 6.96 (br.s, 1 H), 6.89 (d, 2 H), 6.66 (d, 1 H), 6.55 (d, 1 H), 6.52 (s, 1 H), 6.14 (s, 1 H), 4.07 (t, 2 H), 3.94-4.25 (br.s, 4 H), 3.76 (t, 2 H), 3.58 (q, 2 H), 2.90 (s, 3 H), 2.31 (s, 3 H), 1.22 (t, 3 H); MS (ESI+) m/z 561 (M+H)⁺.

30

Example 185N-{3-[benzyl(4-{3-[2-(methylthio)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

35 The product from Example 61F and 2-(methylthio)ethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.14-7.34 (m, 9 H), 7.05 (br.s, 1 H), 6.87 (d, 2 H), 6.64 (d, 1 H), 6.54 (d, 1

H), 6.48 (s, 1 H), 6.12 (s, 1 H), 4.17–4.45 (br.s, 4 H), 4.11 (t, 2 H), 2.85 (m, 5 H), 2.24 (s, 3 H), 2.20 (s, 3 H); MS (ESI⁺) *m/z* 563 (M+H)⁺.

Example 186

5 N-[3-(benzyl){4-[3-(cyclopentylmethoxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide

The product from Example 61F and cyclopentylmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.23 (m, 9 H), 7.00 (br.s, 1 H), 6.88 (d, 2 H), 6.64 (d, 1 H), 6.51 (m, 2 H), 6.09 (s, 1 H), 3.95-4.44 (br.s, 4 H), 3.78 (d, 2 H), 2.86 (s, 3 H), 2.33 (m, 1 H), 2.27 (s, 3 H), 1.81 (m, 2 H), 1.60 (m, 4 H), 1.33 (m, 2 H); MS (ESI+) m/z 571 (M+H)⁺.

Example 187

N-[3-(benzyl{4-[3-(tetrahydrofuran-2-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and tetrahydro-2-furanmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.11-7.39 (m, 9 H), 7.06 (br.s, 1 H), 6.87 (d, 2 H), 6.65 (dd, 1 H), 6.55 (dd, 1 H), 6.42 (br.s, 1 H), 6.27 (br.s, 1 H), 4.07-4.37 (m, 4 H), 3.91 (m, 4 H), 3.81 (m, 1 H), 2.80 (s, 3 H), 2.25 (s, 3 H), 2.06 (m, 1 H), 1.92 (m, 2 H), 1.74 (m, 1 H); MS (ESI+) m/z 573 (M+H)⁺.

Example 188

25 N-[3-(benzyl{4-[3-(tetrahydrofuran-3-ylmethoxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide

The product from Example 61F and tetrahydro-3-furanmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.22 (m, 9 H), 7.01 (br.s, 1 H), 6.88 (d, 2 H), 6.63 (dd, 1 H), 6.54 (dd, 1 H), 6.49 (s, 1 H), 6.14 (s, 1 H), 4.00-4.42 (br.s, 4 H), 3.87 (m, 4 H), 3.77 (dd, 1 H), 3.68 (m, 1 H), 2.87 (s, 3 H), 2.72 (m, 1 H), 2.28 (s, 3 H), 2.10 (m, 1 H), 1.72 (m, 1 H); MS (ESI+) m/z 573 (M+H)⁺.

Example 189

35 N-[3-(benzyl{4-[3-(hexyloxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide

The product from Example 61F and 1-hexanol were processed as described in

Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.09-7.31 (m, 9 H), 6.99 (br.s, 1 H), 6.89 (d, 2 H), 6.63 (d, 1 H), 6.52 (m, 2 H), 6.10 (s, 1 H), 3.96-4.37 (br.s, 4 H), 3.91 (t, 2 H), 2.89 (s, 3 H), 2.29 (s, 3 H), 1.76 (m, 2 H), 1.43 (m, 2 H), 1.31 (m, 4 H), 0.89 (t, 3 H); MS (ESI+) m/z 574 (M+H)⁺.

Example 190

N-[3-(benzyl{4-[3-(3,3-dimethylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 3,3-dimethylbutanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.19 (m, 9 H), 6.99 (br.s, 1 H), 6.89 (d, 2 H), 6.64 (m, 1 H), 6.51 (m, 2 H), 6.10 (s, 1 H), 4.04-4.39 (br.s, 4 H), 3.98 (t, 2 H), 2.88 (s, 3 H), 2.28 (s, 3 H), 1.70 (t, 2 H), 0.97 (s, 9 H); MS (ESI+) m/z 573 (M+H)⁺.

Example 191

N-[3-(benzyl{4-[3-(2-isopropoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 2-isopropoxyethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.18 (m, 9 H), 7.00 (br.s, 1 H), 6.87 (d, 2 H), 6.66 (dd, 1 H), 6.54 (dd, 1 H), 6.51 (m, 1 H), 6.13 (s, 1 H), 4.09-4.35 (br.s, 4 H), 4.05 (m, 2 H), 3.75 (t, 2 H), 3.67 (m, 1 H), 2.87 (s, 3 H), 2.27 (s, 3 H), 1.18 (d, 6 H); MS (ESI+) m/z 575 (M+H)⁺.

Example 192

N-[3-(benzyl{4-[3-(cyclohexylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and cyclohexylmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.12-7.36 (m, 9 H), 7.03 (m, 1 H), 6.87 (d, 2 H), 6.64 (br.s., 1 H), 6.50 (m, 2 H), 6.07 (s, 1 H), 4.00-4.55 (br.s., 4 H), 3.70 (d, 2 H), 2.84 (s, 3 H), 2.26 (s, 3 H), 1.84 (m, 5 H), 1.14-1.36 (m, 4 H), 1.03 (m, 2 H); MS (ESI+) m/z 586 (M+H)⁺.

Example 193

N-[3-(benzyl{4-[3-(3-methoxy-3-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 3-methoxy-3-methyl-1-butanol were processed as described in Example 159A, 6C and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.13-7.35 (m, 9 H), 7.04 (br.s, 1 H), 6.87 (s, 2 H), 6.64 (d, 1 H), 6.52 (d, 1 H), 6.49 (s, 1 H), 6.14 (s, 1 H), 4.06-4.44 (br.s, 4 H), 4.01 (t, 2 H), 3.20 (s, 3 H), 2.84 (s, 3 H), 2.27 (s, 3 H), 1.97 (t, 2 H), 1.22 (s, 6 H); MS (ESI+) m/z 589 (M+H)⁺.

Example 194

1-(4-ethoxy-6-propylpyrimidin-2-yl)isoquinoline

The product from Example 61F and 3-pentyn-1-ol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.10-7.36 (m, 9 H), 6.99 (br.s, 1 H), 6.89 (d, 2 H), 6.64 (dd, 1 H), 6.52 (m, 2 H), 6.08 (s, 1 H), 5.61 (m, 1 H), 5.46 (m, 1 H), 3.97-4.29 (br.s, 4 H), 3.92 (t, 2 H), 2.89 (s, 3 H), 2.52 (dd, 2 H), 2.30 (s, 3 H), 1.64 (d, 3 H); MS (ESI-) m/z 555 (M-H)⁻.

Example 195

N-[3-(benzyl{4-[3-(3-furylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 3-furanmethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.46 (m, 1 H), 7.42 (m, 1 H), 7.07-7.33 (m, 9 H), 6.92 (m, 3 H), 6.70 (m, 1 H), 6.57 (m, 2 H), 6.46 (m, 1 H), 6.09 (m, 1 H), 4.89 (s, 2 H), 4.07 (d, 4 H), 2.93 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 569 (M+H)⁺.

Example 196

N-[3-(benzyl{4-[3-(2-furylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and furfuryl alcohol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.43 (dd, 1 H), 7.24 (m, 3 H), 7.12 (m, 5 H), 6.91 (m, 4 H), 6.71 (m, 1 H), 6.59 (m, 2 H), 6.40 (m, 1 H), 6.36 (m, 1 H), 6.09 (s, 1 H), 4.95 (s, 2 H), 4.04 (d, 4 H), 2.94 (s, 3 H), 2.34 (s, 3 H); MS (ESI+) m/z 569 (M+H)⁺.

Example 197N-[3-(benzyl{4-[3-(thien-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

5 The product from Example 61F and 3-thiophenemethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.32 (m, 1 H), 7.24 (m, 7 H), 7.13 (m, 4 H), 6.91 (m, 3 H), 6.70 (m, 1 H), 6.57 (m, 2 H), 6.10 (s, 1 H), 5.02 (s, 2 H), 4.06 (d, 4 H), 2.93 (s, 3 H), 2.33 (s, 3 H); MS
10 (ESI+) m/z 585 (M+H)⁺.

Example 198N-[3-(benzyl{4-[3-(2-thien-3-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

15 The product from Example 61F and 2-(3-thienyl)ethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.26 (m, 2 H), 7.20 (m, 5 H), 7.13 (m, 3 H), 7.06 (d, 1 H), 7.01 (dd, 1 H), 6.91 (m, 3 H), 6.65 (dd, 1 H), 6.54 (m, 2 H), 6.09 (s, 1 H), 4.13 (t, 2 H), 4.05 (d, 4 H), 3.10
20 (t, 2 H), 2.93 (s, 3 H), 2.32 (s, 3 H); MS (ESI+) m/z 599 (M+H)⁺.

Example 199N-[3-(benzyl{4-[3-(thien-2-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

25 The product from Example 61F and thiophene-2-methanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.31 (dd, 1 H), 7.27 (m, 1 H), 7.21 (m, 5 H), 7.13 (m, 3 H), 7.07 (m, 1 H), 6.99 (m, 1 H), 6.91 (m, 3 H), 6.72 (m, 1 H), 6.59 (m, 2 H), 6.09 (s, 1 H), 5.17 (s, 2 H), 4.06
30 (d, 4 H), 2.93 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 585 (M+H)⁺.

Example 200N-[3-(benzyl{4-[3-(2-thien-2-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

35 The product from Example 61F and 2-(2-thienyl)ethanol were processed as

described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.27 (m, 1 H), 7.20 (m, 5 H), 7.13 (m, 4 H), 6.91 (m, 5 H), 6.66 (m, 1 H), 6.55 (m, 2 H), 6.09 (s, 1 H), 4.15 (t, 2 H), 4.04 (d, 4 H), 3.28 (t, 2 H), 2.94 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 599 (M+H)⁺.

5

Example 201

N-{3-[benzyl(4-{3-[2-(4-methyl-1,3-thiazol-5-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

10 The product from Example 61F and 4-methyl-5-thiazoleethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 8.77 (s, 1 H), 7.27 (m, 1 H), 7.21 (m, 5 H), 7.13 (m, 3 H), 6.91 (m, 3 H), 6.63 (dd, 1 H), 6.57 (dd, 1 H), 6.52 (t, 1 H), 6.17 (s, 1 H), 4.11 (t, 2 H), 4.05 (d, 4 H), 3.23 (t, 2 H), 2.95 (s, 3 H), 2.45 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 614 (M+H)⁺.

15

Example 202

N-{3-[benzyl(4-{3-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

20 The product from Example 61F and N-(2-hydroxyethyl)-2-pyrrolidone were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.24 (m, 6 H), 7.13 (m, 3 H), 6.96 (d, 1 H), 6.89 (d, 2 H), 6.59 (m, 2 H), 6.45 (s, 1 H), 6.42 (t, 1 H), 4.03 (m, 6 H), 3.65 (t, 2 H), 3.55 (t, 2 H), 2.93 (s, 3 H), 2.39 (t, 2 H), 2.32 (s, 3 H), 2.01 (m, 2 H); MS (ESI+) m/z 600 (M+H)⁺.

25

Example 203

N-[3-(benzyl{4-[3-(2-morpholin-4-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

30 The product from Example 61F and N-(2-hydroxyethyl)morpholine were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.20 (m, 9 H), 7.07 (m, 1 H), 6.87 (d, 2 H), 6.79 (s, 1 H), 6.66 (dd, 1 H), 6.57 (dd, 1 H), 6.28 (t, 1 H), 4.26 (m, 2 H), 4.22 (s, 2 H), 4.17 (s, 2 H), 3.99 (s, 2 H), 3.70 (d, 2 H), 3.52 (m, 2 H), 3.05 (br.s, 2 H), 2.89 (s, 3 H), 2.33-2.66 (br.s, 2 H), 2.21 (s, 3 H);
35 MS (ESI+) m/z 602 (M+H)⁺.

Example 204N-[3-(benzyl{4-[3-(2-phenylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and 2-phenylethanol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 11 H), 7.12 (m, 3 H), 6.90 (m, 3 H), 6.64 (dd, 1 H), 6.54 (m, 2 H), 6.09 (s, 1 H), 4.13 (t, 2 H), 4.05 (d, 4 H), 3.07 (t, 2 H), 2.94 (s, 3 H), 2.32 (s, 3 H); MS (ESI+) m/z 593 (M+H)⁺.

Example 205N-{3-[4-[3-(1,3-benzodioxol-5-ylmethoxy)phenoxy]benzyl]}(benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 61F and piperonyl alcohol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.23 (m, 5 H), 7.13 (m, 4 H), 6.91 (m, 4 H), 6.84 (m, 1 H), 6.78 (d, 1 H), 6.69 (dd, 1 H), 6.57 (m, 2 H), 6.10 (s, 1 H), 5.95 (s, 2 H), 4.90 (s, 2 H), 4.08 (d, 4 H), 2.93 (s, 3 H), 2.32 (s, 3 H); MS (ESI+) m/z 623 (M+H)⁺.

Example 206N-[3-(benzyl{4-[3-(benzyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 61F and benzyl alcohol were processed as described in Examples 159A, 6C, and 6D to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.35 (m, 5 H), 7.26 (m, 6 H), 7.13 (m, 3 H), 6.91 (m, 3 H), 6.71 (dd, 1 H), 6.60 (t, 1 H), 6.57 (dd, 1 H), 6.09 (s, 1 H), 5.02 (s, 2 H), 4.05 (d, 4 H), 2.93 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 579 (M+H)⁺.

Example 207(3-{4-[(2-methylbenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83B and 2-methylbenzyl bromide were processed as described in Examples 6B, 6C, 6D, and 137B to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.93 (s, 1 H), 7.25 (m, 4 H), 7.08 (m, 5 H), 6.99 (m, 1 H), 6.92 (d, 2 H), 6.66 (m, 1 H), 6.50 (m, 2 H), 4.58 (s, 2 H), 4.06 (s, 4 H), 2.87 (s, 3 H), 2.27 (s, 3 H), 2.17 (s,

3 H); MS (ESI)⁺ m/z 559 (M-H)⁺.

Example 208

5 (3-{4-[[[(4-methylbenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83B and 4-methylbenzyl bromide were processed as described in Examples 6B, 6C, 6D, and 137B to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.94 (s, 1 H), 7.25 (m, 3 H), 7.13 (m, 2 H), 7.07 (m, 2 H), 7.02 (d, 1 H), 6.94 (m, 4 H), 6.66 (m, 1 H), 6.51 (m, 2 H), 4.64 (s, 2 H), 4.01 (s, 4 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.24 (s, 3 H); MS (ESI+) m/z 562 (M+H)⁺.

Example 209

15 (3-{4-[(2,4-dichlorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83B and 2,4-dichlorobenzyl bromide were processed as described in Examples 6B, 6C, 6D, and 137B to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.94 (s, 1 H), 7.52 (d, 1 H), 7.39 (d, 1 H), 7.32 (dd, 1 H), 7.25 (m, 3 H), 7.08 (t, 1 H), 7.01 (m, 2 H), 6.92 (d, 2 H), 6.67 (m, 1 H), 6.51 (m, 2 H), 4.64 (s, 2 H), 4.17 (s, 2 H), 4.09 (s, 2 H), 2.89 (s, 3 H), 2.30 (s, 3 H); MS (ESI+) m/z 615 (M+H)⁺.

Example 210

25 (3-{4-[(2-chloro-4-fluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83B and 2-chloro-4-fluorobenzyl bromide were processed as described in Examples 6B, 6C, 6D, and 137B to provide the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.93 (s, 1 H), 7.39 (m, 1 H), 7.33 (m, 1 H), 7.25 (m, 3 H), 7.10 (m, 2 H), 7.01 (m, 2 H), 6.92 (d, 2 H), 6.67 (m, 1 H), 6.50 (m, 2 H), 4.64 (s, 2 H), 4.16 (s, 2 H), 4.09 (s, 2 H), 2.89 (s, 3 H), 2.30 (s, 3 H); MS (ESI+) *m/z* 599 (M+H)⁺.

Example 211

35 (3-{4-[(4-benzoylbenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83B and 4-benzoylbenzyl bromide were processed as described in Examples 6B, 6C, 6D, and 137B to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 9.00 (s, 1 H), 7.68 (m, 4 H), 7.54 (t, 2 H), 7.47 (d, 2 H), 7.28 (d, 2 H), 7.24 (t, 1 H), 7.03 (m, 4 H), 6.93 (d, 2 H), 6.65 (d, 1 H), 6.50 (m, 2 H), 4.58 (s, 2 H), 4.18 (s, 2 H), 4.07 (s, 2 H), 2.92 (s, 3 H), 2.41 (s, 3 H); MS (ESI+) m/z 651 (M+H)⁺.

Example 212

(3-{4-[(3,4-difluorobenzyl){2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid

The product from Example 83B and 3,4-difluorobenzyl bromide were processed as described in Examples 6B, 6C, 6D, and 137B to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ 8.97 (s, 1 H), 7.27 (m, 5 H), 7.07 (m, 2 H), 6.95 (m, 4 H), 6.66 (d, 1 H), 6.51 (m, 2 H), 4.63 (s, 2 H), 4.05 (d, 4 H), 2.92 (s, 3 H), 2.38 (s, 3 H); MS (ESI+) m/z 583 (M+H)⁺.

Example 213

N-[3-(benzyl{4-[4-(4-hydrazino-4-oxobutyl)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide

Example-213A

tert-butyl 2-(4-(4-(4-((benzyl(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenyl)butanoyl)hydrazinecarboxyl
ate

The product from Example 100 (56mg, 0.1mmole), 1-[3-(dimethylamino)propyl]-1-ethylcarbodiimide hydrochloride (27mg, 0.14mmole) and 1-hydroxybenzotriazole hydrate (19mg, 0.14mmole) were dissolved in N,N-dimethylformamide (1.5mL) and let stand for 15 minutes at room temperature. Tert-butyl carbazate (26mg, 0.2mmole) and N,N-diisopropylethylamine (35μL, 0.2mmole) in N,N-dimethylformamide (0.5mL) were added to the reaction mixture and stirred overnight at room temperature. The reaction mixture was acidified with 1N aqueous hydrochloric acid (3mL) and extracted with dichloromethane (3mL). The organic layer was washed with saturated NaHCO₃ solution (3mL) and brine (3mL), dried (Na₂SO₄), and concentrated under reduced pressure. The crude product was purified using HPLC.

Example 213B

N-[3-(benzyl{4-[4-(4-hydrazino-4-oxobutyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The purified product from Example 213A was hydrolyzed with 10% trifluoroacetic acid in dichloromethane (2mL) at room temperature for 2 hours and concentrated under reduced pressure. The crude product was purified using HPLC to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 10.71 (s, 1 H), 8.95 (s, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.89 (m, 4 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 2.92 (s, 3 H), 2.58 (m, 2 H), 2.39 (s, 3 H), 2.22 (m, 2 H), 1.82 (m, 2 H); MS (APCI+) m/z 573 (M+H)⁺.

Example 214

N-2-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-asparagine

The product from Example 100 (56mg, 0.1mmole) and H-Asn-OtBu (38mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.11-12.80 (br.s, 1 H), 8.94 (s, 1 H), 7.99 (s, 1 H), 7.28 (m, 7 H), 7.18 (m, 3 H), 7.03 (m, 1 H), 6.96 (m, 2 H), 6.87 (m, 5 H), 4.50 (m, 1 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 2.91 (s, 3 H), 2.55 (m, 2 H), 2.44 (m, 2 H), 2.39 (s, 3 H), 2.11 (t, 2 H), 1.76 (m, 2 H); MS (APCI+) m/z 673 (M+H)⁺.

Example 215

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-valine

The product from Example 100 (56mg, 0.1mmole) and H-D-Val-OtBu (42mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.75-13.22 (br.s, 1 H), 8.95 (s, 1 H), 7.92 (d, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.04 (m, 1 H), 6.97 (m, 2 H), 6.88 (m, 4 H), 4.15 (dd, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 2.91 (s, 3 H), 2.54 (t, 2 H), 2.39 (s, 3 H), 2.19 (m, 2 H), 2.03 (t, 1 H), 1.78 (m, 2 H), 0.89 (s, 3 H), 0.87 (s, 3 H); MS (APCI+) m/z 659 (M+H)⁺.

Example 216

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-tyrosine

The product from Example 100 (56mg, 0.1mmole) and H-Tyr-OtBu (47mg,

0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.81-13.15 (br.s, 1 H), 8.95 (s, 1 H), 8.69-9.58 (br.s, 1 H), 8.03 (d, 1 H), 7.26 (m, 6 H), 7.19 (m, 1 H), 7.12 (d, 2 H), 7.00 (m, 5 H), 6.88 (m, 4 H), 6.64 (d, 2 H), 4.36 (m, 1 H), 4.05 (s, 2 H), 4.02 (s, 2 H), 2.94 (d, 1 H), 2.91 (s, 3 H), 2.72 (dd, 1 H), 2.44 (t, 2 H), 2.39 (s, 3 H), 2.06 (m, 2 H), 1.69 (m, 2 H); MS (APCI+) m/z 723 (M+H)⁺.

Example 217

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-methionine

The product from Example 100 (56mg, 0.1mmole) and H-Met-OtBu HCl (48mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.65-13.33 (br.s, 1 H), 8.95 (s, 1 H), 8.07 (d, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (m, 1 H), 6.97 (m, 2 H), 6.89 (m, 4 H), 4.30 (m, 1 H), 4.05 (s, 2 H), 4.02 (s, 2 H), 2.91 (s, 3 H), 2.55 (m, 2 H), 2.46 (m, 2 H), 2.39 (s, 3 H), 2.14 (m, 2 H), 2.03 (s, 3 H), 1.95 (m, 1 H), 1.80 (m, 3 H); MS (APCI+) m/z 646 (M+H)⁺.

Example 218

N-[2-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-lysine

The product from Example 100 (56mg, 0.1mmole) and H-Lys(Boc)-OtBu HCl (68mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.68-13.46 (br.s, 1 H), 8.95 (s, 1 H), 8.04 (d, 1 H), 7.50-7.74 (br.s, 2 H), 7.27 (m, 6 H), 7.19 (m, 3 H), 7.04 (m, 1 H), 6.97 (m, 2 H), 6.89 (m, 4 H), 4.18 (m, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 2.92 (s, 3 H), 2.75 (m, 2 H), 2.55 (t, 2 H), 2.39 (s, 3 H), 2.14 (t, 2 H), 1.64-1.85 (m, 3 H), 1.43-1.63 (m, 3 H), 1.22-1.43 (m, 2 H); MS (APCI+) m/z 687 (M+H)⁺.

Example 219

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-serine

The product from Example 100 (56mg, 0.1mmole) and H-Ser(tBu)-OtBu (51mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.72-13.24 (br.s, 1 H), 8.95 (s, 1 H), 8.28-8.51 (br.s, 1 H), 7.91 (d, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.88 (m, 4 H), 4.27 (m, 1

H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.64 (m, 2 H), 2.91 (s, 3 H), 2.56 (m, 2 H), 2.39 (s, 3 H), 2.17 (m, 2 H), 1.78 (m, 2 H); MS (APCI+) m/z 646 (M+H)⁺.

Example 220

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-phenylalanine

The product from Example 100 (56mg, 0.1mmole) and H-Phe-OtBu HCl (52mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.22-12.98 (br.s, 1 H), 8.95 (s, 1 H), 8.10 (d, 1 H), 7.22 (m, 12 H), 7.10 (d, 2 H), 7.04 (t, 1 H), 6.96 (m, 2 H), 6.87 (m, 4 H), 4.45 (m, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.06 (dd, 1 H), 2.91 (s, 3 H), 2.84 (dd, 1 H), 2.42 (t, 2 H), 2.39 (s, 3 H), 2.06 (m, 2 H), 1.68 (m, 2 H); MS (APCI+) m/z 706 (M+H)⁺.

Example 221

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-tyrosine

The product from Example 100 (56mg, 0.1mmole) and H-D-Tyr-OtBu (47mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.78-13.25 (br.s, 1 H), 8.95 (s, 1 H), 8.64-9.80 (br.s, 1 H), 8.03 (d, 1 H), 7.26 (m, 6 H), 7.19 (m, 1 H), 7.12 (d, 2 H), 7.00 (m, 5 H), 6.88 (m, 4 H), 6.64 (d, 2 H), 4.35 (m, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 2.94 (d, 1 H), 2.91 (s, 3 H), 2.72 (dd, 1 H), 2.44 (t, 2 H), 2.39 (s, 3 H), 2.06 (m, 2 H), 1.69 (m, 2 H); MS (APCI+) m/z 723 (M+H)⁺.

Example 222

N-2-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-glutamine

The product from Example 100 (56mg, 0.1mmole) and H-Gln-OtBu HCl (48mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.53-13.22 (br.s, 1 H), 8.95 (s, 1 H), 8.06 (d, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.88 (m, 4 H), 6.73 (br.s, 2 H), 4.15 (m, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 2.91 (s, 3 H), 2.54 (t, 2 H), 2.39 (s, 3 H), 2.13 (m, 4 H), 1.93 (m, 1 H), 1.75 (m, 3 H); MS (APCI+) m/z 687 (M+H)⁺.

Example 223N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-isoleucine

The product from Example 100 (56mg, 0.1mmole) and H-Ile-OtBu HCl (45mg, 0.2mmole) were processed as in Example 213A-B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.81-12.98 (br.s, 1 H), 8.95 (s, 1 H), 7.93 (d, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.04 (t, 1 H), 6.96 (m, 2 H), 6.88 (d, 4 H), 4.19 (dd, 1 H), 4.05 (s, 2 H), 4.02 (s, 2 H), 2.91 (s, 3 H), 2.53 (t, 2 H), 2.39 (s, 3 H), 2.18 (m, 2 H), 1.76 (m, 3 H), 1.39 (m, 1 H), 1.20 (m, 1 H), 0.83 (m, 6 H); MS (APCI+) m/z 673 (M+H)⁺

Example 224N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-glutamic acidExample 224A

The product from Example 100 (56mg, 0.1mmole) and L-glutamic acid dimethyl ester hydrochloride (42mg, 0.2mmole) were processed as in Example 213A to give the methyl ester intermediate.

Example 224B

The product from example 224A was hydrolyzed with a mixture of 2N aqueous sodium hydroxide solution (1mL), ethyl alcohol (0.5mL) and tetrahydrofuran (0.5mL) with stirring overnight at room temperature. The reaction mixture was acidified with 2N aqueous hydrochloric acid to pH 2~3 and extracted with ethyl acetate (3mL). The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude product was purified using HPLC to give the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.54-13.13 (br.s, 2 H), 8.95 (s, 1 H), 8.05 (d, 1 H), 7.26 (m, 6 H), 7.19 (m, 3 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.89 (d, 4 H), 4.21 (m, 1 H), 4.04 (d, 4 H), 2.91 (s, 3 H), 2.54 (m, 2 H), 2.39 (s, 3 H), 2.28 (m, 2 H), 2.14 (t, 2 H), 1.96 (m, 1 H), 1.78 (m, 3 H); MS (APCI+) m/z 689 (M+H)⁺.

Example 225N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-histidine

The product from Example 100 (56mg, 0.1mmole) and H-D-His-OMe.2HCl (48mg, 0.2mmole) were processed as in Examples 213A and 224B to provide the title compound.

¹H NMR (500 MHz, DMSO-d₆) δ 13.88-14.43 (m, 1 H), 8.96 (d, 2 H), 8.20 (d, 1 H), 7.38 (s, 1 H), 7.26 (m, 5 H), 7.20 (m, 1 H), 7.13 (d, 4 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.88 (m, 4 H), 4.43-4.70 (m, 1 H), 4.04 (d, 4 H), 3.14 (m, 1 H), 2.98 (m, 1 H), 2.92 (s, 3 H), 2.39 (s, 3 H), 2.36 (m, 2 H), 2.09 (m, 2 H), 1.71 (m, 2 H); MS (APCI+) m/z 696 (M+H)⁺.

5

Example 226

1-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-proline

10 The product from Example 100 (56mg, 0.1mmole) and L-Proline methyl ester hydrochloride (33mg, 0.2mmole) were processed as in Examples 213A and 224B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.66-13.14 (br.s, 1 H), 8.95 (s, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.88 (m, 4 H), 4.22 (dd, 1 H), 4.03 (d, 4 H), 3.46 (m, 2 H), 2.91 (s, 3 H), 2.58 (t, 2 H), 2.39 (s, 3 H), 2.27 (t, 2 H), 1.97-
15 2.24 (m, 2 H), 1.70-1.95 (m, 4 H); MS (APCI+) m/z 656 (M+H)⁺.

Example 227

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-valine

20 The product from Example 100 (56mg, 0.1mmole) and L-Valine methyl ester hydrochloride (34mg, 0.2mmole) were processed as in Examples 213A and 224B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.28-12.62 (br.s, 1 H), 8.94 (s, 1 H), 7.92 (d, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.04 (m, 1 H), 6.96 (m, 2 H), 6.88 (m, 4 H), 4.15 (dd, 1 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 2.91 (s, 3 H), 2.54 (t, 2 H), 2.39 (s, 3 H), 2.19 (td, 2
25 H), 2.02 (m, 1 H), 1.78 (m, 2 H), 0.88 (s, 3 H), 0.87 (s, 3 H); MS (APCI+) m/z 658 (M+H)⁺.

Example 228

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-aspartic acid

30 The product from Example 100 (56mg, 0.1mmole) and L-Aspartic acid dimethyl ester hydrochloride (40mg, 0.2mmole) were processed as in Examples 213A and 224B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.46-13.35 (br.s, 2 H), 8.95 (s, 1 H), 8.13 (d, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.03 (m, 1 H), 6.96 (m, 2 H), 6.88 (m, 4 H), 4.54 (m, 1 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 2.91 (s, 3 H), 2.69 (m, 1 H), 2.55 (m, 3 H),
35 2.39 (s, 3 H), 2.12 (t, 2 H), 1.76 (m, 2 H); MS (APCI+) m/z 674 (M+H)⁺.

Example 229

4-{[4-(4-{4-[(benzyl{2-methyl-3-
 5 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]amino}-1-
 methyl-1H-pyrrole-2-carboxylic acid

The product from Example 100 (56mg, 0.1mmole) and 4-Amino-1-methyl-1H-pyrrole-2-carboxylic acid methyl ester hydrochloride (38mg, 0.2mmole) were processed as in Examples 213A and 224B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.62-12.57 (br.s, 1 H), 9.74 (s, 1 H), 8.95 (s, 1 H), 7.24 (m, 10 H), 7.04 (m, 1 H), 6.96
 10 (m, 2 H), 6.89 (m, 4 H), 6.65 (d, 1 H), 4.05 (s, 2 H), 4.02 (s, 2 H), 3.79 (s, 3 H), 2.91 (s, 3 H), 2.57 (t, 2 H), 2.39 (s, 3 H), 2.23 (t, 2 H), 1.84 (m, 2 H); MS (APCI+) m/z 681 (M+H)⁺.

Example 230

4-(4-{4-[(benzyl{2-methyl-3-
 15 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-oxotetrahydrofuran-
 3-yl)butanamide

The product from Example 100 (56mg, 0.1mmole) and α-Amino-γ-butyrolactone hydrobromide (36mg, 0.2mmole) were processed as in Examples 213A and 224B to provide
 20 the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 8.31 (d, 1 H), 7.26 (m, 6 H), 7.18 (m, 3 H), 7.04 (m, 1 H), 6.97 (m, 2 H), 6.88 (m, 4 H), 4.53 (dd, 1 H), 4.34 (t, 1 H), 4.20 (m, 1 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.17 (s, 3 H), 2.91 (s, 3 H), 2.55 (t, 2 H), 2.39 (m, 2 H), 2.13 (t, 2 H), 1.67-1.90 (m, 2 H); MS (APCI+) m/z 642 (M+H)⁺.

25

Example 231

N-(3-((2,4-difluorobenzyl)(4-(3-(((2S,4R)-6-oxo-4-hydroxytetrahydro-2H-pyran-2-
 yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide

A solution of the product from Example 136B (.2098 g, .274 mmoles) and 3N HCl
 30 (.1 mL) in THF (.68 mL) and EtOH (1.4 mL) was stirred for 16 h at room temperature. The reaction mixture was diluted with pH 7 buffer (10 mL) and extracted with EtOAc (24 mL). The organic layer was rinsed with brine, dried over Na₂SO₄, and concentrated *in vacuo*. The crude products were dissolved in CH₂Cl₂ (.55 mL), cooled to 0 °C, and treated with TFA (.15 mL). After warming to room temperature and stirring for 2 h, the reaction was
 35 cooled to 0 °C and quenched with NaHCO₃ (.136 g). The crude products were diluted with EtOAc, extracted with pH 7 buffer, rinsed with brine, and dried over Na₂SO₄. After concentration *in vacuo*, purification by preparative HPLC (CH₃CN:10 mM NH₄OAc in

H₂O) on a YMC ODS Guardpak column yielded the titled compound (.1321g, 74%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.16 (m, 5 H), 6.96 (m, 1 H), 6.90 (m, 2 H), 6.74 (m, 2 H), 6.61 (dt, 2 H), 6.43 (t, 1 H), 6.30 (s, 1 H), 5.04 (m, 1 H), 4.47 (m, 1 H), 4.09 (s, 2 H), 4.07 (m, 2 H), 4.04 (s, 2 H), 2.91 (s, 3 H), 2.76 (dd, 1 H), 2.67 (m, 1 H), 2.27 (s, 3 H), 2.04 (m, 2 H), 1.60 (br.s, 2 H); MS (ESI) m/z 653 (M+H⁺).

434889 Example 232 James Link

ethyl 4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoate

Example 232A

ethyl 4-(4-(4-((benzyl(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenoxy)butanoate

The product from Example 60E was processed as described in Example 116A to provide the titled compound.

Example 232B

ethyl 4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoate

The product from Example 232A was processed as described in Examples 6C and 6D to provide the titled compound. ¹H NMR (300 MHz, CDCl₃) δ 7.21 (m, 5 H), 7.10 (m, 3 H), 6.87 (m, 8 H), 6.10 (s, 1 H), 4.15 (q, 2 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.98 (t, 2 H), 2.94 (s, 3 H), 2.52 (t, 2 H), 2.33 (s, 3 H), 2.11 (m, 2 H), 1.26 (t, 3 H); MS (ESI) m/z 603 (M+H⁺).

Example 233

4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid

The product from Example 232B was processed as described in Example 50 to provide the titled compound. ¹H NMR (300 MHz, CDCl₃) δ 7.05-7.30 (m, 9 H), 6.76-6.97 (m, 7 H), 6.23 (s, 1 H), 3.88-4.19 (m, 7 H), 2.91 (s, 3 H), 2.59 (t, 2 H), 2.31 (s, 3 H), 2.13 (m, 2 H); MS (ESI) m/z 575 (M+H⁺).

Example 234

5-(4-{4-[(benzyl{2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid

Example 234A

5 ethyl 5-(4-(4-((benzyl(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenoxy)pentanoate

The product from Example 60E and ethyl-5-bromovalerate was processed as described in Example 116A to provide the titled compound.

Example 234B

10 5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid

The product from Example 234A was processed as described in Examples 6C, 6D, and 50 to provide the titled compound. ¹H NMR (300 MHz, CDCL₃) δ 7.22 (m, 8 H), 7.11 (m, 2 H), 6.92 (m, 2 H), 6.82 (m, 4 H), 6.15 (s, 1 H), 4.11 (s, 2 H), 4.06 (s, 2 H), 3.95 (m, 2 H), 2.92 (s, 3 H), 2.46 (m, 2 H), 2.36 (br.s, 1 H), 2.29 (s, 3 H), 1.85 (m, 4 H); MS (APCI) m/z 589 (M+H⁺).

Example 235

20 N-(3-(benzyl(4-(4-(((2S, 4R)-6-oxo-4hydroxytetrahydro-2H-pyran-2-yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide

The product from Example 60G was processed as described in Example 231 to provide the titled compound. ¹H NMR (400 MHz, CDCL₃) δ 7.23 (m, 7 H), 7.10 (m, 3 H), 6.88 (m, 6 H), 6.28 (s, 1 H), 5.05 (m, 1 H), 4.16 (dd, 1 H), 4.09 (dd, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 2.94 (s, 3 H), 2.77 (dd, 1 H), 2.68 (dd, 1 H), 2.34 (s, 3 H), 2.09 (dd, 2 H), 1.99 (s, 1 H), 1 (m, 1 H); MS (ESI) m/z 617 (M+H⁺).

Example 236

30 (5-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-ethylphenoxy)acetic acid

Example 236A

3-(allyloxy)-4-ethylphenyl 4-methylbenzenesulfonate

35 4-Ethyl resorcinol was processed as described in Example 63A.

Example 236B

3-(allyloxy)-4-ethylphenol

The product from Example 236A was processed as described in Example 63B.

Example 236C

5 4-(3-(allyloxy)-4-ethylphenoxy)benzaldehyde

The product from Example 236B and 4-fluorobenzaldehyde were processed as in Example 61C to provide the titled compound. MS (DCI) m/z 289 (M+H)⁺.

Example 236D

10 N-(4-(3-(allyloxy)-4-ethylphenoxy)benzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 236C and 2-methyl-3-nitroaniline were processed as in Example 6A to provide the titled compound. MS (ESI-) m/z 423 (M-H)⁻.

15 Example 236E

The product from Example 236D and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the titled compound. MS (ESI+) m/z 573 (M+H)⁺.

20 Example 236F

The product from Example 236E was processed as described in Example 33A to provide the titled compound.

25 Example 236G

The product from Example 236F and ethyl glycolate were processed as described in Example 62A to provide the titled compound. MS (ESI+) m/z 597 (M+H)⁺.

30 Example 236H

The product from Example 236G was processed as described in Example 6C and D to provide the titled compound.

Example 236I

(5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-ethylphenoxy)acetic acid

The product from Example 236H was processed as described in Example 50 to provide the titled compound. ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.40 (m, 5 H), 6.95 (m, 3 H), 6.77 (d, *J* = 8.5 Hz, 2 H), 6.65 (m, 3 H), 6.43 (d, *J* = 2.4 Hz, 1 H), 4.60 (s, 2 H), 4.31 (s, 2 H), 4.13 (s, 2 H), 2.90 (s, 3 H), 2.50-2.80 (m, 2 H), 1.99 (s, 3 H), 1.21 (t, *J* = 7.5 Hz, 3 H); MS (ESI+) *m/z* 611 (M+H)⁺.

Example 237

ethyl 4-{[4-(3-{4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]amino}butanoate

e

The product from Example 116B and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 68D to provide the titled compound. ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 7 H), 7.15 (m, 3 H), 6.94 (d, 1 H), 6.89 (d, 2 H), 6.61 (dd, 1 H), 6.56 (dd, 1 H), 6.45 (t, 1 H), 6.31 (s, 1 H), 5.77 (s, 1 H), 4.12 (q, 2 H), 4.06 (s, 2 H), 4.02 (s, 2 H), 3.93 (t, 2 H), 3.28 (dd, 2 H), 2.93 (s, 3 H), 2.35 (m, 4 H), 2.33 (s, 3 H), 2.05 (m, 2 H), 1.80 (m, 2 H), 1.24 (t, 3 H); MS (ESI+) *m/z* 688 (M+H)⁺.

Example 238

(5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-hexylphenoxy)acetic acid

Example 238A

3-(allyloxy)-4-hexylphenyl 4-methylbenzenesulfonate

4-Hexylresorcinol was processed as described in Example 63A.

Example 238B

3-(allyloxy)-4-hexylphenol

The product from Example 238A was processed as described in Example 63B.

Example 238C

4-(3-(allyloxy)-4-hexylphenoxy)benzaldehyde

The product from Example 238B and 4-fluorobenzaldehyde were processed as in Example 61C to provide the titled compound. MS (DCI) m/z 289 (M+H)⁺.

5

Example 238D

(2-hexyl-5-(4-(((2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenoxy)acetic acid

The product from Example 238C and 2-methyl-3-nitroaniline were processed as in Example 6A to provide the titled compound. MS (ESI-) m/z 423 (M-H)⁻.

10

Example 238E

N-(4-(3-(allyloxy)-4-hexylphenoxy)benzyl)-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 238D and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the titled compound. MS (ESI+) m/z 573 (M+H)⁺.

15

Example 238F

5-(4-(((2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)-2-hexylphenol

20

The product from Example 238E was processed as described in Example 33A to provide the titled compound.

Example 238G

ethyl (5-(4-(((2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)-2-hexylphenoxy)acetate

25

The product from Example 238F and ethyl glycolate were processed as described in Example 62A to provide the titled compound. MS (ESI+) m/z 597 (M+H)⁺.

30

Example 238H

ethyl (5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-hexylphenoxy)acetate

35

The product from Example 238G was processed as described in Example 6C and D to provide the titled compound.

Example 238I

(5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-hexylphenoxy)acetic acid

The product from Example 238H was processed as described in Example 50 to provide the titled compound. ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 2 H), 7.17 (t, 1 H), 7.09 (d, 2 H), 7.03 (d, 2 H), 6.80 (d, 2 H), 6.75 (m, 3 H), 6.60 (dd, 1 H), 6.42 (d, 1 H), 4.59 (s, 2 H), 4.23 (s, 2 H), 4.10 (s, 2 H), 2.90 (s, 3 H), 2.60 (t, 2 H), 2.08 (s, 3 H), 1.55 (m, 2 H), 1.30 (m, 6 H), 0.88 (t, 3 H); MS (ESI+) m/z 667 (M+H)⁺.

Example 239

ethyl N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-N-
methylglycinate

The product from Example 116B and sarcosine hydrochloride were processed as described in Example 68D to provide the titled compound. ¹H NMR (500 MHz, CDCl₃) δ 7.25 (m, 7 H), 7.15 (m, 3 H), 6.95 (m, 1 H), 6.90 (d, 2 H), 6.63 (dd, 1 H), 6.55 (dd, 1 H), 6.50 (m, 1 H), 6.32 (s, 1 H), 4.17 (q, 2 H), 4.06 (s, 2 H), 4.05 (m, 2 H), 4.02 (s, 2 H), 3.95 (m, 2 H), 3.07 (s, 3 H), 2.93 (s, 3 H), 2.57 (t, 2 H), 2.34 (s, 3 H), 2.15 (m, 2 H), 1.26 (t, 3 H); MS (ESI+) m/z 674 (M+H)⁺.

Example 240

N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-N-
methylglycine

The product from Example 239 was processed as described in Example 50 to provide the titled compound. ¹H NMR (500 MHz, CDCl₃) δ 7.32 (d, 1 H), 7.25 (m, 9 H), 7.15 (m, 2 H), 6.86 (d, 2 H), 6.60 (m, 2 H), 6.26 (t, 1 H), 4.47 (br.s, 4 H), 4.05 (s, 2 H), 3.82 (t, 2 H), 3.07 (s, 3 H), 2.75 (s, 3 H), 2.57 (t, 2 H), 2.10 (m, 2 H), 2.08 (s, 3 H); MS (ESI+) m/z 646 (M+H)⁺.

Example 241

ethyl (2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetate

The product from Example 91G was processed as described in Example 6C and D to

provide the titled compound. ^1H NMR (500 MHz, CDCl_3) δ 7.29 (d, 1 H), 7.20 (m, 3 H), 7.10 (m, 2 H), 6.91 (d, 1 H), 6.91 (d, 2 H), 6.75 (m, 2 H), 6.50 (m, 2 H), 6.22 (s, 1 H), 4.63 (s, 2 H), 4.23 (q, 2 H), 4.07 (s, 2 H), 4.04 (s, 2 H), 2.95 (s, 3 H), 2.31 (s, 3 H), 1.25 (t, 3 H); MS (ESI+) m/z 645 ($\text{M}+\text{H}$) $^+$.

5

Example 242

N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

10

Example 242A

N-di-(3-(benzyl(4-(3-(2-cyclopropylethoxy)benzoyl)benzyl)amino)-2-methylphenyl)methanesulfonamide

15

The product from Example 88F (50mg, 0.087 mmoles) in anhydrous tetrahydrofuran (0.5 ml) was treated with triphenylphosphine (46 mg, 0.174 mmoles) in anhydrous tetrahydrofuran (0.1 ml), di-*t*-butylazodicarboxylate (30 mg, 0.131 mmoles) in anhydrous tetrahydrofuran (0.1 ml), 2-cyclopropylethanol (9.26 mg, 0.109 mmoles) in anhydrous tetrahydrofuran (0.109 ml) at room temperature overnight. Di-*t*-butylazodicarboxylate (30 mg, 0.131 mmoles) in anhydrous tetrahydrofuran (0.1 ml) and 2-cyclopropylethanol (9.26 mg, 0.109 mmoles) in anhydrous tetrahydrofuran (0.109 ml) were added and reaction was continued at room temperature overnight. Reaction mixture was diluted with ethyl acetate and washed with water and brine, dried over sodium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue was purified by HPLC to provide the titled compound.

25

Example 242B

N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

30

The product from Example 242A was treated with 2 N aqueous sodium hydroxide: 1 tetrahydrofuran: 1 ethanol (4 ml) at room temperature overnight. The reaction mixture acidified with 1N hydrochloric acid, extracted with ethyl acetate. The organic layer dried over anhydrous sodium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue purified by HPLC to provide the titled compound. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.96 (s, 1 H), 7.66 (d, 2 H), 7.45 (m, 3 H), 7.28 (d, 4 H), 7.21 (m, 4 H), 7.01 (m, 3 H), 4.17 (s, 2 H), 4.09 (s, 2 H), 4.07 (t, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H), 1.63 (dd, 2 H), 0.83 (m, 1 H), 0.42 (m, 2 H), 0.11 (m, 2 H); MS (ESI+) m/z 569 ($\text{M}+\text{H}$) $^+$.

35

Example 243N-[3-(benzyl{4-[3-(cyclopentylmethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

5 The product from Example 88F and cyclopentylmethanol were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.65 (d, 2 H), 7.45 (m, 3 H), 7.28 (d, 4 H), 7.20 (m, 4 H), 7.01 (m, 3 H), 4.17 (s, 2 H), 4.09 (s, 2 H), 3.89 (d, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H), 2.29 (m, 1 H), 1.77 (m, 2 H), 1.47-1.68 (m, 4 H), 1.33 (m, 2 H); MS (ESI+) m/z 583 (M+H)⁺.

10

Example 244N-[3-(benzyl{4-[3-(tetrahydrofuran-2-ylmethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

15 The product from Example 88F and tetrahydro-2-furanmethanol were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.66 (d, 2 H), 7.45 (m, 3 H), 7.29 (d, 4 H), 7.22 (m, 4 H), 7.01 (m, 3 H), 4.17 (s, 2 H), 4.14 (m, 1 H), 4.09 (s, 2 H), 3.99 (m, 2 H), 3.77 (m, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H), 1.99 (m, 1 H), 1.85 (m, 2 H), 1.68 (m, 1 H); MS (ESI+) m/z 585 (M+H)⁺.

20

Example 245N-[3-(benzyl{4-[3-(tetrahydrofuran-3-ylmethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

25 The product from Example 88F and tetrahydro-3-furanmethanol were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.66 (d, 2 H), 7.45 (m, 3 H), 7.29 (d, 4 H), 7.22 (m, 4 H), 7.01 (m, 3 H), 4.17 (s, 2 H), 4.09 (s, 2 H), 3.71-4.05 (m, 4 H), 3.64 (m, 1 H), 3.53 (m, 1 H), 2.92 (s, 3 H), 2.66 (m, 1 H), 2.42 (s, 3 H), 2.01 (m, 1 H), 1.66 (m, 1 H); MS (ESI+) m/z 585 (M+H)⁺.

30

Example 246N-[3-(benzyl{4-[3-(2-phenylethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

35 The product from Example 88F and 2-phenyl ethanol were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.65 (d, 2 H), 7.44 (m, 3 H), 7.30 (m, 8 H), 7.20 (m, 5 H), 7.01 (m, 3 H), 4.25 (t, 2 H), 4.16 (s, 2 H), 4.09 (s, 2 H), 3.04 (t, 2 H), 2.91 (s, 3 H), 2.41 (s, 3 H); MS (ESI+) m/z 605 (M+H)⁺.

Example 247N-[3-(benzyl{4-[3-(benzyloxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide

5 The product from Example 88F and benzyl alcohol were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.62 (d, 2 H), 7.45 (m, 5 H), 7.39 (m, 2 H), 7.31 (m, 7 H), 7.22 (m, 2 H), 7.01 (m, 3 H), 5.17 (s, 2 H), 4.17 (s, 2 H), 4.09 (s, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H); MS (ESI+) m/z 591 (M+H)⁺.

Example 248N-[3-[benzyl(4-{3-[2-(2-oxopyrrolidin-1-yl)ethoxy]benzoyl}benzyl)amino]-2-methylphenyl]methanesulfonamide

15 The product from Example 88F and N-(2-hydroxyethyl)-2-pyrrolidone were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.66 (d, 2 H), 7.45 (m, 3 H), 7.28 (d, 4 H), 7.21 (m, 4 H), 7.01 (m, 3 H), 4.17 (s, 2 H), 4.13 (t, 2 H), 4.09 (s, 2 H), 3.55 (t, 2 H), 3.44 (t, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H), 2.20 (t, 2 H), 1.90 (m, 2 H); MS (ESI+) m/z 612 (M+H)⁺.

Example 249N-[3-[benzyl(4-{3-[2-(4-methyl-1,3-thiazol-5-yl)ethoxy]benzoyl}benzyl)amino]-2-methylphenyl]methanesulfonamide

25 The product from Example 88F and 4-methyl-5-thiazoleethanol were processed as described in Example 242A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 8.82 (s, 1 H), 7.65 (d, 2 H), 7.45 (m, 3 H), 7.28 (d, 4 H), 7.21 (m, 4 H), 7.01 (m, 3 H), 4.20 (t, 2 H), 4.17 (s, 2 H), 4.09 (s, 2 H), 3.23 (t, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H), 2.34 (s, 3 H); MS (ESI+) m/z 626 (M+H)⁺.

Example 250N-(3-{benzyl[4-(4-{[(2R)-2,3-dihydroxypropyl]oxy}benzoyl)benzyl]amino}-2-methylphenyl)methanesulfonamideExample 250A

1-(allyloxy)-4-bromobenzene

4-Bromophenol was processed as described in Example 88C to provide the title compound.

5

Example 250B

N-(3-((4-(4-(allyloxy)benzoyl)benzyl)(benzyl)amino)-2-methylphenyl)methanesulfonamide

10 The product Example 88B was processed as described in Example 88D using the product from Example 250A instead of 1-(allyloxy)-3-bromobenzene to provide the title compound.

Example 250C

15 di-N-(3-((4-(4-(allyloxy)benzoyl)benzyl)(benzyl)amino)-2-methylphenyl)methanesulfonamide

The product from Example 250B was processed as described in Example 6D to provide the title compound.

20

Example 250D

di-N-(3-(benzyl(4-(4-hydroxybenzoyl)benzyl)amino)-2-methylphenyl)methanesulfonamide

The product from Example 250C was processed as described in Example 33A to provide the title compound.

25

Example 250E

N-(3-{benzyl[4-(4-{[(2R)-2,3-dihydroxypropyl]oxy}benzoyl)benzyl]amino}-2-methylphenyl)methanesulfonamide

30 The product from Example 250D (50mg, 0.087mmole), (4S)-2,2-dimethyl-1,3-dioxolane-4-methanol (79.3mg, 0.6mmole), di-tert-butyl azodicarboxylate (30mg, 0.13mmole), and triphenylphosphine (45mg, 0.17mmole) were dissolved in tetrahydrofuran (2mL) in a capped test-tube and shaken at room temperature overnight. Ethyl acetate (2mL) was added to dilute the reaction mixture. Then, the diluted organic solution was washed with water (2mL) and brine (2mL), dried (Na₂SO₄), concentrated *in vacuo*, and purified by
35 HPLC to give the ethylene glycol protected compound. To the ethylene glycol protected compound was added 2N NaOH solution (2mL), tetrahydrofuran (1mL), and ethyl alcohol (1mL). The mixture was shaken at room temperature overnight. Then 1N HCl solution was

used to adjust the PH to 2~3. Ethyl acetate (4mL) was used to extract the product. The extraction solution was concentrated *in vacuo* and purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO- d₆) δ ppm 8.96 (s, 1 H), 7.70 (m, 2 H), 7.60 (m, 2 H), 7.44 (m, 2 H), 7.28 (m, 4 H), 7.21 (m, 1 H), 7.03 (m, 5 H), 4.17 (s, 2 H), 4.11 (m, 1 H), 4.10 (s, 2 H), 3.97 (dd, 1 H), 3.82 (m, 1 H), 3.46 (d, 2 H), 2.92 (s, 3 H), 2.42 (s, 3 H); MS (APCI+) m/z 575 (M+H)⁺.

Example 251

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]glycine

Example 251A

methyl N-(4-(4-(4-((benzyl(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl]glycinate

The product from Example 233 (28.7 mg, 0.05 mmoles) in anhydrous dimethylformamide (0.5 ml) was treated with 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (14 mg, 0.07 mmoles) and 1-hydroxybenzotriazole hydrate (10 mg, 0.07 mmoles) at room temperature for 15 minutes. A mixture of glycine methyl ester hydrochloride (12.6 mg, 0.1mmole) and N,N-diisopropylethylamine (0.014 ml, 0.1 mmole) in anhydrous dimethylformamide (0.235 ml) was added to the reaction and shaken at room temperature overnight. The reaction mixture was acidified with 1N aqueous hydrochloric acid and extracted with dichloromethane. The organic layer was washed with saturated aqueous sodium bicarbonate solution and brine, dried over anhydrous sodium sulfate, filtered and filtrate concentrated *in vacuo*. The residue purified by HPLC to provide the titled compound.

Example 251B

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]glycine

The product from Example 251A was treated with 2 2N aqueous sodium hydroxide: 1 tetrahydrofuran: 1 ethanol (2 ml) at room temperature overnight. The reaction mixture acidified with 1N hydrochloric acid to pH 3-4, extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, filtered and the filtrate concentrated *in vacuo*. The residue purified by HPLC to provide the titled compound. ¹H NMR (500 MHz,

DMSO-d₆) δ 12.13-12.81 (br.s, 1 H), 8.94 (s, 1 H), 8.18 (t, 1 H), 7.14-7.34 (m, 7 H), 6.98 (m, 7 H), 6.82 (d, 2 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.95 (t, 2 H), 3.74 (d, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.30 (t, 2 H), 1.93 (m, 2 H); MS (ESI+) m/z 632 (M+H)⁺.

5

Example 252

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-beta-alanine

The product from Example 233 and beta-alanine ethyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.91-12.41 (br.s, 1 H), 8.94 (s, 1 H), 7.90 (m, 1 H), 7.22 (m, 7 H), 6.88-7.09 (m, 7 H), 6.82 (d, 2 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.92 (t, 2 H), 3.24 (dd, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.37 (t, 2 H), 2.21 (t, 2 H), 1.91 (m, 2 H); MS (ESI+) m/z 646 (M+H)⁺.

15

Example 253

4-{[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]amino}butanoic acid

The product from Example 233 and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.75-12.25 (br.s, 1 H), 8.94 (s, 1 H), 7.83 (t, 1 H), 7.24 (m, 7 H), 6.98 (m, 7 H), 6.82 (d, 2 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.93 (t, 2 H), 3.05 (dd, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.20 (m, 4 H), 1.91 (m, 2 H), 1.62 (m, 2 H); MS (ESI+) m/z 660 (M+H)⁺.

25

Example 254

N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-N-methylglycine

30

The product from Example 233 and N-methylglycine ethyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.13-13.26 (br.s, 1 H), 8.94 (s, 1 H), 7.23 (m, 7 H), 6.99 (m, 7 H), 6.82 (d, J = 8.7 Hz, 2 H), 4.12 (s, 1 H), 4.04 (s, 2 H), 3.96 (m, 5 H), 3.01 (s, 3 H), 3.01 (s, 2 H), 2.82 (s, 1 H), 2.50 (m, 1 H), 2.37 (m, 4 H), 1.92 (m, 2 H); MS (ESI+) m/z 646 (M+H)⁺.

35

Example 255

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamic
acid

The product from Example 233 and L-glutamic acid dimethyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.33-12.71 (br.s, 1 H), 11.89-12.26 (br.s, 1 H), 8.94 (s, 1 H), 8.12 (d, 1 H), 7.23 (m, 7 H), 6.99 (m, 7 H), 6.82 (d, 2 H), 4.22 (m, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.94 (t, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.28 (m, 4 H), 1.95 (m, 3 H), 1.77 (m, 1 H); MS (ESI+) m/z 704 (M+H)⁺.

Example 256

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-valine

The product from Example 233 and L-valine methyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.18-12.73 (br.s, 1 H), 8.94 (s, 1 H), 7.98 (d, 1 H), 7.23 (m, 7 H), 6.97 (m, 7 H), 6.82 (d, 2 H), 4.14 (m, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.94 (t, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.34 (m, 2 H), 2.03 (m, 1 H), 1.92 (m, 2 H), 0.87 (dd, 6 H); MS (ESI+) m/z 674 (M+H)⁺.

Example 257

4-{[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]amino}-1-
methyl-1H-pyrrole-2-carboxylic acid

The product from Example 233 and 4-amino-1-methyl-1H-pyrrole-2-carboxylic acid methyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.89-12.33 (br.s, 1 H), 9.81 (s, 1 H), 8.94 (s, 1 H), 7.24 (m, 8 H), 6.98 (m, 7 H), 6.82 (d, 2 H), 6.65 (d, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.97 (t, 2 H), 3.79 (s, 3 H), 2.91 (s, 3 H), 2.38 (m, 5 H), 1.99 (m, 2 H); MS (ESI+) m/z 697 (M+H)⁺.

Example 258

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-serine

Example 258A

tert-butyl N-(4-(4-(4-((benzyl(2-methyl-3-
5 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-L-serinate

The product from Example 233 and O-tert-butyl L-serine tert-butyl ester hydrochloride were processed as described in Example 251A to provide the titled compound.

10 Example 258B

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-serine

The product from Example 258A was treated with 10% trifluoroacetic acid in dichloromethane (2 ml) for 5 hours at room temperature and concentrated in vacuo. Residue
15 purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.04-12.95 (br.s, 1 H), 8.94 (s, 1 H), 8.00 (d, 1 H), 7.24 (m, 7 H), 6.88-7.09 (m, 7 H), 6.82 (d, 2 H), 4.29 (m, 1 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.95 (t, 2 H), 3.65 (m, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.32 (m, 2 H), 1.94 (m, 2 H); MS (ESI+) m/z 662 (M+H)⁺.

20

Example 259

N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-isoleucine

The product from Example 233 and L-isoleucine tert-butyl ester hydrochloride were
25 processed as described in Example 251A and 258B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.06-12.82 (br.s, 1 H), 8.94 (s, 1 H), 7.99 (d, 1 H), 7.23 (m, 7 H), 6.87-7.09 (m, 7 H), 6.82 (d, 2 H), 4.19 (dd, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.93 (m, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.33 (m, 2 H), 1.92 (m, 2 H), 1.76 (m, 1 H), 1.40 (m, 1 H), 1.18 (m, 1 H), 0.83 (m, 6 H); MS (ESI+) m/z 688 (M+H)⁺.

30

Example 260

N-[2-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamine

35 The product from Example 233 and L-glutamine tert-butyl ester hydrochloride were processed as described in Example 251A and 258B to provide the titled compound. ¹H

NMR (500 MHz, DMSO- d_6) δ 12.00-12.89 (br.s, 1 H), 8.94 (s, 1 H), 8.13 (d, 1 H), 7.24 (m, 8 H), 6.98 (m, 7 H), 6.82 (d, 2 H), 6.73 (s, 1 H), 4.16 (m, 1 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.95 (m, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.29 (t, 2 H), 2.12 (m, 2 H), 1.94 (m, 3 H), 1.74 (m, 1 H); MS (ESI+) m/z 703 (M+H)⁺.

5

Example 261

N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]glycine

10

The product from Example 234B and glycine methyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO- d_6) δ 12.13-12.77 (br.s, 1 H), 8.94 (s, 1 H), 8.12 (t, 1 H), 7.24 (m, 7 H), 7.03 (t, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.93 (t, 2 H), 3.73 (d, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.19 (t, 2 H), 1.55-1.79 (m, 4 H); MS (ESI+) m/z 646 (M+H)⁺.

15

Example 262

N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-beta-alanine

20

The product from Example 234B and beta-alanine ethyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO- d_6) δ 11.84-12.51 (br.s, 1 H), 8.94 (s, 1 H), 7.85 (t, 1 H), 7.22 (m, 7 H), 7.03 (t, $J = 7.8$ Hz, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.92 (t, 2 H), 3.23 (dd, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.36 (t, 2 H), 2.11 (t, 2 H), 1.64 (m, 4 H); MS (ESI+) m/z 660 (M+H)⁺.

25

Example 263

4-{[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]amino}butanoic acid

30

The product from Example 234B and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO- d_6) δ 11.33-12.51 (br.s, 1 H), 8.94 (s, 1 H), 7.79 (t, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.93 (t, 2 H), 3.04 (dd, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.21 (t, 2 H), 2.12 (t, 2 H), 1.63 (m, 6 H); MS (ESI+) m/z

35

674 (M+H)⁺.

Example 264

5 N-[5-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-N-
 methylglycine

 The product from Example 234B and N-methylglycine ethyl ester hydrochloride
 were processed as described in Example 251A and B to provide the titled compound. ¹H
10 NMR (500 MHz, DMSO-d₆) δ 12.20-13.15 (br.s, 1 H), 8.94 (s, 1 H), 7.23 (m, 7 H), 7.03 (t,
 1 H), 6.95 (m, 6 H), 6.82 (d, 2 H), 4.12 (s, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.94 (m, 3 H),
 3.01 (s, 2 H), 2.91 (s, 3 H), 2.81 (s, 1 H), 2.40 (m, 4 H), 2.27 (t, 1 H), 1.58-1.79 (m, 4 H);
 MS (ESI+) m/z 660 (M+H)⁺.

15

Example 265

N-[5-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-glutamic
 acid

20 The product from Example 234B and L-glutamic acid dimethyl ester hydrochloride
 were processed as described in Example 251A and B to provide the titled compound. ¹H
 NMR (500 MHz, DMSO-d₆) δ 11.60-13.04 (br.s, 2 H), 8.94 (s, 1 H), 8.06 (d, 1 H), 7.23 (m,
 7 H), 7.03 (t, 1 H), 6.95 (m, 6 H), 6.82 (d, 2 H), 4.21 (m, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H),
 3.93 (t, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.28 (m, 2 H), 2.19 (t, 2 H), 1.96 (m, 1 H), 1.60-
25 1.82 (m, 5 H); MS (ESI+) m/z 718 (M+H)⁺.

Example 266

30 N-[5-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-valine

 The product from Example 234B and L-valine methyl ester hydrochloride were
 processed as described in Example 251A and B to provide the titled compound. ¹H NMR
 (500 MHz, DMSO-d₆) δ 12.15-12.77 (br.s, 1 H), 8.94 (s, 1 H), 7.92 (d, 1 H), 7.23 (m, 7 H),
 7.03 (t, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.15 (dd, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.94 (t,
35 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.23 (m, 2 H), 2.04 (m, 1 H), 1.67 (m, 4 H), 0.88 (dd, 6 H);
 MS (ESI+) m/z 688 (M+H)⁺.

Example 267

1-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-proline

5 The product from Example 234B and L-proline methyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.95-13.15 (br.s, 1 H), 8.94 (s, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.95 (m, 6 H), 6.82 (d, 2 H), 4.21 (dd, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.93 (m, 2 H), 3.52 (m, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.34 (t, 2 H), 1.99-2.27 (m, 2 H), 1.87 (m, 2 H), 1.68 (m, 4 H; MS (ESI+) m/z 686 (M+H)⁺.

Example 268

5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-oxotetrahydrofuran-3-yl)pentanamide

15 The product from Example 234B and alpha-amino-gamma-butyrolactone hydrobromide were processed as described in Example 251A to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.94 (s, 1 H), 8.33 (d, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.53 (dd, 1 H), 4.34 (t, 1 H), 4.20 (m, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.94 (t, 2 H), 3.44 (m, 1 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.16 (m, 3 H), 1.55-1.78 (m, 4 H); MS (ESI+) m/z 672 (M+H)⁺.

Example 269

N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-serine

25 The product from Example 234B and O-tert-butyl L-serine tert-butyl ester hydrochloride were processed as described in Example 251A and 258B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.98-13.00 (br.s, 1 H), 8.95 (s, 1 H), 7.93 (d, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.27 (m, 1 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.94 (t, 2 H), 3.65 (m, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.22 (m, 2 H), 1.66 (m, 4 H); MS (ESI+) m/z 676 (M+H)⁺.

Example 270

N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-isoleucine

35 The product from Example 234B and L-isoleucine tert-butyl ester hydrochloride

were processed as described in Example 251A and 258B to provide the titled compound.

¹H NMR (500 MHz, DMSO-d₆) δ 12.09-12.71 (br.s, 1 H), 8.94 (s, 1 H), 7.93 (d, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.94 (m, 6 H), 6.82 (d, 2 H), 4.19 (dd, 1 H), 4.04 (s, 2 H), 4.00 (s, 2 H), 3.93 (t, 2 H), 2.91 (s, 3 H), 2.38 (s, 3 H), 2.21 (m, 2 H), 1.59-1.80 (m, 5 H), 1.39 (m, 1 H), 1.20 (m, 1 H), 0.84 (m, 6 H); MS (ESI+) m/z 702 (M+H)⁺.

Example 271

4-[[4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy]acetyl]amino}butanoic
acid

The product from Example 95C and ethyl 4-aminobutyrate hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.69-12.36 (br.s, 1 H), 8.94 (s, 1 H), 8.09 (t, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.96 (m, 6 H), 6.83 (d, 2 H), 4.43 (s, 2 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 3.15 (m, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.20 (t, 2 H), 1.66 (m, 2 H); MS (ESI+) m/z 632 (M+H)⁺.

Example 272

N-[[4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy]acetyl]-N-methylglycine

The product from Example 95C and N-methylglycine ethyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.40-13.15 (br.s, 1 H), 8.94 (s, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.93 (m, 6 H), 6.82 (dd, 2 H), 4.84 (s, 1 H), 4.69 (s, 1 H), 4.17 (s, 1 H), 4.05 (s, 2 H), 4.00 (m, 3 H), 3.04 (s, 2 H), 2.91 (s, 3 H), 2.85 (s, 1 H), 2.39 (s, 3 H); MS (ESI+) m/z 618 (M+H)⁺.

Example 273

N-[[4-[(benzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy]acetyl]-L-valine

The product from Example 95C and L-valine methyl ester hydrochloride were processed as described in Example 251A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.36-13.13 (br.s, 1 H), 8.94 (s, 1 H), 7.99 (d, 1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.95 (m, 6 H), 6.82 (d, 2 H), 4.58 (d, 2 H), 4.21 (dd, 1 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.10 (m, 1 H), 0.87 (dd, 6 H); MS (ESI+) m/z 646 (M+H)⁺.

Example 274

2-(4-{4-[(benzyl{2-methyl-3-
5 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-
oxotetrahydrofuran-3-yl)acetamide

The product from Example 95C and alpha-amino-gamma-butyrolactone hydrobromide were processed as described in Example 251A to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.94 (s, 1 H), 8.62 (d, 1 H), 7.23 (m, 7 H),
10 7.00 (m, 7 H), 6.84 (d, 2 H), 4.69 (dd, 1 H), 4.52 (d, 2 H), 4.36 (m, 1 H), 4.23 (dd, 1 H),
4.05 (s, 2 H), 4.00 (s, 2 H), 2.91 (s, 3 H), 2.39 (m, 4 H), 2.27 (m, 1 H); MS (ESI+) m/z 630 (M+H)⁺.

Example 275

15 N-[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-L-serine

The product from Example 95C and O-tert-butyl L-serine tert-butyl ester hydrochloride were processed as described in Example 251A and 258B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.26-13.13 (br.s, 1 H), 8.94 (s, 1 H), 8.01
20 (d, 1 H), 7.23 (m, 7 H), 7.00 (m, 7 H), 6.84 (d, 2 H), 4.54 (s, 2 H), 4.36 (m, 1 H), 4.05 (s, 2
H), 4.00 (s, 2 H), 3.78 (m, 1 H), 3.68 (m, 1 H), 2.91 (s, 3 H), 2.39 (s, 3 H); MS (ESI+) m/z
634 (M+H)⁺.

Example 276

25 N-2~[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-L-asparagine

The product from Example 95C and L-asparagine tert-butyl ester hydrochloride were processed as described in Example 251A and 258B to provide the titled compound. ¹H
30 NMR (500 MHz, DMSO-d₆) δ 12.18-13.09 (br.s, 1 H), 8.94 (s, 1 H), 8.28 (d, 1 H), 7.39 (s,
1 H), 7.23 (m, 7 H), 7.03 (t, 1 H), 6.94 (m, 6 H), 6.91 (s, 1 H), 6.84 (d, 2 H), 4.60 (dd, 1 H),
4.49 (s, 2 H), 4.05 (s, 2 H), 4.00 (s, 2 H), 2.91 (s, 3 H), 2.60 (d2 H), 2.39 (s, 3 H); MS
(ESI+) m/z 661 (M+H)⁺.

35

Example 277

N-[3-(benzyl{4-[4-(2-hydrazino-2-oxoethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

The product from Example 95C and tert-butyl carbazate were processed as described in Example 251A and 258B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.94 (s, 1 H), 7.22 (m, 7 H), 6.99 (m, 7 H), 6.83 (d, 2 H), 4.64 (m, 2 H), 4.05 (s, 2 H), 4.01 (s, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H); MS (ESI+) m/z 561 (M+H)⁺.

Example 278

N-(3-{benzyl[4-(3-{[(2S)-5-oxopyrrolidin-2-yl]methoxy}phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

Example-278A

(5S)-5-((3-(4-((benzyl(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenoxy)methyl)pyrrolidin-2-one

The product from Example 61F (88mg, 0.2mmole), (S)-5-(hydroxymethyl)-2-pyrrolidinone (58mg, 0.5mmole), di-tert-butyl azodicarboxylate (69mg, 0.3mmole), and triphenylphosphine (77mg, 0.3mmole) were dissolved in tetrahydrofuran (1.5mL) in a capped test-tube and shaken at room temperature overnight. Ethyl acetate (2mL) was added to dilute the reaction mixture. Then, the diluted organic solution was washed with water (2mL) and saturated sodium chloride solution (2mL), dried *in vacuo*, and purified by HPLC to give the pure intermediate.

Example-278B

(5S)-5-((3-(4-(((3-amino-2-methylphenyl)(benzyl)amino)methyl)phenoxy)phenoxy)methyl)pyrrolidin-2-one

To the product from Example 278A in a capped test-tube were added iron powder (78.4mg, 1.4mmoles), NH₄Cl (7.5mg, 1.4mmoles), ethyl alcohol (3mL) and water (1mL). The reaction mixture was shaken violently at 80°C overnight. CH₂Cl₂ (4mL) was added to extract the product. Then, the reaction mixture was filtered. The CH₂Cl₂ layer was separated and dried *in vacuo*.

Example-278C

N-(3-{benzyl[4-(3-{[(2S)-5-oxopyrrolidin-2-yl]methoxy}phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The dried product from example 278B was dissolved in anhydrous pyridine (3mL) at 0°C for 30 minutes. Then methanesulfonyl chloride (50μL, 6mmoles) was injected. The

reaction mixture was let to stand at room temperature for 3 hours. Then the reaction mixture was dried *in vacuo*, dissolved in CH₂Cl₂ (4mL), filtered and dried *in vacuo*. The final residue was purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.76 (s, 1 H), 7.27 (m, 7 H), 7.20 (m, 1 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.71 (m, 1 H), 6.53 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.92 (m, 1 H), 3.82 (m, 2 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.04-2.29 (m, 3 H), 1.80 (m, 1 H); MS (APCI+) m/z 586 (M+H)⁺.

Example 279

1-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-3-carboxylic acid

Example 279A

ethyl 1-(4-(3-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)piperidine-3-carboxylate

The product from Example 116 (57mg, 0.10mmole), 1-[3-(dimethylamino)propyl]-1-ethylcarbodiimide hydrochloride (27mg, 0.14mmole), and 1-hydroxybenzotriazole hydrate (19mg, 0.14mmole) were dissolved in N,N-dimethylformamide (1.5mL) and let stand for 15 minutes at room temperature. Ethyl nipecotate (31mg, 0.20mmole) and N,N-diisopropylethylamine (35μL, 0.20mmole) in N,N-dimethylformamide (0.5mL) were added. The reaction mixture was shaken at room temperature overnight. Then the reaction mixture was acidified with 1N hydrochloric acid (3mL) and extracted with dichloromethane (3mL). The organic layer was washed with saturated NaHCO₃ solution (3mL) and brine (3mL), dried (MgSO₄), and concentrated *in vacuo*. The crude product was purified using HPLC to provide the ester compound.

Example 279B

1-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-3-carboxylic acid

The product from Example 279A was treated with 2N NaOH aqueous solution (1mL), ethyl alcohol (0.5mL) and tetrahydrofuran (0.5mL) at room temperature overnight. 1N hydrochloric acid was dropped to the reaction mixture until pH=2~3. Ethyl acetate

(3mL) was used to extract the product. The organic layer was separated, dried (Na₂SO₄), concentrated *in vacuo*, and purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.03-12.70 (br.s, 1 H), 8.95 (s, 1 H), 7.24 (m, 8 H), 7.04 (m, 1 H), 6.94 (m, 4 H), 6.69 (m, 1 H), 6.49 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (t, *J* = 6.4 Hz, 2 H), 3.82 (m, 1 H), 3.72 (m, 1 H), 3.31 (dd, *J* = 13.6, 9.2 Hz, 1 H), 2.98 (t, *J* = 11.1 Hz, 1 H), 2.91 (s, 3 H), 2.38 (m, 5 H), 2.24 (m, 1 H), 1.90 (m, 3 H), 1.65 (m, 1 H), 1.52 (m, 1 H), 1.34 (m, 1 H); MS (APCI+) *m/z* 686.1 (M+H)⁺.

10

Example 280

1-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-4-carboxylic acid

The product from Example 116B (57mg, 0.10mmole) and ethyl isonipecotate (31mg, 0.20mmole) were processed as in Example 279A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.05-12.35 (br.s, 1 H), 8.95 (s, 1 H), 7.23 (m, 8 H), 7.03 (m, 1 H), 6.94 (m, 4 H), 6.69 (dd, 1 H), 6.49 (m, 2 H), 4.20 (d, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (t, 2 H), 3.77 (d, 1 H), 3.05 (t, 1 H), 2.91 (s, 3 H), 2.70 (t, 1 H), 2.47 (m, 1 H), 2.43 (m, 2 H), 2.39 (s, 3 H), 1.90 (m, 2 H), 1.79 (m, 2 H), 1.45 (m, 1 H), 1.34 (m, 1 H); MS (APCI+) *m/z* 686.4 (M+H)⁺.

20

Example 281

1-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-2-carboxylic acid

The product from Example 116B (57mg, 0.10mmole) and ethyl pipecolate (31mg, 0.20mmole) were processed as in Example 279A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.29-13.30 (br.s, 1 H), 8.95 (s, 1 H), 7.24 (m, 8 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.69 (m, 1 H), 6.49 (m, 2 H), 5.07 (d, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (m, 2 H), 3.77 (d, 1 H), 3.09 (m, 1 H), 2.91 (s, 3 H), 2.49 (m, 2 H), 2.40 (s, 3 H), 2.12 (m, 1 H), 1.90 (m, 2 H), 1.62 (m, 2 H), 1.49 (m, 1 H), 1.35 (m, 1 H), 1.21 (m, 1 H); MS (APCI+) *m/z* 685.9 (M+H)⁺.

30

Example 282

1-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]proline

The product from Example 116B (57mg, 0.10mmole) and H-PRO-OME hydrochloride (33mg, 0.20mmole) were processed as in Example 279A and B to provide the
 5 titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.66-13.20 (br.s, 1 H), 8.95 (s, 1 H), 7.24 (m, 8 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.69 (m, 1 H), 6.50 (m, 2 H), 4.46 (m, 0 H), 4.22 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.93 (m, 2 H), 3.50 (m, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.39 (m, 2 H), 1.98-2.25 (m, 2 H), 1.76-1.97 (m, 4 H); MS (APCI+) m/z 672.5 (M+H)⁺.

10

Example 283

N-{3-[benzyl(4-{3-[(3-methyloxetan-3-yl)methoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 61F (88mg, 0.2mmole) and 3-methyl-3-oxetanemethanol
 15 (51mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.23 (m, 8 H), 7.04 (m, 1 H), 6.94 (m, 4 H), 6.72 (m, 1 H), 6.53 (m, 2 H), 4.45 (d, 1 H), 4.27 (d, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 4.01 (s, 1 H), 3.78 (dd, 1 H), 3.64 (dd, 1 H), 3.38 (d, 1 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 1.33 (s, 2
 20 H), 0.99 (s, 1 H); MS (APCI+) m/z 573 (M+H)⁺.

Example 284

2-[(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)methyl]cyclopropanecarb
 25 oxylic acid

25

Example 284A

ethyl 2-((3-(4-((benzyl(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)methyl)cyclopropanecarb
 30 oxylate

30

The product from Example 61F (88mg, 0.2mmole) and ethyl 2-hydroxymethyl-1-cyclopropanecarboxylate (51mg, 0.5mmole) were processed as in Example 278A-C to provide the ester compound.

35

Example 284B

2-[(3-{4-[(benzyl{2-methyl-3-

[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)methyl]cyclopropanecarb
oxylic acid

To the product from Example 284A was added 2N hydrochloric acid (1mL), tetrahydrofuran (0.5mL) and water (0.5mL). The reaction mixture was stirred at room temperature overnight. Water (2mL) was added. The solution was neutralized with 1N hydrochloric acid until pH=2~3. The product was extracted with ethyl acetate (2mL), dried *in vacuo*, and purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.01-12.29 (br.s, 1 H), 8.95 (s, 1 H), 7.23 (m, 8 H), 7.03 (m, 1 H), 6.94 (m, 4 H), 6.69 (m, 1 H), 6.50 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (m, 1 H), 3.80 (m, 1 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 1.67 (m, 1 H), 1.57 (m, 1 H), 1.05 (m, 1 H), 0.92 (m, 1 H); MS (APCI+) m/z 587.1 (M+H)⁺

Example 285

N-(3-{benzyl[4-(3-{[(2R)-2,3-dihydroxypropyl]oxy}phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

Example 285A

N-(3-(benzyl(4-(3-(((4S)-2,2-dimethyl-1,3-dioxolan-4-yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide

The product from Example 61F (88mg, 0.2mmole) and (S)-(+)-2,2-dimethyl-1,3-dioxolane-4-methanol (66mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound.

Example 285B

N-(3-{benzyl[4-(3-{[(2R)-2,3-dihydroxypropyl]oxy}phenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide

The product from Example 285A was treated with tetrafluoroacetic acid (0.20mL) and dichloromethane (1.8mL). The reaction mixture was stirred for 3 hours at room temperature, then dried *in vacuo*, and purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.23 (m, 8 H), 7.04 (m, 1 H), 6.94 (m, 4 H), 6.69 (m, 1 H), 6.49 (m, 2 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.95 (m, 1 H), 3.81 (m, 1 H), 3.75 (m, 1 H), 3.41 (m, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H); MS (APCI+) m/z 563.2 (M+H)⁺.

Example 286

4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)cyclohexanecarboxylic
acid

The product from Example 61F (88mg, 0.2mmole) and ethyl 4-
5 hydroxycyclohexanecarboxylate (86mg, 0.5mmole) were processed as in Example 278 A-C
and 284B to provide the title compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.88-12.20 (m,
1 H), 8.95 (s, 1 H), 7.23 (m, 8 H), 7.03 (m, 1 H), 6.94 (m, 4 H), 6.69 (m, 1 H), 6.52 (m, 1
H), 6.46 (m, 1 H), 4.20-4.52 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 2.91 (s, 3 H), 2.40 (s, 3
H), 2.17-2.38 (m, 1 H), 1.85-2.06 (m, 3 H), 1.57-1.83 (m, 2 H), 1.29-1.55 (m, 3 H); MS
10 (APCI+) m/z 615.1 (M+H)⁺.

Example 287

N-[3-(benzyl{4-[3-(tetrahydro-2H-pyran-4-yloxy)phenoxy]benzyl}amino)-2-
15 methylphenyl]methanesulfonamide

The product from Example 61F (88mg, 0.2mmole) and tetrahydro-2H-pyran-4-ol
(51mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound.
¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.24 (m, 8 H), 7.04 (m, 1 H), 6.95 (m, 4 H),
6.73 (dd, 1 H), 6.56 (m, 1 H), 6.48 (dd, 1 H), 4.53 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.81
20 (m, 2 H), 3.45 (m, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 1.92 (m, 2 H), 1.55 (m, 2 H); MS
(APCI+) m/z 573.7 (M+H)⁺.

Example 288

N-[3-(benzyl{4-[3-(tetrahydrofuran-3-yloxy)phenoxy]benzyl}amino)-2-
25 methylphenyl]methanesulfonamide

The product from Example 61F (88mg, 0.2mmole) and 3-hydroxytetrahydrofuran
(44mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound.
¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.24 (m, 8 H), 6.99 (m, 5 H), 6.67 (m, 1 H),
30 6.50 (m, 2 H), 4.97 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.79 (m, 4 H), 2.91 (s, 3 H), 2.39
(s, 3 H), 2.17 (m, 1 H), 1.93 (m, 1 H); MS (APCI+) m/z 559.6 (M+H)⁺.

Example 289

N-(3-{benzyl[4-(3-{[(2S)-1-methylpyrrolidin-2-yl]methoxy}phenoxy)benzyl]amino}-2-
35 methylphenyl)methanesulfonamide

The product from Example 61F (88mg, 0.2mmole) and (S)-(-)-2-hydroxymethyl-1-

5 methylpyrrolidine (58mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.25 (m, 8 H), 7.04 (t, 1 H), 6.95 (m, 4 H), 6.79 (d, 1 H), 6.64 (dd, 1 H), 6.58 (dd, 1 H), 4.31 (dd, 1 H), 4.16 (dd, 2 H), 4.06 (s, 2 H), 4.04 (s, 2 H), 2.93 (d, 2 H), 2.92 (s, 3 H), 2.77 (d, 2 H), 2.50 (s, 3 H), 2.40 (s, 3 H), 1.86 (m, 2 H); MS (APCI+) m/z 586 (M+H)⁺.

Example 290

N-[3-(benzyl{4-[3-(pyridin-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide

10 The product from Example 61F (88mg, 0.2mmole) and pyridine-3-methanol (55mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 8.81 (s, 1 H), 8.71 (d, 1 H), 8.20 (d, 1 H), 7.73 (dd, 1 H), 7.28 (m, 7 H), 7.20 (m, 1 H), 7.03 (m, 1 H), 6.94 (m, 4 H), 6.81 (dd, 1 H), 6.66 (t, 1 H),
15 6.55 (dd, 1 H), 5.21 (s, 2 H), 4.07 (s, 2 H), 4.04 (s, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H); MS (APCI+) m/z 580 (M+H)⁺.

Example 291

N-{3-[benzyl(4-{3-[(2S)-pyrrolidin-2-ylmethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

20 The product from Example 61F (88mg, 0.2mmole) and 1-methyl-3-pyrrolidinol (51mg, 0.5mmole) were processed as in Example 278A-C and 285B to provide the title compound processed as in Example 278A-C and 285B to provide the title compound. ¹H
25 NMR (500 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.28 (m, 7 H), 7.20 (m, 1 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.75 (m, 1 H), 6.58 (m, 2 H), 4.23 (m, 1 H), 4.06 (s, 2 H), 4.04 (s, 2 H), 4.03 (m, 1 H), 3.89 (m, 1 H), 3.20 (m, 2 H), 2.92 (s, 3 H), 2.40 (s, 3 H), 2.10 (m, 1 H), 1.92 (m, 2 H), 1.70 (m, 1 H); MS (APCI+) m/z 572 (M+H)⁺.

Example 292

N-{3-[benzyl(4-{3-[(1-methylpyrrolidin-3-yl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide

30 The product from Example 61F (88mg, 0.2mmole) and 1-methyl-3-pyrrolidinol (51mg, 0.5mmole) were processed as in Example 278A-C to provide the titled compound.
35 ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.29 (m, 7 H), 7.20 (m, 1 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.72 (m, 1 H), 6.56 (m, 2 H), 5.12 (m, 1 H), 4.06 (s, 2 H), 4.04 (s, 2 H), 3.33

(m, 1 H), 3.23 (m, 1 H), 3.11 (m, 1 H), 2.92 (s, 3 H), 2.85 (m, 3 H), 2.57 (m, 1 H), 2.40 (s, 3 H), 2.24 (m, 1 H), 2.03 (m, 1 H); MS (APCI+) m/z 572.6 (M+H)⁺.

5

Example 293N-{3-[benzyl(4-chloro-2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide

The product from Example 6A and 2-fluoro-4-chloro-1-(bromomethyl) benzene were processed as described in Example 6B-D to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 7.25 (m, 8 H), 7.04 (t, 1 H), 6.99 (d, 1 H), 6.94 (d, 1 H), 4.08 (s, 4 H), 2.89 (s, 3 H), 2.34 (s, 3 H); MS (APCI+) m/z 433 (M+H)⁺.

10

Example 294

15 N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamic acid

The product from Example 116B (57mg, 0.10mmole) and H-GLU(OME)-OME hydrochloride (42mg, 0.20mmole) were processed as in Example 279 A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.32-12.71 (br.s, 1 H), 11.89-12.32 (br.s, 1 H), 8.95 (s, 1 H), 8.10 (d, 1 H), 7.24 (m, 8 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.68 (dd, 1 H), 6.49 (m, 2 H), 4.20 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.93 (t, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.26 (m, 4 H), 1.93 (m, 3 H), 1.75 (m, 1 H); MS (APCI+) m/z 703.9 (M+H)⁺.

20

25

Example 295

30 4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-oxotetrahydrofuran-3-yl)butanamide

The product from Example 116B (57mg, 0.10mmole) and α-Amino-γ-butyrolactone hydrobromide (36mg, 0.20mmole) were processed as in Example 279A to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 8.36 (d, 1 H), 7.23 (m, 8 H), 7.04 (m, 1 H), 6.94 (m, 4 H), 6.69 (dd, 1 H), 6.50 (m, 2 H), 4.53 (d, 1 H), 4.33 (d, 1 H), 4.19 (ddd, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (t, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.35 (m, 1 H), 2.27 (m, 2 H), 2.13 (m, 1 H), 1.91 (m, 2 H); MS (APCI+) m/z 702.9 (M+H)⁺.

35

Example 296

N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-leucine

The product from Example 116B (57mg, 0.10mmole) and L-leucine methyl ester hydrochloride (36mg, 0.20mmole) were processed as in Example 279A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.22-12.61 (br.s, 1 H), 8.95 (s, 1 H), 8.06 (d, 1 H), 7.24 (m, 8 H), 7.03 (m, 1 H), 6.94 (m, 4 H), 6.67 (dd, 1 H), 6.49 (m, 2 H), 4.22 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.91 (m, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.26 (m, 2 H), 1.90 (m, 2 H), 1.60 (m, 1 H), 1.47 (m, 2 H), 0.85 (d, 3 H), 0.81 (d, 3 H); MS (APCI+) m/z 688.7 (M+H)⁺.

Example 297

N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-aspartic acid

The product from Example 116B (57mg, 0.10mmole) and H-ASP(OME)-OME hydrochloride (36mg, 0.20mmole) were processed as in Example 279A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.68-13.28 (br.s, 2 H), 8.95 (s, 1 H), 8.18 (d, 1 H), 7.24 (m, 8 H), 7.04 (m, 1 H), 6.95 (m, 4 H), 6.68 (dd, 1 H), 6.49 (m, 2 H), 4.53 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.92 (m, 2 H), 2.91 (s, 3 H), 2.68 (m, 1 H), 2.55 (m, 1 H), 2.39 (s, 3 H), 2.24 (m, 2 H), 1.89 (m, 2 H); MS (APCI+) m/z 690.7 (M+H)⁺.

Example 298

N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-valine

The product from Example 116B (57mg, 0.10mmole) and L-valine methyl ester hydrochloride (34mg, 0.20mmole) were processed as in Example 279A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.03-12.85 (br.s, 1 H), 8.95 (s, 1 H), 7.96 (d, 1 H), 7.23 (m, 8 H), 7.04 (m, 1 H), 6.94 (m, 4 H), 6.68 (dd, 1 H), 6.49 (m, 2 H), 4.14 (dd, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.92 (t, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.31 (m, 2 H), 2.02 (m, 1 H), 1.90 (m, 2 H), 0.86 (s, 3 H), 0.85 (s, 3 H); MS (APCI+) m/z 674.7 (M+H)⁺.

Example 299

1-[(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]piperidine-4-
carboxamide

The product from Example 91 (62 mg, 0.1 mmoles) in anhydrous
5 dimethylformamide (1.0 ml) was treated with 1-[3-(dimethylamino)propyl]-3-
ethylcarbodiimide hydrochloride (27 mg, 0.14 mmoles) and 1-hydroxybenzotriazole hydrate
(19 mg, 0.14 mmoles) at room temperature for 15 minutes. A mixture of isonipecotamide
(24.4 mg, 0.21 mmole) and N,N-diisopropylethylamine (0.056 ml, 0.4 mmoles) in anhydrous
dimethylformamide (0.2 ml) was added to the reaction and shaken at room temperature
10 overnight. The reaction mixture was acidified with 1N aqueous hydrochloric acid and
extracted with dichloromethane. The organic layer was washed with saturated aqueous
sodium bicarbonate solution and brine, dried over anhydrous sodium sulfate, filtered and
filtrate concentrated in vacuo. The residue purified by HPLC to provide the titled
compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.94 (s, 1 H), 7.37 (d, 1 H), 7.26 (m, 4 H),
15 7.09 (m, 2 H), 6.96 (m, 5 H), 6.78 (d, 1 H), 6.75 (s, 1 H), 6.45 (dd, 1 H), 4.92 (m, 2 H), 4.23
(d, 1 H), 4.07 (d, 4 H), 3.79 (d, 1 H), 3.00 (t, 1 H), 2.90 (s, 3 H), 2.64 (m, 1 H), 2.32 (m, 4
H), 1.71 (m, 2 H), 1.51 (m, 1 H), 1.33 (m, 1 H); MS (ESI+) m/z 727 (M+H)⁺.

20

Example 300

2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-[3-(2-oxopyrrolidin-
1-yl)propyl]acetamide

The product from Example 91 and 1-(3-aminopropyl)-2-pyrrolidinone were
25 processed as described in Example 299 to provide the titled compound. ¹H NMR (500
MHz, DMSO-d₆) δ 8.94 (s, 1 H), 7.92 (t, 1 H), 7.40 (d, 1 H), 7.26 (m, 3 H), 7.09 (m, 2 H),
6.96 (m, 5 H), 6.74 (d, 1 H), 6.51 (dd, 1 H), 4.56 (s, 2 H), 4.07 (d, 4 H), 3.27 (m, 2 H), 3.13
(t, 2 H), 3.07 (dd, 2 H), 2.90 (s, 3 H), 2.33 (s, 3 H), 2.18 (t, 2 H), 1.88 (m, 2 H), 1.56 (m, 2
H); MS (ESI+) m/z 741 (M+H)⁺.

30

Example 301

2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(1,3-thiazol-5-
ylmethyl)acetamide

35

The product from Example 91 and 5-(aminomethyl)thiazole were processed as
described in Example 299 to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆)

δ 9.02 (s, 1 H), 8.95 (s, 1 H), 8.47 (t, 1 H), 7.39 (m, 2 H), 7.27 (m, 3 H), 7.09 (m, 2 H), 6.96 (m, 5 H), 6.79 (d, 1 H), 6.50 (dd, 1 H), 4.66 (s, 2 H), 4.46 (d, 2 H), 4.07 (d, 4 H), 2.90 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 713 (M+H)⁺.

5

Example 302

2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(3-morpholin-4-ylpropyl)acetamide

10 The product from Example 91 and N-(3-aminopropyl)morpholine were processed as described in Example 299 to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 8.95 (s, 1 H), 8.08 (t, 1 H), 7.41 (d, 1 H), 7.26 (m, 3 H), 7.10 (m, 2 H), 6.96 (m, 5 H), 6.79 (d, 1 H), 6.50 (dd, 1 H), 4.59 (s, 2 H), 4.07 (d, 4 H), 3.96 (d, 2 H), 3.61 (m, 2 H), 3.39 (d, 2 H), 3.20 (dd, 2 H), 3.05 (m, 4 H), 2.90 (s, 3 H), 2.33 (s, 3 H), 1.82 (m, 2 H); MS (ESI+) m/z 743 (M+H)⁺.

15

Example 303

N-[4-(3-{4-[(benzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-serine

20

Example 303A

The product from Example 116 (57mg, 0.10mmole) and O-tert-butyl-L-serine tert-butyl ester hydrochloride (60mg, 0.20mmole) were processed as in Example 279A to provide the titled compound.

25

Example 303B

The product from example 303A was treated with trifluoroacetic acid (0.2mL) and dichloromethane (1.8mL) at room temperature for about 4 hours with shaking. Saturated NaHCO₃ solution (2mL) was added to neutralize the reaction mixture 1N hydrochloric acid was used to adjust the pH value to 2~3. The organic layer was separated, dried (MgSO₄), concentrated *in vacuo*, and purified by HPLC to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 11.94-12.89 (br.s, 1 H), 8.95 (s, 1 H), 7.98 (d, 1 H), 7.24 (m, 8 H), 7.03 (m, 1 H), 6.94 (m, 4 H), 6.68 (dd, 1 H), 6.49 (m, 2 H), 4.27 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.94 (t, 2 H), 3.67 (dd, 1 H), 3.61 (dd, 1 H), 2.91 (s, 3 H), 2.39 (s, 3 H), 2.30 (t, 2 H), 1.90 (m, 2 H); MS (APCI+) m/z 662.1 (M+H)⁺.

30

35

Example 304

N~2~[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamine

5 The product from Example 116 (57mg, 0.10mmole) and L-glutamine tert-butyl ester hydrochloride (26mg, 0.20mmole) were processed as in Example 303A and B to provide the titled compound. ¹H NMR (500 MHz, DMSO-d₆) δ 12.07-12.91 (br.s, 1 H), 8.95 (s, 1 H), 8.11 (d, 1 H), 7.23 (m, 9 H), 7.05 (m, 1 H), 6.95 (m, 4 H), 6.71 (m, 2 H), 6.49 (m, 2 H), 4.15 (m, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.93 (t, 2 H), 2.91 (s, 3 H), 2.40 (s, 3 H), 2.27 (t, 2 H),
10 2.11 (t, 2 H), 1.92 (m, 1 H), 1.90 (t, 2 H), 1.73 (m, 1 H); MS (APCI+) m/z 703.7 (M+H)⁺.

Example 305

(5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
15 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-fluorophenoxy)acetic acid

Example 305A4-fluorobenzene-1,3-diol

20 [1-(Chloromethyl)-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate)] (SELECTFLUOR) (1.61 g., 4.54 mmoles) in anhydrous acetonitrile (50 mL) was treated with resorcinol (0.500 g., 4.54 mmoles) and heated at 100°C overnight. The reaction was diluted with diethyl ether, washed with H₂O (2X), saturated NaHCO₃ (2X), brine, dried (Na₂SO₄), filtered, and the filtrate concentrated under reduced pressure. MS (DCI) m/z 129
25 (M+H)⁺.

Example 305B3-(allyloxy)-4-fluorophenyl 4-methylbenzenesulfonate

30 The product from Example 305A was processed as described in Example 63A. MS (DCI) m/z 300 (M+NH₄)⁺.

Example 305C3-(allyloxy)-4-fluorophenol

35 The product from Example 305B was processed as described in Example 63B. MS

(DCI) m/z 168 (M+H)⁺.

Example 305D

5 4-(3-(allyloxy)-4-fluorophenoxy)benzaldehyde

The product from Example 305C and 4-fluorobenzaldehyde were processed as in Example 61C to provide the title compound.

10 Example 305E

N-(4-(3-(allyloxy)-4-fluorophenoxy)benzyl)-N-(2-methyl-3-nitrophenyl)amine

The product from Example 305D and 2-methyl-3-nitroaniline were processed as in Example 6A to provide the title compound. MS (ESI-) m/z 407 (M-H)⁻.

15 Example 305F

N-(4-(3-(allyloxy)-4-fluorophenoxy)benzyl)-N-(2,4-difluorobenzyl)-N-(2-methyl-3-nitrophenyl)amine

20 The product from Example 305E and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound. MS (ESI+) m/z 535 (M+H)⁺.

Example 305G

25 5-(4-(((2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)-2-fluorophenol

The product from Example 305F was processed as described in Example 33A to provide the title compound.

Example 305H

30 ethyl (5-(4-(((2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)-2-fluorophenoxy)acetate

The product from Example 305G and ethyl glycolate were processed as described in Example 62A to provide the title compound.

35 Example 305I

ethyl (5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-fluorophenoxy)acetate

The product from Example 305H was processed as described in Example 6C and D to provide the title compound.

5

Example 305J

(5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-fluorophenoxy)acetic acid

The product from Example 305I was processed as described in Example 50 to provide the title compound. ¹H NMR (500 MHz, DMSO) δ 8.95 (s, 1 H), 7.26 (d, 2 H), 7.21 (m, 2 H), 7.11 (m, 1 H), 7.05 (d, 1 H), 7.00 (d, 1 H), 6.95 (m, 2 H), 6.88 (d, 2 H), 6.82 (m, 1 H), 6.48 (m, 1 H), 4.75 (s, 2 H), 4.07 (s, 2 H), 4.05 (s, 2 H), 2.90 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 601 (M+H)⁺.

15

Example 306

(2-chloro-5-(4-(((2-fluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

20

Example 306A

ethyl (2-chloro-5-(4-(((2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenoxy)acetate

The product from Example 91D was processed as in Examples 33A and 62A to provide the title compound.

25

Example 306B

ethyl (2-chloro-5-(4-(((2-fluorobenzyl)(2-methyl-3-
nitrophenyl)amino)methyl)phenoxy)phenoxy)acetate

The product from Example 306A (86 mg, 0.18 mmoles) in anhydrous dimethylformamide (1 mL) was treated with Hunig's base (0.107 mL, 0.5 mmoles) and 1-(bromomethyl)-2-fluorobenzene (0.198 mL, 0.9 mmoles) in anhydrous dimethylformamide (1 mL) and heated at 90°C for 72 hours. The reaction was diluted with ether, washed with saturated NH₄Cl, water, brine, dried (Na₂SO₄), filtered, and filtrate concentrated under reduced pressure. The residue purified by using ethyl acetate and hexane gradient on 10 g silica column to provide the title compound.

35

Example 306C

ethyl (5-(4-(((3-amino-2-methylphenyl)(2-fluorobenzyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetate

5 The product from Example 306B in ethanol (1.2 mL) and H₂O (0.6 mL) was treated with iron powder (100 mg, 1.8 mmoles) and NH₄Cl (10 mg, 0.18 mmoles) at 80°C overnight. The reaction mixture filtered, residue washed with methanol and dichloromethane, the filtrate concentrated in vacuo to provide the title compound.

Example 306D

10 ethyl (2-chloro-5-(4-(((2-fluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetate compound with ethane (1:1)

15 The product from Example 306C in anhydrous pyridine (1.5 mL) was cooled at 0°C and treated with methane sulfonyl chloride (0.042 mL, 0.54 mmoles) at 0°C for 1 hour, then room temperature for 1 hour. The reaction mixture concentrated in vacuo, residue dissolved in dichloromethane, filtered and filtrate concentrated in vacuo to provide the title compound.

Example 306E

20 The product from Example 306D was treated with 2M aqueous NaOH (2 mL), tetrahydrofuran (1 mL) and ethanol (1 mL) at room temperature overnight. The reaction was acidified with 1 M aqueous hydrochloric acid to pH 3-4, extracted with ethyl acetate, dried (Na₂SO₄), filtered and filtrate concentrated in vacuo. The residue was purified by HPLC to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.76-13.43 (br.s, 1 H), 8.94 (s, 1 H), 7.38 (d, 1 H), 7.26 (m, 4 H), 7.07 (m, 3 H), 6.95 (m, 4 H), 6.77 (d, 1 H), 6.47 (dd, 1 H), 4.78 (s, 2 H), 4.11 (s, 2 H), 4.07 (s, 2 H), 2.90 (s, 3 H), 2.35 (s, 3 H); MS (ESI+) m/z 599 (M+H).

Example 307

30 (5-(4-(((2-bromobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid

35 The product from Example 306A and 1-bromo-2-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 8.93 (s, 1 H), 7.55 (d, 1 H), 7.42 (d, 1 H), 7.38 (d, 1 H), 7.28 (m, 3 H), 7.15 (m, 1 H), 7.15 (br.s, 1 H), 7.07 (t, 1 H), 7.00 (t, 2 H), 6.93 (d, 2 H), 6.76 (d, 1 H),

6.46 (dd, 1 H), 4.78 (s, 2 H), 4.17 (s, 2 H), 4.11 (s, 2 H), 2.89 (s, 3 H), 2.32 (s, 3 H); MS (ESI+) m/z 681 (M+22).

5

Example 308

(5-(4-(((4-bromobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid

The product from Example 306A and 1-bromo-4-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.84-13.29 (br.s, 1 H), 8.95 (s, 1 H), 7.45 (d, 2 H), 7.38 (d, 1 H), 7.26 (d, J = 8.7 Hz, 2 H), 7.20 (d, 2 H), 7.04 (t, 1 H), 6.95 (m, 4 H), 6.76 (d, 1 H), 6.48 (dd, 1 H), 4.78 (s, 2 H), 4.03 (d, 4 H), 2.92 (s, 3 H), 2.38 (s, 3 H); MS (ESI+) m/z 661 (M+H).

15

Example 309

(2-chloro-5-(4-(((2,4-dichlorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

The product from Example 306A and 2,4-dichloro-1-(iodomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.78-13.45 (br.s, 1 H), 8.94 (s, 1 H), 7.52 (d, 1 H), 7.38 (d, 2 H), 7.32 (dd, 1 H), 7.26 (d, 2 H), 7.07 (t, 1 H), 7.00 (d, 2 H), 6.93 (d, 2 H), 6.76 (d, 1 H), 6.46 (dd, 1 H), 4.78 (s, 2 H), 4.16 (s, 2 H), 4.09 (s, 2 H), 2.90 (s, 3 H), 2.30 (s, 3 H); MS (ESI+) m/z 651 (M+H).

25

Example 310

(2-chloro-5-(4-(((4-chloro-2-fluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

The product from Example 306A and 2-fluoro-4-chloro-1-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.86-13.21 (br.s, 1 H), 8.95 (s, 1 H), 7.38 (d, 1 H), 7.28 (m, 4 H), 7.17 (d, 1 H), 7.06 (t, 1 H), 6.96 (m, 4 H), 6.77 (d, 1 H), 6.47 (dd, 1 H), 4.78 (s, 2 H), 4.07 (d, 4 H), 2.90 (s, 3 H), 2.33 (s, 3 H); MS (ESI+) m/z 633 (M+H).

35

Example 311

(2-chloro-5-(4-(((2-chloro-4-fluorobenzyl)(2-methyl-3-

((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

The product from Example 306A and 2-chloro-4-fluoro-1-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.86-13.32 (br.s, 1 H), 8.93 (s, 1 H), 7.39 (m, 2 H), 7.34 (dd, 1 H), 7.26 (d, 2 H), 7.11 (t, 1 H), 7.07 (d, 1 H), 7.01 (t, 2 H), 6.93 (d, 2 H), 6.76 (d, 1 H), 6.46 (dd, 1 H), 4.78 (s, 2 H), 4.15 (s, 2 H), 4.09 (s, 2 H), 2.89 (s, 3 H), 2.30 (s, 3 H); MS (ESI+) m/z 633 (M+H).

Example 312

(5-(4-(((2-bromo-4-chlorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid

The product from Example 306A and 2-bromo-4-chloro-1-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.84-13.32 (br.s, 1 H), 8.96 (s, 1 H), 7.67 (s, 1 H), 7.38 (m, 3 H), 7.24 (d, 2 H), 7.07 (d, 1 H), 7.00 (d, 2 H), 6.92 (d, 2 H), 6.74 (s, 1 H), 6.46 (dd, 1 H), 4.75 (s, 2 H), 4.15 (s, 2 H), 4.10 (s, 2 H), 2.90 (s, 3 H), 2.30 (s, 3 H); MS (ESI+) m/z 695 (M+H).

Example 313

(2-chloro-5-(4-(((3,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

The product from Example 306A and 3,4-difluoro-1-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.91-13.25 (br.s, 1 H), 8.96 (s, 1 H), 7.38 (d, 1 H), 7.27 (m, 4 H), 7.08 (m, 2 H), 6.95 (m, 4 H), 6.76 (d, 1 H), 6.47 (dd, 1 H), 4.78 (s, 2 H), 4.05 (d, 4 H), 2.92 (s, 3 H), 2.38 (s, 3 H); MS (ESI-) m/z 695 (M+79).

Example 314

(5-(4-(((4-benzoylbenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid

The product from Example 306A and 4-(bromomethyl)benzophenone were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.76-13.43 (br.s, 1 H), 8.97 (s, 1 H), 7.67 (m, 5 H), 7.55 (t, 2 H),

7.46 (d, 2 H), 7.38 (d, 1 H), 7.29 (d, 2 H), 7.06 (t, 1 H), 7.00 (t, 2 H), 6.94 (d, 2 H), 6.76 (d, 1 H), 6.48 (dd, 1 H), 4.78 (s, 2 H), 4.18 (s, 2 H), 4.07 (s, 2 H), 2.92 (s, 3 H), 2.41 (s, 3 H); MS (ESI-) m/z 685 (M+79).

5

Example 315

(5-(4-(((4-bromo-2-fluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid

10 The product from Example 306A and 2-fluoro-4-bromo-1-(bromomethyl)benzene were processed as described in Example 306B-E to provide the title compound. ¹H NMR (500 MHz, DMSO-D₆) δ ppm 12.67-13.36 (br.s, 1 H), 8.95 (s, 1 H), 7.43 (dd, 1 H), 7.39 (d, 1 H), 7.29 (t, 3 H), 7.18 (t, 1 H), 7.06 (t, 1 H), 7.00 (d, 1 H), 6.96 (d, 1 H), 6.93 (d, 2 H), 6.76 (d, 1 H), 6.47 (dd, 1 H), 4.78 (s, 2 H), 4.07 (d, 4 H), 2.90 (s, 3 H), 2.33 (s, 3 H); MS (ESI-) m/z 679 (M+H).

15

Example 316

N-((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)glycine

20 The product from Example 91I was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 8.18 (t, 1 H), 7.40 (d, 1 H), 6.90-7.33 (m, 11 H), 6.82 (d, 1 H), 6.49 (dd, 1 H), 4.62 (s, 2 H), 4.08 (s, 2 H), 4.06 (s, 2 H), 3.83 (d, 2 H), 2.90 (s, 3 H), 2.33 (s, 3 H); MS (ESI) m/z 674 (M+H⁺).

25

Example 317

N-((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)-beta-alanine

30 The product from Example 91I and β-alanine ethyl ester hydrochloride was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.94 (t, 1 H), 7.40 (d, 1 H), 6.87-7.35 (m, 10 H), 6.76 (d, 1 H), 6.50 (dd, 1 H), 4.56 (s, 2 H), 4.08 (s, 2 H), 4.06 (s, 2 H), 3.32 (q, 2 H), 2.90 (s, 3 H), 2.39 (t, 3 H), 2.33 (s, 3 H); MS (ESI) m/z 688 (M+H⁺).

35

Example 318

4-(((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)amino)butanoic
acid

The product from Example 91I and ethyl 4-aminobutyrate hydrochloride was
processed as described in Example 104B to provide the title compound. ¹H NMR (300
MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.97 (t, 1 H), 7.39 (d, 1 H), 6.88-7.33 (m, 11 H), 6.72 (d, 1
H), 6.49 (dd, 1 H), 4.55 (s, 2 H), 4.07 (s, 2 H), 4.05 (s, 2 H), 3.11 (q, 2 H), 2.90 (s, 3 H),
2.33 (s, 3 H), 2.19 (t, 2 H), 1.61 (m, 2 H); MS (ESI) m/z 702 (M+H⁺).

Example 319

4-((4-(5-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-
chlorophenoxy)butanoyl)amino)butanoic acid

Example 319A

N-(4-(3-(allyloxy)-4-chlorophenoxy)benzyl)-N-benzyl-N-(2-methyl-3-nitrophenyl)amine

The product from Example 91D and benzyl bromide was processed as described in
Example 6B to provide the title compound.

Example 319B

ethyl 4-(5-(4-((benzyl(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)-2-
chlorophenoxy)butanoate

The product from Example 319A was processed as described in Examples 33A and
116A to provide the title compound.

Example 319C

4-(5-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-
chlorophenoxy)butanoic acid

The product from Example 319B was processed as described in Examples 6C, 6D,
and 50 to provide the title compound.

Example 319D

4-((4-(5-(4-((benzyl(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-

chlorophenoxy)butanoyl)amino)butanoic acid

The product from Example 319C and ethyl 4-aminobutyrate hydrochloride was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.84 (t, 1 H), 7.36 (d, 1 H), 7.24 (m, 7 H), 6.99 (m, 5 H), 6.83 (d, 1 H), 6.45 (dd, 1 H), 4.06 (s, 2 H), 4.03 (s, 2 H), 3.99 (t, 2 H), 3.04 (dd, 2 H), 2.92 (s, 3 H), 2.40 (s, 3 H), 2.22 (m, 5 H), 1.92 (m, 2 H), 1.60 (m, 2 H); MS (ESI) m/z 694 (M+H⁺).

Example 320

4-((4-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic acid

Example 320A

ethyl 4-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenoxy)butanoate

The product from Example 91F was processed as described in Example 116A to provide the title compound.

Example 320B

4-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoic acid

The product from Example 320A was processed as described in Examples 6C, 6D, and 50 to provide the title compound.

Example 320C

4-((4-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic acid

The product from Example 320B and ethyl 4-aminobutyrate hydrochloride was processed as described in Example 104B to provide the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 12.03 (s, 1 H), 8.97 (s, 1 H), 7.85 (t, 1 H), 7.37 (d, 1 H), 6.89-7.33 (m, 10 H), 6.83 (d, 1 H), 6.45 (dd, 1 H), 4.08 (s, 2 H), 4.06 (s, 2 H), 3.99 (t, 2 H), 3.04 (m, 2 H),

2.90 (s, 3 H), 2.33 (s, 3 H), 2.21 (m, 4 H), 1.91 (m, 2 H), 1.60 (m, 2 H); MS (ESI) m/z 730 ($M+H^+$).

5

Example 321

(2R)-2-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)propanoic acid

10

Example 321A

methyl (2R)-2-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino)methyl)phenoxy)phenoxy)propanoate

The product from Example 91F and methyl-(*R*)-lactate was processed as described in Example 62A to provide the title compound.

15

Example 321B

(2R)-2-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)propanoic acid

20

The product from Example 321A was processed as described in Examples 6C, 6D and 50 to provide the title compound. 1H NMR (300 MHz, DMSO- d_6) δ 13.12 (s, 1 H), 8.97 (s, 1 H), 7.38 (d, 1 H), 6.82-7.34 (m, 10 H), 6.66 (d, 1 H), 6.47 (dd, 1 H), 4.90 (q, 1 H), 4.07 (s, 2 H), 4.05 (s, 2 H), 2.90 (s, 3 H), 2.33 (s, 3 H), 1.51 (d, 3 H); MS (ESI) m/z 631 ($M+H^+$).

25

Example 322

N-(3-((4-(4-chloro-3-hydroxyphenoxy)benzyl)(2,4-difluorobenzyl)amino)-2-
methylphenyl)methanesulfonamide

30

The product from Example 91F was processed as described in Examples 6C and 6D to provide the title compound. 1H NMR (300 MHz, $CDCl_3$) δ 7.07-7.29 (m, 5 H), 6.91 (m, 3 H), 6.73 (m, 3 H), 6.59 (d, 1 H), 6.52 (dd, 1 H), 6.10 (s, 1 H), 5.57 (s, 1 H), 4.09 (s, 2 H), 4.06 (s, 2 H), 2.95 (s, 3 H), 2.29 (s, 3 H); MS (ESI) m/z 559 ($M+H^+$).

35

Example 323

(2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

5

Example 323A

N-(4-(3-(allyloxy)-4-bromophenoxy)benzyl)-N-(2,4-difluorobenzyl)-N-(2-methyl-3-
nitrophenyl)amine

The product from Example 63D and 2,4-difluorobenzyl bromide were processed as described in Example 6B to provide the title compound. MS (ESI+) m/z 597 (M+H)⁺.

10

Example 323B

2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-nitrophenyl)amino)methyl)phenoxy)phenol

The product from Example 323A was processed as described in Example 33A to provide the title compound. MS (ESI+) m/z 557 (M+H)⁺.

15

Example 323C

ethyl (2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
nitrophenyl)amino)methyl)phenoxy)phenoxy)acetate

20

The product from Example 323B and ethyl glycolate were processed as described in Example 62A to provide the title compound. MS (ESI+) m/z 641 (M+H)⁺.

25

Example 323D

ethyl (2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetate

The product from Example 323C was processed as described in Example 6C and D to provide the title compound. MS (ESI+) m/z 689 (M+H)⁺.

30

Example 323E

(2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid

The product from Example 323D was processed as described in Example 50 to provide the title compound. ¹H NMR (500 MHz, CDCl₃) δ 7.48 (d, 1 H), 7.35 (dd, 1 H), 7.28 (dd, 1 H), 7.21 (m, 2 H), 7.09 (s, 1 H), 7.03 (d, 2 H), 7.03 (d, 2 H), 6.78 (dd, 1 H), 6.73 (m, 1 H), 6.63 (dd, 1 H), 6.55 (d, 1 H), 4.64 (s, 2 H), 4.35 (s, 2 H), 4.20 (s, 2 H), 2.89 (s, 3

35

H), 2.01 (s, 3 H)); MS (ESI+) m/z 663 (M+H)⁺.

Example 324

5 N-(4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-beta-alanine

 The product from Example 120D and β-alanine ethyl ester hydrochloride was
processed as described in Example 104B to provide the title compound. ¹H NMR (300
MHz, DMSO-d₆) δ 8.97 (s, 1 H), 7.90 (t, 1 H), 6.87-7.35 (m, 11 H), 6.68 (m, 1 H), 6.48 (m,
10 2 H), 4.08 (s, 2 H), 4.05 (s, 2 H), 3.90 (t, 2 H), 3.64 (s, 1 H), 3.22 (dd, 2 H), 2.90 (s, 3 H),
2.35 (m, 2 H), 2.33 (s, 3 H), 2.19 (t, , 2 H), 1.87 (m, 2 H); MS (ESI) m/z 682 (M+H)⁺.

Example 325

15 4-((4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic
 acid

 The product from Example 120D and ethyl 4-aminobutyrate hydrochloride was
processed as described in Example 104B to provide the title compound. ¹H NMR (300
20 MHz, DMSO-d₆) δ 8.96 (s, 1 H), 7.83 (t, 1 H), 6.88-7.32 (m, 11 H), 6.68 (m, 1 H), 6.49 (m,
2 H), 4.08 (s, 2 H), 4.05 (s, 2 H), 3.91 (t, 2 H), 3.03 (dd, 2 H), 2.90 (s, 3 H), 2.33 (s, 3 H),
2.19 (t, 4 H), 2.07 (s, 1 H), 1.89 (m, 2 H), 1.59 (m, 2 H); MS (ESI) m/z 696 (M+H)⁺.

Example 326

25 N-(4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-N-
 methylglycine

 The product from Example 120D and sarcosine ethyl ester hydrochloride was
30 processed as described in Example 104B to provide the title compound. ¹H NMR (300
MHz, DMSO-d₆) δ 8.96 (s, 1 H), 6.85-7.34 (m, 11 H), 6.69 (m, 1 H), 6.49 (m, 2 H), 4.08 (s,
2 H), 4.05 (s, 2 H), 3.94 (m, 4 H), 2.98 (s, 3 H), 2.89 (s, 3 H), 2.80 (s, 1 H), 2.50 (m, 2 H),
2.33 (s, 3 H), 1.90 (m, 2 H); MS (ESI) m/z 682 (M+H)⁺.

35 Compounds of the present invention may exist as stereoisomers wherein,

asymmetric or chiral centers are present. These stereoisomers are "R" or "S" depending on the configuration of substituents around the chiral carbon atom. The terms "R" and "S" used herein are configurations as defined in IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem., 1976, 45: 13-30. The present invention
5 contemplates various stereoisomers and mixtures thereof and are specifically included within the scope of this invention. Stereoisomers include enantiomers and diastereomers, and mixtures of enantiomers or diastereomers. Individual stereoisomers of compounds of the present invention may be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of racemic mixtures
10 followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary or (2) direct separation of the mixture of optical enantiomers on chiral chromatographic columns.

15 Geometric isomers can also exist in the compounds of the present invention. The present invention contemplates the various geometric isomers and mixtures thereof resulting from the arrangement of substituents around a carbon-carbon double bond or arrangement of substituents around a ring. Substituents around a carbon-carbon double bond are designated as being in the Z or E configuration where the term "Z" represents substituents
20 on the same side of the carbon-carbon double bond and the term "E" represents substituents on opposite sides of the carbon-carbon double bond. The arrangement of substituents around a ring are designated as cis or trans where the term "cis" represents substituents on the same side of the plane of the ring and the term "trans" represents substituents on opposite sides of the plane of the ring. Mixtures of compounds where the substituents are
25 disposed on both the same and opposite sides of plane of the ring are designated cis/trans.

The term "pharmaceutically acceptable prodrugs" represents those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals with
30 undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention.

The term "prodrug" represents compounds which are rapidly transformed in vivo to a compound of formula (I-IV), for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of
35 the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987.

The term "pharmaceutically acceptable salt" represents those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans

and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, et al. describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 66:1-19 (1977). The salts can be prepared in situ during the final isolation and purification of the compounds of the invention, or separately by reacting the free base function with a suitable organic acid. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphersulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like.

The term "pharmaceutically acceptable carrier," as used herein, means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols; such a propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

The present invention also provides pharmaceutical compositions, which comprise compounds of the present invention formulated together with one or more non-toxic pharmaceutically acceptable carriers. The pharmaceutical compositions may be specially formulated for oral administration in solid or liquid form, for parenteral injection, or for

rectal administration.

Further included within the scope of the present invention are pharmaceutical compositions comprising one or more of the compounds of formula (I-IV) prepared and formulated in combination with one or more non-toxic pharmaceutically acceptable compositions. The pharmaceutical compositions can be formulated for oral administration in solid or liquid form, for parenteral injection or for rectal administration.

The pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray. The term "parenteral" administration refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. Conversely, reduced particle size may maintain biological activity.

These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of

drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides) Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

5 The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

10 Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating
15 agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid
20 polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

25 The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding
30 compositions which can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

35 Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut,

corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers, or propellants which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions, and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated, and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required for to achieve the desired therapeutic effect and to gradually increase the dosage until the desired

effect is achieved.


Generally, dosage levels of about 1 to about 50 mg/kg/day, more preferably of about 5 to about 25 mg/kg/day of active compound are administered orally to a mammalian patient. If desired, the effective daily dose may be divided into multiple doses for purposes of administration, e.g. two to four separate doses per day.

It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention, which is defined solely by the appended claims and their equivalents. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art.

Such changes and modifications, including without limitation those relating to the chemical structures, substituents, derivatives, intermediates, syntheses, formulations and/or methods of use of the invention, may be made without departing from the spirit and scope thereof.

What is claimed is:

1. A compound having formula (I)



5 or a pharmaceutically acceptable salt or prodrug thereof, wherein

L_A is a covalent bond or $C(X_{A1})(X_{A2})$;

L_D is a covalent bond or $C(X_{D1})(X_{D2})$;

X_{A1} is selected from the group consisting of hydrogen, alkoxy, alkoxycarbonyl, alkoxyalkyl, alkyl, alkylcarbonyl, cyano, halogen, hydroxy, hydroxyalkyl, nitro and
10 $(NR_{10}R_{11})$ carbonyl;

X_{A2} is selected from the group consisting of hydrogen and alkyl;

or X_{A1} and X_{A2} together are oxo;

X_{D1} is selected from the group consisting of hydrogen, alkoxy, alkoxycarbonyl, alkoxyalkyl, alkyl, alkylcarbonyl, cyano, halogen, hydroxy, hydroxyalkyl, nitro and
15 $(NR_{10}R_{11})$ carbonyl;

X_{D2} is selected from the group consisting of hydrogen and alkyl;

or X_{D1} and X_{D2} together are oxo;

L_B is absent or selected from the group consisting of a covalent bond, alkenylene, alkylene, alkynylene, $-(CH_2)_mC(O)(CH_2)_n-$, $-(CH_2)_mC(O)(CH_2)_nCH=CH-$,

20 $-(CH_2)_mC(O)(CH_2)_nC\equiv C-$, $-(CH_2)_mCH(OH)(CH_2)_n-$, $-(CH_2)_mCH(OH)(CH_2)_nCH_2CH=CH-$, $-(CH_2)_mCH(OH)(CH_2)_nCH_2C\equiv C-$, $-(CH_2)_mO(CH_2)_n-$, $-(CH_2)_mO(CH_2)_nCH_2CH=CH-$,

$-(CH_2)_mO(CH_2)_nCH_2C\equiv C-$, $-(CH_2)_mS(CH_2)_n-$, $-(CH_2)_mS(O)(CH_2)_n-$, $-(CH_2)_mS(O)_2(CH_2)_n-$, $-(CH_2)_mN(R_7)(CH_2)_n-$, $-(CH_2)_mN(R_7)C(O)(CH_2)_n-$, $-(CH_2)_mC(O)N(R_7)(CH_2)_n-$,

$O(CH_2)_mC(O)N(R_7)(CH_2)_n-$, $CH_2)_mC(O)O(CH_2)_n-$, $-O(CH_2)_mC(O)(CH_2)_n-$,

25 $(CH_2)_mN(R_7)S(O)_2(CH_2)_n-$, $-(CH_2)_mS(O)_2N(R_7)(CH_2)_n-$, $-(CH_2)_mS(O)_2O(CH_2)_n-$ and

$-(CH_2)_mOS(O)_2(CH_2)_n-$ wherein each group is inserted as drawn with the left end attached to A and the right end attached to B;

m is an integer of 0-6;

n is an integer of 0-6;

30 L_C is absent or selected from the group consisting of a covalent bond, alkenylene,

alkylene, alkynylene, $-(CH_2)_pC(O)(CH_2)_q-$, $-(CH_2)_pC(O)(CH_2)_qCH=CH-$,
 $-(CH_2)_pC(O)(CH_2)_qC\equiv C-$, $-(CH_2)_pCH(OH)(CH_2)_q-$, $-(CH_2)_pCH(OH)(CH_2)_qCH_2CH=CH-$, -
 $(CH_2)_pCH(OH)(CH_2)_qCH_2C\equiv C-$, $-(CH_2)_pO(CH_2)_q-$, $-(CH_2)_pO(CH_2)_qCH_2CH=CH-$, -
 $(CH_2)_pO(CH_2)_qCH_2C\equiv C-$, $-(CH_2)_pS(CH_2)_q-$, $-(CH_2)_pS(O)(CH_2)_q-$, $-(CH_2)_pS(O)_2(CH_2)_q-$, -
5 $(CH_2)_pN(R_7)(CH_2)_q-$, $-(CH_2)_pN(R_7)C(O)(CH_2)_q-$, $-(CH_2)_pC(O)N(R_7)(CH_2)_q-$, -
 $O(CH_2)_pC(O)N(R_7)(CH_2)_q-$, $-(CH_2)_pC(O)O(CH_2)_q-$, $-O(CH_2)_pC(O)(CH_2)_q-$, -
 $(CH_2)_pN(R_7)S(O)_2(CH_2)_q-$, $-(CH_2)_pS(O)_2N(R_7)(CH_2)_q-$, $-(CH_2)_pS(O)_2O(CH_2)_q-$ and -
 $(CH_2)_pOS(O)_2(CH_2)_q-$ wherein each group is inserted as drawn with the left end attached to
B and the right end attached to C;

10 p is an integer of 0-6;

q is an integer of 0-6;

L_E is absent or selected from the group consisting of a covalent bond, alkenylene,
alkylene, alkynylene, $-(CH_2)_rC(O)(CH_2)_s-$, $-(CH_2)_rC(O)(CH_2)_sCH=CH-$,
 $-(CH_2)_rC(O)(CH_2)_sC\equiv C-$, $-(CH_2)_rCH(OH)(CH_2)_s-$, $-(CH_2)_rCH(OH)(CH_2)_sCH_2CH=CH-$,
15 $-(CH_2)_rCH(OH)(CH_2)_sCH_2C\equiv C-$, $-(CH_2)_rO(CH_2)_s-$, $-(CH_2)_rO(CH_2)_sCH_2CH=CH-$,
 $-(CH_2)_rO(CH_2)_sCH_2C\equiv C-$, $-(CH_2)_rS(CH_2)_s-$, $-(CH_2)_rS(O)(CH_2)_s-$,
 $-(CH_2)_rS(O)_2(CH_2)_s-$, $-(CH_2)_rN(R_7)(CH_2)_s-$, $-(CH_2)_rN(R_7)C(O)(CH_2)_s-$,
 $-(CH_2)_rC(O)N(R_7)(CH_2)_s-$, $-O(CH_2)_rC(O)N(R_7)(CH_2)_s-$, $-(CH_2)_rC(O)O(CH_2)_s-$, -
 $O(CH_2)_rC(O)(CH_2)_s-$,
20 $-(CH_2)_rN(R_7)S(O)_2(CH_2)_s-$, $-(CH_2)_rS(O)_2N(R_7)(CH_2)_s-$, $-(CH_2)_rS(O)_2O(CH_2)_s-$ and
 $-(CH_2)_rOS(O)_2(CH_2)_s-$ wherein each group is inserted as drawn with the left end attached to
D and the right end attached to E;

r is an integer of 0-6;

s is an integer of 0-6;

25 L_F is absent or selected from the group consisting of a covalent bond, alkenylene,
alkylene, alkynylene, $-(CH_2)_uC(O)(CH_2)_v-$, $-(CH_2)_uC(O)(CH_2)_vCH=CH-$,
 $-(CH_2)_uC(O)(CH_2)_vC\equiv C-$, $-(CH_2)_uCH(OH)(CH_2)_v-$, $-(CH_2)_uCH(OH)(CH_2)_vCH_2CH=CH-$,
 $-(CH_2)_uCH(OH)(CH_2)_vCH_2C\equiv C-$, $-(CH_2)_uO(CH_2)_v-$, $-(CH_2)_uO(CH_2)_vCH_2CH=CH-$,
 $-(CH_2)_uO(CH_2)_vCH_2C\equiv C-$, $-(CH_2)_uS(CH_2)_v-$, $-(CH_2)_uS(O)(CH_2)_v-$,
30 $-(CH_2)_uS(O)_2(CH_2)_v-$, $-(CH_2)_uN(R_7)(CH_2)_v-$, $-(CH_2)_uN(R_7)C(O)(CH_2)_v-$,
 $-(CH_2)_uC(O)N(R_7)(CH_2)_v-$, $-O(CH_2)_uC(O)N(R_7)(CH_2)_v-$, $-(CH_2)_uC(O)O(CH_2)_v-$, -
 $O(CH_2)_uC(O)(CH_2)_v-$,
 $-(CH_2)_uN(R_7)S(O)_2(CH_2)_v-$, $-(CH_2)_uS(O)_2N(R_7)(CH_2)_v-$, $-(CH_2)_uS(O)_2O(CH_2)_v-$ and
 $-(CH_2)_uOS(O)_2(CH_2)_v-$ wherein each group is inserted as drawn with the left end attached to
35 E and the right end attached to F;

u is an integer of 0-6;

v is an integer of 0-6;

A and D are each independently selected from the group consisting of aryl,

R₈ and R₉ are each independently selected from the group consisting of hydrogen, alkenyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkylcarbonyl, alkylsulfonyl, alkynyl, carboxyalkylcarbonyl; cyanoalkyl, formyl, hydroxy, hydroxyalkyl, -NR₁₀R₁₁, (NR₁₀R₁₁)carbonyl and carboxyalkyl wherein the alkyl portion of carboxyalkyl is optionally substituted with one or two substituents selected from the group consisting of alkylthio, aryl, heterocycle, hydroxy, carboxy, -NR₁₀R₁₁ and (NR₁₀R₁₁)carboxy;

R₁₀ and R₁₁ are each independently selected from the group consisting of hydrogen and alkyl;

R₁₂ is selected from the group consisting of alkoxy, alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl; and

R₁₃ and R₁₄ are each independently selected from the group consisting of hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl.

2. A compound according to claim 1 wherein

L_A is a covalent bond;

L_D is a covalent bond;

A is aryl; and

D is aryl.

3. A compound according to claim 1 wherein

L_A is a covalent bond;

L_D is a covalent bond;

A is heterocycle; and

D is aryl.

4. A compound according to claim 1 wherein

L_A is a covalent bond;

L_D is a covalent bond;

A is heterocycle; and

D is heterocycle.

5. A compound according to claim 1 wherein

L_A is C(X_{A1})(X_{A2});

L_D is a covalent bond;

A is aryl; and

D is aryl.

6. A compound according to claim 1 wherein

L_A is $C(X_{A1})(X_{A2})$;
 L_D is a covalent bond;
A is heterocycle; and
D is aryl.

5

7. A compound according to claim 1 wherein
 L_A is $C(X_{A1})(X_{A2})$;
 L_D is a covalent bond;
A is heterocycle; and
D is heterocycle.

10

8. A compound according to claim 1 wherein
 L_A is $C(X_{A1})(X_{A2})$;
 L_D is a covalent bond;
A is aryl; and
D is heterocycle.

15

9. A compound according to claim 1 wherein
 L_A is $C(X_{A1})(X_{A2})$;
 L_D is $C(X_{D1})(X_{D2})$;
A is heterocycle; and
D is heterocycle.

20

10. A compound according to claim 1 wherein
 L_A is $C(X_{A1})(X_{A2})$;
 L_D is $C(X_{D1})(X_{D2})$;
A is aryl; and
D is heterocycle.


25

11. A compound according to claim 1 wherein
 L_A is $C(X_{A1})(X_{A2})$;
 L_D is $C(X_{D1})(X_{D2})$;
A is aryl; and
D is aryl.

30

12. A compound according to claim 1 of formula (II)

35



or a pharmaceutically acceptable salt or prodrug thereof.

- 5 13. A compound according to claim 12 wherein
 R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
- 10 14. A compound according to claim 13 selected from the group consisting of
N-{3-[bis(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-(dibenzylamino)-2-methylphenyl}methanesulfonamide;
N-{3-(dibenzylamino)-2-methylphenyl}ethanesulfonamide;
N-{3-(dibenzylamino)-2-methylphenyl}-2-propanesulfonamide;
15 N-{3-[benzyl(4-methoxycarbonylbenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-nitrobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamid;
20 N-{3-[benzyl(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-cyanobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-bromobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(3-methoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
25 N-{3-[(1,3-benzodioxol-5-ylmethyl)(benzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[benzyl(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-chlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
30 N-{3-[benzyl(2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[[4-(allyloxy)benzyl](benzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[[4-(allyloxy)benzyl](2,4-difluorobenzyl)amino]-2-


- methylphenyl}methanesulfonamide;
N-{3-[(2-cyanobenzyl)(2-fluoro-4-methoxybenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[(2,4-difluorobenzyl)(4-methoxybenzyl)amino]-2-
5 methylphenyl}methanesulfonamide;
N-{3-[(2,4-difluorobenzyl)(2-fluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[bis(2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[(2-cyanobenzyl)(2,4-difluorobenzyl)amino]-2-
10 methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-methylbenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-methylbenzyl)amino]-2-methylphenyl}methanesulfonamide;
N-{3-[benzyl(4-chlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
15 N-(3-{benzyl[4-(trifluoromethyl)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
N-(3-{benzyl[2-fluoro-4-(trifluoromethyl)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
N-{3-[benzyl(2,4-dichlorobenzyl)amino]-2-methylphenyl}methanesulfonamide;
20 N-{3-[benzyl(4-{[3-bromo-2-propenyl]oxy}benzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-(3-{benzyl[4-(methoxymethyl)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
N-(3-{benzyl[4-(hydroxymethyl)benzyl]amino}-2-
25 methylphenyl)methanesulfonamide;
N-(3-{benzyl[4-(2-hydroxyethoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
N-{3-[(2,4-difluorobenzyl)(4-propoxybenzyl)amino]-2-
methylphenyl}methanesulfonamide;
30 N-[3-(dibenzylamino)-2-ethylphenyl]methanesulfonamide;
N-{3-[benzyl(4-bromo-2-fluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[benzyl(2-chloro-4-fluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
35 N-{3-[benzyl(4-{[(2Z)-3-bromoprop-2-enyl]oxy}benzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-{3-[(2,4-difluorobenzyl)(4-ethoxybenzyl)amino]-2-
methylphenyl}methanesulfonamide; and

N-{3-[benzyl(4-chloro-2-fluorobenzyl)amino]-2-methylphenyl}methanesulfonamide.

15. A compound according to claim 12 wherein
5 R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen;
R₄ is alkyl; and
R₆ is (NR₁₂R₁₃)sulfonyl.
16. A compound according to claim 15 that is N'-[3-(dibenzylamino)-2-methylphenyl]-
10 N,N-dimethylsulfamide.
17. A compound according to claim 12 wherein
R₂, R₃, R₄, R₅, X_{A1} and X_{D1} are each hydrogen;
R₁ is alkoxy; and
15 R₆ is alkylsulfonyl.
18. A compound according to claim 17 that is N-{3-[bis(2-bromobenzyl)amino]-4-methoxyphenyl}methanesulfonamide.
19. A compound according to claim 12 wherein
20 X_{A1} is cyano;
R₁, R₂, R₃, R₅ and X_{D1} are each hydrogen;
R₄ is alkyl; and
R₆ is alkylsulfonyl.
20. A compound according to claim 19 that is N-(3-{(2-bromobenzyl)[cyano(phenyl)methyl]amino}-2-methylphenyl)methanesulfonamide.
21. A compound according to claim 12 wherein
30 R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen;
R₄ is selected from the group consisting of alkenyl, alkoxyalkyl, alkoxycarbonylalkenyl and hydroxyalkyl; and
R₆ is alkylsulfonyl.
22. A compound according to claim 21 selected from the group consisting of
35 N-[3-(dibenzylamino)-2-((E)-3-ethoxy-3-oxo-1-propenyl)phenyl]methanesulfonamide;
N-[3-(dibenzylamino)-2-(hydroxymethyl)phenyl]methanesulfonamide;

N-[3-(dibenzylamino)-2-vinylphenyl]methanesulfonamide;
 N-[3-(dibenzylamino)-2-(methoxymethyl)phenyl]methanesulfonamide; and
 N-[3-(dibenzylamino)-2-(ethoxymethyl)phenyl]methanesulfonamide.

- 5 23. A compound according to claim 1 of formula (III)



or a pharmaceutically acceptable salt or prodrug thereof.

- 10 24. A compound according to claim 23 wherein

B is aryl wherein said aryl is phenyl;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m and n are each 0;

R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;

15 R_4 is alkyl; and

R_6 is alkylsulfonyl.

25. A compound according to claim 24 selected from the group consisting of

20 N-{3-[[4-(4-bromophenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[[4-(4-(3-ethoxy-3-oxopropyl)phenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

N-{3-[[4-(4-(2-carboxyethyl)phenoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide;

25 N-{3-[(2,4-difluorobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;

methyl 4-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate;

N-{3-[(2,4-difluorobenzyl)(2-fluoro-4-phenoxybenzyl)amino]-2-

- methylphenyl}methanesulfonamide;
 N-(3-{(2,4-difluorobenzyl)[4-(3-methoxyphenoxy)benzyl]amino}-2-
 methylphenyl)methanesulfonamide;
 N-{3-[(2-fluorobenzyl)(4-phenoxybenzyl)amino]-2-
 5 methylphenyl}methanesulfonamide;
 N-{3-[(4-methoxybenzyl)(4-phenoxybenzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 3-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoic acid;
 10 N-[3-(benzyl{4-[3-(2-methoxyethoxy)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 (3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
 4-[4-{4-[(benzyl{2-methyl-3-
 15 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-(2-
 methoxyethoxy)phenyl]butanoic acid;
 ethyl 4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoate;
 N-[3-(benzyl{4-[3-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-
 20 methylphenyl]methanesulfonamide;
 methyl 4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate;
 ethyl N-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoyl)-beta-alaninate;
 25 methyl 3-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}benzoate;
 N-{3-[benzyl(4-{3-[2-(2-methoxyethoxy)ethoxy]phenoxy}benzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 N-[3-(3-{4-[(benzyl{2-methyl-3-
 30 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl)acetamide;
 N-[3-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propyl]methanesulfonami
 de;
 N-{3-[benzyl(4-{3-[3-(dimethylamino)propoxy]phenoxy}benzyl)amino]-2-
 35 methylphenyl}methanesulfonamide;
 4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoic acid;
 3-(4-{4-[(2-bromobenzyl){2-methyl-3-

- [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)propanoic acid;
 (5-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-2-bromophenoxy)acetic acid;
 4-{2-(3-{4-[(benzyl{2-methyl-3-
 5 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)ethyl]amino}-4-
 oxobutanoic acid;
 5-{2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)ethyl]amino}-5-
 oxopentanoic acid;
 10 N-[2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)ethyl]acetamide;
 N-[2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)ethyl]methanesulfonamid
 e;
 15 methyl 2-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)ethylcarbamate;
 (3-{4-[(4-chloro-2-fluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetic acid;
 3-(4-{4-[(2-fluorobenzyl){2-methyl-3-
 20 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)propanoic acid;
 3-(4-{4-[(4-fluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)propanoic acid;
 3-(4-{4-[(4-chloro-2-fluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenyl)propanoic acid;
 25 (3-{4-[(2,4-difluorobenzyl){2-ethyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetic acid;
 (3-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-5-methylphenoxy)acetic acid;
 (3-{4-[(2,4-difluorobenzyl){2-methyl-3-
 30 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}-5-methoxyphenoxy)acetic acid;
 (2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetic acid;
 (4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}phenoxy)acetic acid;
 35 4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy}benzoic acid;
 N-[3-(benzyl{4-[4-(methoxymethoxy)phenoxy]benzyl} amino)-2-
 methylphenyl]methanesulfonamide;

- N-[3-(benzyl{4-[4-(3-hydroxypropyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoic acid;
- 5 N-[3-(benzyl{4-[4-(4-hydroxybutyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[(3-bromobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide
- N-{3-[(4-bromobenzyl)(4-phenoxybenzyl)amino]-2-methylphenyl}methanesulfonamide;
- 10 N-[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]glycine;
- N-[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]-beta-alanine;
- 15 4-{[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]amino}butanoic acid;
- N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]glycine;
- 20 N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-beta-alanine;
- 4-{[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]amino}butanoic acid;
- 25 (3-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
- 2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)propanoic acid;
- 2-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-4-hydroxybutanoic acid;
- 30 N-(3-{benzyl[4-(3-hydroxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- 4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid;
- 35 5-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid;
- N-[3-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoyl]-N-

- methylglycine;
 N-[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-N-
 methylglycine;
- 5 4-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid;
 5-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid;
 N-[(3-{4-[(benzyl{2-methyl-3-
 10 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]glycine;
 N-[(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-beta-alanine;
 N-[(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-N-methylglycine;
- 15 4-{[(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]amino}butanoic
 acid;
 ethyl 4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoate;
- 20 ethyl 5-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoate;
 N-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]glycine;
 N-[4-(3-{4-[(benzyl{2-methyl-3-
 25 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-beta-alanine;
 4-{[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]amino}butanoic
 acid;
 N-[5-(3-{4-[(benzyl{2-methyl-3-
 30 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]glycine;
 N-[5-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-beta-alanine;
 4-{[5-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]amino}butanoi
 35 c acid;
 N-[3-(benzyl{4-[3-(2-hydroxyethoxy)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 [2-(3-{4-[(benzyl{2-methyl-3-

- [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethoxy]acetic acid;
2,4-dideoxy-6-O-(3-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-D-erythro-hexonic acid;
(3-{4-[(2-bromobenzyl){2-methyl-3-
5 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
N-{3-[[4-(3-acetylphenoxy)benzyl](2,4-difluorobenzyl)amino]-2-
methylphenyl}methanesulfonamide;
N-(3-{(2,4-difluorobenzyl)[4-(3,4-dimethoxyphenoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
10 N-(3-{(2,4-difluorobenzyl)[4-(pent-3-ynyloxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
N-(3-{(4-chloro-2-fluorobenzyl)[4-(methylthio)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
N-{3-[(4-chloro-2-fluorobenzyl)(2-fluorobenzyl)amino]-2-
15 methylphenyl}methanesulfonamide;
N-(3-{benzyl[(2'-cyano-1,1'-biphenyl-4-yl)methyl]amino}-2-
methylphenyl)methanesulfonamide;
3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(5-
20 hydroxypentyl)propanamide;
3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(4-
hydroxybutyl)propanamide;
3-(4-{4-[(benzyl{2-methyl-3-
25 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(3-
hydroxypropyl)propanamide;
3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-
hydroxyethyl)propanamide;
30 3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(6-
hydroxyhexyl)propanamide;
N-(3-{benzyl[4-(3-isopropoxyphenoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
35 N-[3-(benzyl{4-[3-(cyclobutyloxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide;
N-(3-{benzyl[4-(3-sec-butoxyphenoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;

- N-[3-(benzyl{4-[3-(cyclopentyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(1-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 5 N-[3-(benzyl{4-[3-(2-methoxy-1-methylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(cyclohexyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[benzyl(4-{3-[(3-methylcyclopentyl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 10 N-[3-(benzyl{4-[3-(2-ethoxy-1-methylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[benzyl(4-{3-[(4-methylcyclohexyl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 15 N-[3-(benzyl{4-[3-(cycloheptyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-(3-{benzyl[4-(3-methoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- N-(3-{benzyl[4-(3-ethoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- 20 N-(3-{benzyl[4-(3-propoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- N-[3-(benzyl{4-[3-(cyclopropylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 25 N-(3-{benzyl[4-(3-butoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- N-(3-{benzyl[4-(3-isobutoxyphenoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;
- N-[3-(benzyl{4-[3-(pent-3-ynyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 30 N-{3-[benzyl(4-{3-[(2E)-pent-2-enyloxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- N-{3-[benzyl(4-{3-[(1-methylcyclopropyl)methoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 35 N-[3-(benzyl{4-[3-(cyclobutylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

- N-[3-(benzyl{4-[3-(pentyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(2-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 5 N-[3-(benzyl{4-[3-(3-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(2-ethoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 10 N-[3-(benzyl{4-[3-(2-(methylthio)ethoxy]phenoxy}benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(cyclopentylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(hexyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 15 N-[3-(benzyl{4-[3-(3,3-dimethylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(2-isopropoxyethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(cyclohexylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 20 N-[3-(benzyl{4-[3-(3-methoxy-3-methylbutoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- (3-{4-[(2-methylbenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
- 25 (3-{4-[(4-methylbenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
- (3-{4-[(2,4-dichlorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
- (3-{4-[(2-chloro-4-fluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
- 30 (3-{4-[(3,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid;
- N-[3-(benzyl{4-[4-(4-hydrazino-4-oxobutyl)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 35 N~2~[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-asparagine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-valine;

- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-tyrosine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-methionine;
- 5 N-[2-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-lysine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-serine;
- 10 N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-phenylalanine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-tyrosine;
- N-[2-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-glutamine;
- 15 N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-isoleucine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-glutamic acid;
- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-D-histidine;
- 20 N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-valine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-aspartic acid;
- 25 ethyl 4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoate;
- 4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoic acid;
- 5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoic acid;
- 30 (5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-ethylphenoxy)acetic acid;
- ethyl 4-{[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]amino}butanoat
- 35 e;
- (5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}-2-hexylphenoxy)acetic acid;
- ethyl N-[4-(3-{4-[(benzyl{2-methyl-3-

[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-N-methylglycinate;

- 5 N-[4-(3-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-N-methylglycine;
- ethyl (2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)acetate;
- 10 N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]glycine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-beta-alanine;
- 4-{[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]amino} butanoic acid;
- 15 N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-N-methylglycine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-glutamic acid;
- 20 N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-valine;
- N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-serine;
- 25 N-[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-isoleucine;
- N-2~[4-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)butanoyl]-L-glutamine;
- N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]glycine;
- 30 N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-beta-alanine;
- 4-{[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]amino} butanoic acid;
- 35 N-[5-(4-{4-[(benzyl{2-methyl-3-[(methylsulfonyl)amino]phenyl} amino)methyl]phenoxy} phenoxy)pentanoyl]-N-methylglycine;

- N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-glutamic
acid;
- 5 N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-valine;
- N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-serine;
- N-[5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-isoleucine;
- 10 4-{[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]amino}butanoic
acid;
- N-[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-N-methylglycine;
- 15 N-[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-L-valine;
- N-[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-L-serine;
- N-2-~[(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]-L-asparagine;
- 20 N-[3-(benzyl{4-[4-(2-hydrazino-2-oxoethoxy)phenoxy]benzyl}amino)-2-
methylphenyl]methanesulfonamide;
- N-(3-{benzyl[4-(3-{[(2R)-2,3-dihydroxypropyl]oxy}phenoxy)benzyl]amino}-2-
methylphenyl)methanesulfonamide;
- 25 N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamic
acid;
- N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-leucine;
- 30 N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-aspartic
acid;
- N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-valine;
- 35 N-[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-serine;
- N-2-~[4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]-L-glutamine;

- (5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-fluorophenoxy)acetic acid;
 (2-chloro-5-(4-(((2-fluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
 5 (5-(4-(((2-bromobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
 (5-(4-(((4-bromobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
 (2-chloro-5-(4-(((2,4-dichlorobenzyl)(2-methyl-3-
 10 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
 (2-chloro-5-(4-(((2-chloro-2-fluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
 (2-chloro-5-(4-(((2-chloro-4-fluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
 15 (5-(4-(((2-bromo-4-chlorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
 (2-chloro-5-(4-(((3,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;
 (5-(4-(((4-bromo-2-fluorobenzyl)(2-methyl-3-
 20 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-chlorophenoxy)acetic acid;
 N-((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)glycine;
 N-((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)-beta-alanine;
 25 4-(((2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetyl)amino)butanoic
 acid;
 4-((4-(5-(4-(benzyl(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)-2-
 30 chlorophenoxy)butanoyl)amino)butanoic acid;
 4-((4-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic
 acid;
 (2R)-2-(2-chloro-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-
 35 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)propanoic acid;
 N-(3-((4-(4-chloro-3-hydroxyphenoxy)benzyl)(2,4-difluorobenzyl)amino)-2-
 methylphenyl)methanesulfonamide;
 (2-bromo-5-(4-(((2,4-difluorobenzyl)(2-methyl-3-

((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)acetic acid;

N-(4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-

((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-beta-alanine;

4-((4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-

5 ((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)amino)butanoic acid; and

N-(4-(3-(4-(((2,4-difluorobenzyl)(2-methyl-3-

((methylsulfonyl)amino)phenyl)amino)methyl)phenoxy)phenoxy)butanoyl)-N-methylglycine.

10

26. A compound according to claim 23 wherein

B is aryl wherein said aryl is phenyl;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m is 0;

15

n is 1-6;

R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;

R_4 is alkyl; and

R_6 is alkylsulfonyl.

20

27. A compound according to claim 26 selected from the group consisting of

N-{3-[[4-(benzyloxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide; and

N-(3-{(2,4-difluorobenzyl)[4-(2-phenylethoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide.

25

28. A compound according to claim 23 wherein

B is heterocycle;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m is 0;

30

n is 1-6;

R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;

R_4 is alkyl; and

R_6 is alkylsulfonyl.

35

29. A compound according to claim 28 selected from the group consisting of

N-(3-{(2,4-difluorobenzyl)[4-(3-furylmethoxy)benzyl]amino}-2-methylphenyl)methanesulfonamide;

N-(3-{(2,4-difluorobenzyl)[4-(2-furylmethoxy)benzyl]amino}-2-

methylphenyl)methanesulfonamide;

N-{3-[[4-(1,3-benzodioxol-5-ylmethoxy)benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide ;and

5 N-{3-[[4-[(6-chloro-1,3-benzodioxol-5-yl)methoxy]benzyl](2,4-difluorobenzyl)amino]-2-methylphenyl}methanesulfonamide.

30. A compound according to claim 23 wherein
 B is aryl wherein said aryl is phenyl;
 L_B is $-(CH_2)_mO(CH_2)_nCH_2CH=CH-$
 10 m is 0;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
- 15 31. A compound according to claim 30 that is N-{3-[(2,4-difluorobenzyl)(4-{[3-phenyl-2-propenyl]oxy}benzyl)amino]-2-methylphenyl}methanesulfonamide.
- 20 32. A compound according to claim 23 wherein
 B is aryl wherein said aryl is phenyl;
 L_B is $-(CH_2)_mC(O)(CH_2)_n-$;
 m and n are each 0;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
- 25 33. A compound according to claim 32 selected from the group consisting of
 (3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]benzoyl}phenoxy)acetic acid;
 N-{3-[(4-benzoylbenzyl)(benzyl)amino]-2-methylphenyl}methanesulfonamide;
 30 N-[3-(benzyl{4-[3-(2-cyclopropylethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[3-(cyclopentylmethoxy)benzoyl]benzyl}amino)-2-methylphenyl]methanesulfonamide; and
 N-(3-{benzyl[4-(4-{[(2R)-2,3-dihydroxypropyl]oxy}benzoyl)benzyl]amino}-2-
 35 methylphenyl)methanesulfonamide.
34. A compound according to claim 23 wherein
 B is aryl wherein said aryl is phenyl;

L_B is $-(CH_2)_mS(CH_2)_n-$;

m and n are each 0;

$R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;

R_4 is alkyl; and

5 R_6 is alkylsulfonyl.

35. A compound according to claim 34 that is
N-(3-{benzyl[4-(phenylthio)benzyl]amino}-2-methylphenyl)methanesulfonamide.


10 36. A compound according to claim 23 wherein
B is aryl wherein said aryl is phenyl;
 L_B is alkylene;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
15 R_6 is alkylsulfonyl.

37. A compound according to claim 23 wherein
B is aryl wherein said aryl is phenyl;
 L_B is $-(CH_2)_mC(OH)(CH_2)_n-$;
20 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.

38. A compound according to claim 23 wherein
25 B is cycloalkyl;
 L_B is $-(CH_2)_mO(CH_2)_n-$;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.

30 39. A compound according to claim 23 wherein
B is heterocycle;
 L_B is $-(CH_2)_mC(O)(CH_2)_n-$;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
35 R_4 is alkyl; and
 R_6 is alkylsulfonyl.

40. A compound according to claim 1 of formula (IV)



or a pharmaceutically acceptable salt or prodrug thereof.

5

41. A compound according to claim 40 wherein

B is aryl wherein said aryl is phenyl;

C is aryl wherein said aryl is phenyl;

L_B is $-(CH_2)_mO(CH_2)_n-$;

10

m and n are each 0;

L_C is $-(CH_2)_pO(CH_2)_q-$;

p is 0;

R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;

R_4 is alkyl; and

15

R_6 is alkylsulfonyl.

42. A compound according to claim 41 selected from the group consisting of

N-[3-(benzyl{4-[3-(2-phenylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;

20

N-{3-[{4-[3-(1,3-benzodioxol-5-ylmethoxy)phenoxy]benzyl}(benzyl)amino]-2-methylphenyl}methanesulfonamide; and

N-[3-(benzyl{4-[3-(benzyloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide.

25

43. A compound according to claim 40 wherein

B is aryl wherein said aryl is phenyl;

- C is aryl wherein said aryl is phenyl;
 L_B is $-(CH_2)_mO(CH_2)_n-$;
 m and n are each 0;
 L_C is $-(CH_2)_pO(CH_2)_q-$;
 5 p and q are each 0;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
- 10 45. A compound according to claim 40 wherein
 B is aryl wherein said aryl is phenyl;
 C is heterocycle;
 L_B is $-(CH_2)_mO(CH_2)_n-$;
 m and n are each 0;
 15 L_C is $-(CH_2)_pO(CH_2)_q-$;
 p is 0;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
- 20 46. A compound according to claim 45 selected from the group consisting of
 N-[3-(benzyl{4-[3-(3-morpholin-4-ylpropoxy)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 1-[2-(3-{4-[(benzyl{2-methyl-3-
 25 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)ethyl]piperidine-2-
 carboxylic acid;
 N-{3-[benzyl(4-{3-[(1-methylpyrrolidin-3-yl)methoxy]phenoxy}benzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 N-{3-[(4-{3-[(1-acetylpyrrolidin-3-yl)methoxy]phenoxy}benzyl)(benzyl)amino]-2-
 30 methylphenyl}methanesulfonamide;
 N-{3-[benzyl(4-{3-[(2-oxotetrahydrofuran-3-yl)oxy]phenoxy}benzyl)amino]-2-
 methylphenyl}methanesulfonamide;
 N-[3-(benzyl{4-[3-(tetrahydrofuran-2-ylmethoxy)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 35 N-[3-(benzyl{4-[3-(tetrahydrofuran-3-ylmethoxy)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;
 N-[3-(benzyl{4-[3-(3-furylmethoxy)phenoxy]benzyl}amino)-2-
 methylphenyl]methanesulfonamide;

- N-[3-(benzyl{4-[3-(2-furylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(thien-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 5 N-[3-(benzyl{4-[3-(2-thien-3-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(thien-2-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(2-thien-2-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- 10 N-{3-[benzyl(4-{3-[2-(4-methyl-1,3-thiazol-5-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- N-{3-[benzyl(4-{3-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 15 N-[3-(benzyl{4-[3-(2-morpholin-4-ylethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N*-(3-((2,4-difluorobenzyl)(4-(3-(((2*S*,4*R*)-6-oxo-4-hydroxytetrahydro-2*H*-pyran-2-yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide;
- N*-(3-(benzyl(4-(4-(((2*S*,4*R*)-6-oxo-4-hydroxytetrahydro-2*H*-pyran-2-yl)methoxy)phenoxy)benzyl)amino)-2-methylphenyl)methanesulfonamide;
- 20 N-(3-{benzyl[4-(3-{[(2*S*)-5-oxopyrrolidin-2-yl]methoxy}phenoxy]benzyl)amino]-2-methylphenyl)methanesulfonamide;
- N-{3-[benzyl(4-{3-[(3-methyloxetan-3-yl)methoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide;
- 25 N-[3-(benzyl{4-[3-(tetrahydro-2*H*-pyran-4-yloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-[3-(benzyl{4-[3-(tetrahydrofuran-3-yloxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-(3-{benzyl[4-(3-{[(2*S*)-1-methylpyrrolidin-2-yl]methoxy}phenoxy]benzyl)amino]-2-methylphenyl)methanesulfonamide;
- 30 N-[3-(benzyl{4-[3-(pyridin-3-ylmethoxy)phenoxy]benzyl}amino)-2-methylphenyl]methanesulfonamide;
- N-{3-[benzyl(4-{3-[(2*S*)-pyrrolidin-2-ylmethoxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide; and
- 35 N-{3-[benzyl(4-{3-[(1-methylpyrrolidin-3-yl)oxy]phenoxy}benzyl)amino]-2-methylphenyl}methanesulfonamide.

47. A compound according to claim 40 wherein

B is aryl wherein said aryl is phenyl;

C is cycloalkyl;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m and n are each 0;

5 L_C is $-(CH_2)_pO(CH_2)_q-$;

p is 0;

R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;

R_4 is alkyl; and

R_6 is alkylsulfonyl.

10

48. A compound according to claim 47 selected from the group consisting of

2-[(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)methyl]cyclopropanecarboxylic acid; and

15 4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)cyclohexanecarboxylic acid.

20

49. A compound according to claim 40 wherein

B is aryl wherein said aryl is phenyl;

C is aryl wherein said aryl is phenyl;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m and n are each 0;

L_C is $-(CH_2)_pC(O)O(CH_2)_q-$;

25

R_1 , R_2 , R_3 , R_5 , X_{A1} and X_{D1} are each hydrogen;

R_4 is alkyl; and

R_6 is alkylsulfonyl.

30

50. A compound according to claim 49 that is

benzyl 3-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)propanoate.

35

51. A compound according to claim 40 wherein

B is aryl wherein said aryl is phenyl;

C is heterocycle;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m and n are each 0;

L_C is $-(CH_2)_pC(O)(CH_2)_q-$;

q is 0;
 R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen;
 R₄ is alkyl; and
 R₆ is alkylsulfonyl.

5

52. A compound according to claim 51 that is
 1-[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]-L-proline.

10

53. A compound according to claim 40 wherein
 B is aryl wherein said aryl is phenyl;
 C is heterocycle;
 L_B is -(CH₂)_mO(CH₂)_n;
 m and n are each 0;
 L_C is -O(CH₂)_pC(O)(CH₂)_q;
 q is 0;
 R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen;
 R₄ is alkyl; and
 R₆ is alkylsulfonyl.

15

20

54. A compound according to claim 53 selected from the group consisting of
 1-[5-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)pentanoyl]-L-proline;
 1-[4-(3-{4-[(benzyl{2-methyl-3-
 25 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-3-
 carboxylic acid;
 1-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-4-
 carboxylic acid;
 30 1-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]piperidine-2-
 carboxylic acid;
 1-[4-(3-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]proline; and
 35 1-[(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetyl]piperidine-4-
 carboxamide.

55. A compound according to claim 40 wherein
 B is aryl wherein said aryl is phenyl;
 C is aryl wherein said aryl is phenyl;
 L_B is $-(CH_2)_mO(CH_2)_n-$;
 L_C is $-(CH_2)_pC(O)N(R_7)(CH_2)_q-$;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
56. A compound according to claim 40 wherein
 B is aryl wherein said aryl is phenyl;
 C is heterocycle;
 L_B is $-(CH_2)_mO(CH_2)_n-$;
 m and n are each 0;
 L_C is $-(CH_2)_pC(O)N(R_7)(CH_2)_q-$;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
57. A compound according to claim 56 selected from the group consisting of
 4-(4-{4-[(2,4-difluorobenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-[3-(2-oxopyrrolidin-1-
 yl)propyl]butanamide;
 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-morpholin-4-
 ylethyl)propanamide;
 3-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-[3-(2-oxopyrrolidin-1-
 yl)propyl]propanamide;
 4-{[4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)butanoyl]amino}-1-
 methyl-1H-pyrrole-2-carboxylic acid; and
 4-(4-{4-[(benzyl{2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenyl)-N-(2-oxotetrahydrofuran-
 3-yl)butanamide.
58. A compound according to claim 40 wherein
 B is aryl wherein said aryl is phenyl;

C is heterocycle;

L_B is $-(CH_2)_mO(CH_2)_n-$;

m and n are each 0;

L_C is $-O(CH_2)_pC(O)N(R_7)(CH_2)_q-$;

5 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;


R_4 is alkyl; and

R_6 is alkylsulfonyl.

59. A compound according to claim 58 selected from the group consisting of
- 10 2-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-[3-(2-oxopyrrolidin-
1-yl)propyl]acetamide;
- 4-{[4-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)butanoyl]amino}-1-
15 methyl-1H-pyrrole-2-carboxylic acid;
- 5-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-
oxotetrahydrofuran-3-yl)pentanamide;
- 2-(4-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-
20 oxotetrahydrofuran-3-yl)acetamide;
- 4-(3-{4-[(benzyl{2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(2-
oxotetrahydrofuran-3-yl)butanamide;
- 25 2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-[3-(2-oxopyrrolidin-
1-yl)propyl]acetamide;
- 2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(1,3-thiazol-5-
30 ylmethyl)acetamide; and
- 2-(2-chloro-5-{4-[(2,4-difluorobenzyl){2-methyl-3-
[(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)-N-(3-morpholin-4-
ylpropyl)acetamide.

- 35 60. A compound according to claim 40 wherein
- B is aryl wherein said aryl is phenyl;
- C is aryl wherein said aryl is phenyl;
- L_B is $-(CH_2)_mC(O)(CH_2)_n-$;

- m and n are each 0;
L_C is -(CH₂)_pO(CH₂)_q;
p is 0;
R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen;
5 R₄ is alkyl; and
R₆ is alkylsulfonyl.
61. A compound according to claim 60 selected from the group consisting of
N-[3-(benzyl{4-[3-(2-phenylethoxy)benzoyl]benzyl}amino)-2-
10 methylphenyl]methanesulfonamide; and
N-[3-(benzyl{4-[3-(benzyloxy)benzoyl]benzyl}amino)-2-
methylphenyl]methanesulfonamide.
62. A compound according to claim 40 wherein
15 B is aryl wherein said aryl is phenyl;
C is heterocycle;
L_B is -(CH₂)_mC(O)(CH₂)_n;
m and n are each 0;
L_C is -(CH₂)_pO(CH₂)_q;
20 p is 0;
R₁, R₂, R₃, R₅, X_{A1} and X_{D1} are each hydrogen;
R₄ is alkyl; and
R₆ is alkylsulfonyl.
63. A compound according to claim 62 selected from the group consisting of
N-[3-(benzyl{4-[3-(tetrahydrofuran-2-ylmethoxy)benzoyl]benzyl}amino)-2-
methylphenyl]methanesulfonamide;
N-[3-(benzyl{4-[3-(tetrahydrofuran-3-ylmethoxy)benzoyl]benzyl}amino)-2-
methylphenyl]methanesulfonamide;
30 N-[3-[benzyl(4-[3-[2-(2-oxopyrrolidin-1-yl)ethoxy]benzoyl]benzyl)amino]-2-
methylphenyl]methanesulfonamide; and
N-[3-[benzyl(4-[3-[2-(4-methyl-1,3-thiazol-5-yl)ethoxy]benzoyl]benzyl)amino]-2-
methylphenyl]methanesulfonamide.
64. A compound according to claim 1 of formula (V)
- 35



(V),

or a pharmaceutically acceptable salt or prodrug thereof.

65. A compound according to claim 64 wherein
 B is aryl wherein said aryl is phenyl;
 E is aryl wherein said aryl is phenyl;
 L_B is $-(CH_2)_mO(CH_2)_n-$;
 m and n are each 0;
 L_E is $-(CH_2)_rC(O)(CH_2)_s-$
 r and s are each 0;
 $R_1, R_2, R_3, R_5, X_{A1}$ and X_{D1} are each hydrogen;
 R_4 is alkyl; and
 R_6 is alkylsulfonyl.
66. A compound according to claim 65 selected from the group consisting of
 (3-{4-[(4-benzoylbenzyl){2-methyl-3-
 [(methylsulfonyl)amino]phenyl}amino)methyl]phenoxy}phenoxy)acetic acid; and
 (5-{4-[(4-benzoylbenzyl)(2-methyl-3-
 [(methylsulfonyl)amino]phenyl)amino)methyl]phenoxy}-2-chlorophenoxy)acetic acid.
67. A pharmaceutical composition comprising a therapeutically effective amount of a
 compound of claim 1 in combination with a pharmaceutically acceptable carrier.
68. A method of selectively modulating the effects of the glucocorticoid receptor in a
 mammal comprising administering an effective amount of a compound of claim 1.
69. A method of treating type II diabetes in a mammal comprising administering a
 therapeutically effective amount of a glucocorticoid receptor antagonist.

70. A method of treating type II diabetes in a mammal comprising administering a therapeutically effective amount of a compound of claim 1.
- 5 71. A method of treating symptoms related to type II diabetes wherein said symptoms are selected from the group consisting of hyperglycemia, hyperinsulinemia, inadequate, glucose clearance, obesity, hypertension and high glucocorticoid levels in a mammal comprising administering a therapeutically effective amount of a compound of claim 1.
- 10 72. A method of treating diseases associated with an excess or deficiency of glucocorticoids, said diseases selected from the group consisting of diabetes, obesity, Syndrome X, Cushing's Syndrome, Addison's disease, inflammatory diseases such as asthma, rhinitis and arthritis, allergy, autoimmune disease, immunodeficiency, anorexia, cachexia, bone loss or bone frailty, and wound healing comprising administering a
15 therapeutically effective amount of a compound of claim 1.
73. A method of treating a disease, comprising administering a compound of claim 1 in combination with an anti-diabetic agent, wherein said disease is selected from the group consisting of diabetes and Syndrom X.
- 20 74. A method according to claim 73, wherein said anti-diabetic agent is selected from the group consisting of insulin, mecasemin, nateglinide, metformin, chlorpropamide, glipizide, glyburide, troglitazone, pioglitazone, rosiglitazone, acarbose, voglibose, miglitol, zopolrestat and repaglinide.
- 25 75. A method of treating obesity, comprising administering a compound of claim 1 in combination with an anti-obesity agent.
- 30 76. A method according to claim 75, wherein said anti-obesity agent is selected from the group consisting of orlistat, sibutramine, dexfenfluramine, bromocryptine, phentermine, phendimetrazine and mazindol.

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/US 02/04501

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C307/04 C07C311/08 A61K31/18 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 929 058 A (DEISHER THERESA A) 27 July 1999 (1999-07-27) cited in the application the whole document	1
A	US 6 166 013 A (COGHLAN MICHAEL J ET AL) 26 December 2000 (2000-12-26) the whole document	1

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

12 June 2002

Date of mailing of the international search report

24/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Goetz, G

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 02/04501

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 68 - 76 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

Inter nal Application No
PCT/US 02/04501

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
US 5929058	A	27-07-1999	AU	5729098 A	17-07-1998
			WO	9827986 A1	02-07-1998
<hr/>					
US 6166013	A	26-12-2000	NONE		
<hr/>					

THIS PAGE BLANK (USPTO)